arXiv:0704.1121v2 [math.OA] 27 Apr 2007

CLASSIFICATION OF NONCOMMUTING QUADRILATERALS OF FACTORS

PINHAS GROSSMAN AND MASAKI IZUMI

Abstract. A quadrilateral of factors is an irreducible inclusion of factors $N \subset M$ with intermediate subfactors P and Q such that P and Q generate M and the intersection of P and Q is N. We investigate the structure of a noncommuting quadrilateral of factors with all the elementary inclusions $P \subset M$, $Q \subset M$, $N \subset P$, and $N \subset Q$ 2-supertransitive. In particular we classify noncommuting quadrilaterals with the indices of the elementary subfactors less than or equal to 4. We also compute the angles between P and Q for quadrilaterals coming from α-induction and asymptotic inclusions.

1. Introduction

If M is a factor with a finite group G acting by outer automorphisms, then $N = M^G$, the algebra of fixed points of the action, is an irreducible subfactor of M with index equal to the order of G. It is well known that in this case the intermediate subalgebras $N \subset P \subset M$ are precisely the fixed point algebras of the subgroups of G [48].

In this spirit one thinks of the classification of intermediate subfactors as a "noncommutative Galois theory". Motivated by von Neumann's study of projection lattices in Hilbert space as a "continuous geometry", Watatani proposed studying lattices of intermediate subfactors, which are finite for irreducible finite-index inclusions, as a quantization [56]. Sano and Watatani introduced the notion of angles between subfactors, a numerical invariant which measures the degree of noncommutativity of pairs of subfactors [54].

A major step towards the classification of intermediate subfactors was Bisch's characterization of intermediate subfactors as *biprojections* in planar algebras- elements of the standard invariant which are projections in both of the dual algebraic structures [4]. Bisch and Jones then described a generic construction of intermediate subfactors by constructing a universal planar algebra generated by a single biprojection, the Fuss-Catalan algebra with parameters corresponding to the indices [7]. These intermediate subfactors are *supertransitive* in the sense that the standard invariants are minimal.

It was hoped that a similar generic planar algebra for multiple intermediate subfactors could be described in terms of indices and invariants such as angles. Indeed, a tensor product does yield a generic construction of commuting pairs of intermediate subfactors, but constructing pairs with nontrivial angles has proven more difficult.

Work supported by JSPS.

In [13], Jones and the first-named author showed that there are essentially only 2 examples of noncommuting irreducible quadrilaterals of factors such that all of the elementary inclusions are supertransitive. One is a quadrilateral of subgroups of the symmetric group $\mathfrak{S}_3$ and the other comes from the GHJ family of subfactors and all the elementary inclusions have index $2+\sqrt{2}$.

It is known that the structure of an intermediate subfactor $N \subset P \subset M$ is determined by the relationship between the two systems of P-P bimodules arising from $N \subset P$ and $P \subset M$ (see [6],[26]). After [13] appeared, the second-named author gave a short proof of the above mentioned result using simple computation of $P-P$ bimodules (see Remark 4.2). It turns out from the proof that the assumption in [13] is too restrictive from the view point of bimodules. Therefore to capture more interesting and general structure, one should relax the assumption of supertransitivity. Indeed, the first named author [12] recently obtained a complete classification result of irreducible non-commuting quadrilaterals with $N \subset P$ and $N \subset Q$ isomorphic to the Jones subfactor of index less than 4 (without posing any assumption on $P \subset M$ or $Q \subset M$) and showed that a series of such quadrilaterals exists.

In this paper we investigate quadrilaterals whose elementary inclusions $P \subset M$, $Q \subset M$, $N \subset P$, and $N \subset Q$ are each 2-supertransitive (an inclusion $L \subset K$ is 2-supertransitive if the complement of L in the $L-L$ bimodule decomposition of K is irreducible.) In this case the systems of $P-P$ bimodules for $N \subset P$ and for $P \subset M$ are each generated by a single irreducible $P-P$ bimodule, and the question is how these two bimodules are related.

A key tool in this analysis is the notion of second cohomology for subfactors, introduced by the second named author and Kosaki in [27], which counts the inner conjugacy classes of subfactors sharing the same basic extension (as a bimodule class). For a quadrilateral with $N \subset P$ and $N \subset Q$ 2-supertransitive, noncommutativity is equivalent to the existence of an $N-N$ bimodule isomorphism from P to Q [13]. If $N \subset P$ (dual) has trivial second cohomology, this $N-N$ bimodule isomorphism can actually be realized as an algebra isomorhism. In particular they showed that any 3-supertransitive subfactor has trivial second cohomology.

Ultimately this leads to a classification of noncommuting quadrilaterals whose elementary subfactors are 2-supertransitive with trivial second cohomology into two basic types: those which are not cocommuting ("cocommuting" means that the quadrilateral of commutants is a commuting square) and have $[M:P]=[P:N]$, like the $(2+\sqrt{2})^2$ example, and those which are cocommuting and have $[M:P]=[P:N]-1$, like the $\mathfrak{S}_3$ example. The latter type is further subdivided according to the Galois group of $N \subset M$, which must be a subgroup of $\mathfrak{S}_3$. In the case that the elementary subfactors have index less than or equal to 4 we examine all the possibilities and arrive at the following result.

Theorem 1.1. *There are exactly seven noncommuting irreducible quadrilaterals of hyperfinite II_1 factors whose elementary inclusions all have indices less than or equal to 4, up to conjugacy.*

If $(G_{N\subset P}, G_{P\subset M})$ are the principal graphs of the elementary subfactors, the possible configurations are:

$$(A_7, A_7), \quad (E_7^{(1)}, E_7^{(1)})$$
$$(A_5, A_3), \quad (D_6, A_4), \quad (E_7^{(1)}, A_5), \quad (E_6^{(1)}, D_4)$$
$$(D_6^{(1)}, A_3)$$

The fully supertransitive cases are (A_7, A_7), which is the $(2+\sqrt{2})^2$ example, and (A_5, A_3), which is the $\mathfrak{S}_3$ example, and all other cases have some extra structure. The new examples include a noncocommuting quadrilateral whose elementary subfactors have index 4 $(E_7^{(1)}, E_7^{(1)})$, a cocommuting quadrilateral whose indices are $(3+\sqrt{5})/2$ and $(5+\sqrt{5})/2$ (D_6, A_4), and several cocommuting quadrilaterals arising from group actions. Note that any quadrilateral is either noncommuting, noncocommuting (and hence dual to a noncommuting quadrilateral), or else is both commuting and cocommuting, in which case there is no obstruction to the choices of $N \subset P$ and $N \subset Q$.

It is also interesting to study such quadrilaterals with larger indices. In fact the exotic Haagerup subfactor appears as an elementary subfactor with maximal supertransitivity in both types of quadrilaterals and the Asaeda-Haagerup subfactor may appear as well. It is still unknown how many examples there are of the various types of quadrilaterals (although there is an infinite series of cocommuting quadrilaterals with 3-supertransitive elementary subfactors, coming from symmetric group actions.)

Second cohomolgy also turns out to be closely related to angles. For a noncommuting quadrilateral with $N \subset P$ and $N \subset Q$ 2-supertransitive, there are at most two possibilities for the angle for each cohomolgy class of $N \subset P$ (dual). In particular, if $N \subset P$ is 3-supertransitive, the angle is always $\cos^{-1} 1/([P:N]-1)$. We also compute the angles in quadrilaterals arising from α-induction (which are identified with the GHJ pairs of [13]) and asymptotic inclusions, the former of which includes the forked Temperley-Lieb quadrilaterals of [12].

The paper is organized as follows:

Section 2 sets forth the basic notation of quadrilaterals, sectors and Q-systems which are used throughout. We often deal with infinite factors since we are actually studying properties of the standard invariant.

Section 3 studies the relationship between second cohomology for subfactors and angles.

Section 4 proves that quadrilaterals whose elementary subfactors are 2-supertransitive with trivial second cohomology fall into the two basic types discussed above, and that for the cocommuting type the Galois group of the total inclusion is a subgroup of $\mathfrak{S}_3$.

Section 5 analyzes several classes of quadrilaterals according to the results of the previous section and for each class provides an example with maximal supertransitivity.

Section 6 classifies all noncommuting irreducible quadrilaterals whose elementary subfactors have indices less than or equal to 4.

Section 7 computes the angles for quadrilaterals coming from α-induction and asymptotic inclusions.

The Appendix constructs a Q-system from a sector in the Haagerup principal graph and shows that it is unique, filling in a gap for one of the examples in Section 5.

2. Preliminaries

We first fix notation used throughout this paper. We always assume that Hilbert spaces are separable and von Neumann algebras have separable preduals.

2.1. **Quadrilaterals.** For an inclusion of II_1 factors $N \subset M$, the Jones index, the Jones projection, and the trace preserving conditional expectation onto N are denoted by $[M : N]$, e_N and E_N respectively [29]. We use the same symbols for an inclusion of properly infinite factors $N \subset M$ considering a unique minimum conditional expectation [38], [15]. Although the Jones index $[M : N]$ does not necessarily coincides with the minimum index for a II_1 inclusion in general, no confusion arises as we always consider extremal inclusions throughout this paper. For a II_1 factor, we denote by tr the unique normalize trace. For an inclusion of general factors $N \subset M$ of finite index, we use the same symbol tr for the restriction of E_N to the relative commutant $M \cap N'$.

Let $N \subset M$ be an inclusion of factors of finite index with associated tower

$$N = M_{-1} \subset M = M_0 \subset M_1 \subset M_2 \subset \cdots,$$

where M_{k+1}, $k \geq 0$ is the von Neumann algebra on the standard space of M_k generated by M_k and the Jones projection $e_{k+1} = e_{M_{k-1}}$. Each e_k commutes with N, so $\{1, e_1, .., e_k\}$ generates a *-subalgebra, which we will call TL_{k+1}, of the k^{th} relative commutant $N' \cap M_k$. The following definition first appeared in [32].

Definition 2.1. Call a finite-index subfactor $N \subset M$ *k-supertransitive* (for $k > 1$) if $N' \cap M_{k-1} = TL_k$. We will say $N \subset M$ is *supertransitive* if it is k-supertransitive for all k.

Note that $N \subset M$ is k-supertransitive if and only if $M \subset M_1$ is k-supertransitive. Sano and Watatani [54] introduced the notion of angles between two subfactors.

Definition 2.2. For two subfactors P and Q of a factor M, the *set of angles* $\mathrm{Ang}(P, Q)$ between P and Q is the spectrum of the angle operator of the two Jones projections e_P and e_Q, that is, the spectrum of $\cos^{-1} \sqrt{e_P e_Q e_P - e_P \wedge e_Q}$ where $e_P e_Q e_P - e_P \wedge e_Q$ is regarded as the operator acting on its support.

Note that $\mathrm{Ang}(P, Q) = \mathrm{Ang}(Q, P)$ always holds.

Definition 2.3. A *quadrilateral* of factors $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ is an inclusion of factors $N \subset M$ of finite index with two intermediate subfactors $P \neq Q$ such that P and Q generate M and $P \cap Q = N$. The subfactors $N \subset P$, $N \subset Q$, $P \subset M$, and $Q \subset M$ are said to be the elementary subfactors for $\mathfrak{Q}$. When $M \cap N' = \mathbb{C}$, the quadrilateral $\mathfrak{Q}$ is said to be *irreducible.* When E_P commutes with E_Q, the quadrilateral $\mathfrak{Q}$ is said to be *commuting.* Let $M_1 = J_M N' J_M$, $\hat{P} = J_M P' J_M$, and $\hat{Q} = J_M Q' J_M$ be the basic extensions of M by N, P, and Q respectively. Then $\hat{P}$ and $\hat{Q}$ are intermediate subfactors between $M \subset M_1$, which form the *dual quadrilateral* $\hat{\mathfrak{Q}} = \begin{array}{ccc} \hat{Q} & \subset & M_1 \\ \cup & & \cup \\ N & \subset & \hat{P} \end{array}$.

The quadrilateral $\mathfrak{Q}$ is said to be *cocommuting* if $\hat{\mathfrak{Q}}$ is commuting. The quadrilateral $\mathfrak{Q}$ is said to be (p,q)-*supertransitive* if $P \subset M$ and $Q \subset M$ are p-supertransitive and $N \subset P$ and $N \subset Q$ are q-supertransitive. Two quadrilaterals $\mathfrak{Q} = \begin{matrix} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{matrix}$ and $\tilde{\mathfrak{Q}} = \begin{matrix} \tilde{P} & \subset & \tilde{M} \\ \cup & & \cup \\ \tilde{N} & \subset & \tilde{Q} \end{matrix}$ are said to be *conjugate* if there exists an isomorphism π from M onto $\tilde{M}$ such that $\pi(P) = \tilde{P}$ and $\pi(Q) = \tilde{Q}$. The two quadrilateral $\mathfrak{Q}$ and $\tilde{\mathfrak{Q}}$ are said to be *flip conjugate* if either they are conjugate or there exists an isomorphism π from M onto $\tilde{M}$ such that $\pi(P) = \tilde{Q}$ and $\pi(Q) = \tilde{P}$.

We denote by $\widehat{P \subset M}$ the inclusion $M \subset \hat{P}$ and by $\widecheck{N \subset P}$ the inclusion $\check{P} \subset N$, where $\check{P} \subset N \subset P$ is the downward basic construction. Note that $\check{P}$ is uniquely determined up to inner conjugacy in N.

When $\mathfrak{Q} = \begin{matrix} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{matrix}$ is a non-commuting quadrilateral of subfactors such that $N \subset P$ and $N \subset Q$ are 2-supertransitive, the two projections e_P and e_Q are of rank 2 in $M_1 \cap N'$ and $e_P \wedge e_Q = e_N$. Thus $\mathrm{Ang}(P,Q)$ consists of at most one point, which will be denoted by $\Theta(P,Q)$ if $\mathfrak{Q}$ is non-commuting.

The following lemma is essentially proved in [13, Proposition 3.2.11, Theorem 3.3.4] using Landau's result [41]:

Lemma 2.4. *Let* $\mathfrak{Q} = \begin{matrix} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{matrix}$ *be an irreducible cocommuting quadrilaterals of subfactors such that* $N \subset P$ *and* $N \subset Q$ *are 2-supertransitive. If* $[M:P] = [P:N]$, *the quadrilateral* $\mathfrak{Q}$ *is commuting, and if* $[M:P] \neq [P:N]$,

$$\cos^2 \Theta(P,Q) = \frac{[P:N] - [M:P]}{[M:P]([P:N]-1)}.$$

We show an easy example of a noncommuting quadrilateral coming from a finite group action. For an automorphism group G of a factor R, we denote by R^G the fixed point subalgebra of R under G. The next lemma follows from Lemma 2.4.

Lemma 2.5. *Let* G *be a finite group and* H *and* K *be subgroups of* G *such that the natural actions of* G *on* G/H *and* G/K *are 2-transitive. Assume that* $[G:H] = [G:K]$, $[G:H] \neq [H:H \cap K]$, *and* G *acts on a factor* R *as an outer automorphism group. We set* $M = R^{K \cap H}$, $P = R^H$, $Q = R^K$, *and* $N = R^G$. *Then* $\mathfrak{Q} = \begin{matrix} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{matrix}$ *is an irreducible noncommuting and cocommuting quadrilateral of factors such that* $N \subset P$ *and* $N \subset Q$ *are 2-supertransitive. The angle between* P *and* Q *is given by*

$$\cos^2 \Theta(P,Q) = \frac{[G:H] - [H:H \cap K]}{[H:H \cap K]([G:H]-1)}.$$

Example 2.6. For a finite set F, we denote by $\mathfrak{S}_F$ the set of permutations of F and by $\mathfrak{A}_F$ the set of even permutations of F. Let $F_0 = \{1, 2, \cdots, n\}$, $F_1 = \{1, 2, \cdots, n-1\}$, and $F_2 = \{2, 3, \cdots, n\}$. Then $G = \mathfrak{S}_{F_0}$ ($G = \mathfrak{A}_{F_0}$), $H = \mathfrak{S}_{F_1}$ ($H = \mathfrak{A}_{F_1}$), and $K = \mathfrak{S}_{F_2}$ ($K = \mathfrak{A}_{F_2}$) with $n \geq 3$ ($n \geq 4$) satisfies the assumption of Lemma 2.5 and $\Theta(P, Q) = \cos^{-1}(1/(n-1))$.

In [13] Jones and the first-named author obtained the following theorem:

Theorem 2.7. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *be an irreducible noncommuting quadrilateral of factors. If $\mathfrak{Q}$ is (6,6)-supertransitive, one of the following two occurs:*

1. *All the elementary subfactors of $\mathfrak{Q}$ are the A_7 subfactors and $\Theta(P, Q) = \cos^{-1}(\sqrt{2} - 1)$. When M is the hyperfinite II_1 factor, such a quadrilateral exists and is unique up to conjugacy.*
2. *$[M : P] = [M : Q] = 2$, $[P : N] = [Q : N] = 3$, $\Theta(P, Q) = \pi/3$, and the principal graphs of $N \subset P$ and $N \subset Q$ are A_5. In this case, there exists an outer action of the symmetric group $\mathfrak{S}_3$ of degree 3 on M such that N is the fixed point algebra of the action. When M is the hyperfinite II_1 factor, such a quadrilateral is unique up to conjugacy.*

In Section 4, we give a new proof of the above theorem, except for the existence, relaxing the assumption. The theorem still holds if we assume that $\mathfrak{Q}$ is (3,4)-supertransitive.

2.2. **Sectors.**

To compute the angle between P and Q, it is more convenient to work with properly infinite factors using sectors as we will see below. Thus we recall a few basic facts about sectors here. The reader is referred to [20] for details. Note that the structure of the intermediate subfactor lattice of an inclusion $N \subset M$, including information of the angle, only depends on the standard invariant (or paragroup, plannar algebra) of $N \subset M$. Every standard invariant realized in the type II_1 case is also realized in the properly infinite case and vice versa.

For the reason stated above, we always assume in the proofs that the subfactors involved are properly infinite though we state results for general quadrilaterals of factors. When we show uniqueness results, we need to deal with quadrilaterals of hyperfinite II_1 factors. Since the classification theory of subfactors for the hyperfinite II_1 factor and the hyperfinite II_∞ factor is the same as far as strongly amenable subfactors are concerned [52], we may assume in the proofs that the factors involved are isomorphic to the hyperfinite II_∞ factors in this case.

Remark 2.8. When one discusses sectors, it is customary to assume that every factor involved is of type III. However the whole theory also works for general properly infinite factors. This is based on the following facts: (1) for an inclusion of properly infinite factors $N \subset M$, every non-zero projection in the relative commutant $N' \cap M$ is an infinite projection in M, and (2) a properly infinite factor has a unique representation whose commutant is also properly infinite.

Let M and N be properly infinite factors. We denote by $\mathrm{Mor}(N,M)$ the set of normal unital homomorphisms from N to M. For two morphisms $\rho,\sigma\in\mathrm{Mor}(N,M)$, we denote by (ρ,σ) the set of intertwiners between ρ and σ

$$(\rho,\sigma)=\{v\in M;\ v\rho(x)=\sigma(x)v,\ \forall x\in N\}.$$

When ρ is irreducible, that is $M\cap\rho(N)'=\mathbb{C}$, the space (ρ,σ) is a Hilbert space with an inner product $\langle v_1,v_2\rangle=v_2^*v_1$ for $v_1,v_2\in(\rho,\sigma)$. The two morphisms ρ and σ are said to be equivalent if there exists a unitary $u\in(\rho,\sigma)$. We denote by $\mathrm{Sect}(M,N)$ the quotient of $\mathrm{Mor}(N,M)$ by this equivalence relation. We denote by $[\rho]$ the equivalence class of ρ, which we call a sector. However, we often omit the quare bracket when no confusion arises. When $M=N$, we use the notation $\mathrm{Mor}(M,M)=\mathrm{End}(M)$ and $\mathrm{Sect}(M,M)=\mathrm{Sect}(M)$.

For $M-N$ bimodule X and $\rho\in\mathrm{Mor}(L,N)$, we denote by X_ρ the $M-L$ bimodule defined by

$$x\cdot\xi\cdot y=x\xi\rho(y),\quad \xi\in X,\ x\in M,\ y\in L.$$

For $\sigma\in\mathrm{Mor}(L,M)$, we denote by ${}_\sigma X$ the $L-N$ bimodule similarly defined.

Let $L^2(M)$ be the standard Hilbert space of M, which is a $M-M$ bimodule with $x\xi y=xJ_My^*J_M\xi$, where J_M is the modular conjugation. Then it is known that for every $M-N$ bimodule X, there exists a unique sector $[\rho]\in\mathrm{Sect}(M,N)$ such that X is equivalent to $L^2(M)_\rho$. With this correspondence, the relative tensor product of two bimodules is transformed into the composition of two morphisms. A direct sum of bimodules is easily translated into the sector language too (see [20]).

We denote by $\mathrm{Mor}_0(N,M)$ the set of $\rho\in\mathrm{Mor}(N,M)$ whose image has finite index. For $\mathrm{Mor}_0(N,M)$, we denote by $d(\rho)$ the square root of $[M:\rho(N)]$, called the (statistical) dimension of ρ, which is multiplicative under composition of two sectors and is additive under direct sum. We use the notation $\mathrm{Sect}_0(M,N)$ etc. in an obvious sense.

Corresponding to the complex conjugate bimodule of $L^2(M)_\rho$, the conjugate sector of ρ is defined. For $\rho\in\mathrm{Mor}_0(N,M)$, the (equivalence class of) conjugate morphism $\bar\rho\in\mathrm{Mor}_0(M,N)$ is characterized by existence of two isometries $r_\rho\in(\mathrm{id}_N,\bar\rho\rho)$ and $\bar r_\rho\in(\mathrm{id}_M,\rho\bar\rho)$ satisfying the following equation

$$r_\rho^*\bar\rho(\bar r_\rho)=\frac{1}{d(\rho)},\quad \bar r_\rho^*\rho(r_\rho)=\frac{1}{d(\rho)}.$$

Throughout this paper, we fix such r_ρ and $\bar r_\rho$ for each ρ. When ρ is not self-conjugate, we can and do assume $r_{\bar\rho}=\bar r_\rho$. This is not the case for a self-conjugate ρ. In this case $r_\rho^*\rho(r_\rho)=\pm1/d(\rho)$ holds and so $r_\rho=\pm\bar r_\rho$ [42]. A self-conjugate endomorphism ρ is said to be real if $r_{\bar\rho}=\bar r_\rho$ and said to be pseudo-real if $r_{\bar\rho}=-\bar r_\rho$.

We set $\phi_\rho(x)=r_\rho^*\bar\rho(x)r_\rho$ for $x\in N$, which is called the standard left inverse of ρ. We denote by E_ρ the minimum conditional expectation from M onto the image of ρ. Then we have $E_\rho=\rho\phi_\rho$.

For an inclusion of properly infinite factors $N\subset M$ of finite index, we denote by $\iota_{M,N}$ the inclusion map $\iota_{M,N}:N\hookrightarrow M$. Then the $M-N$ bimodule $X={}_ML^2(M)_N$ is nothing but $L^2(M)\iota_{M,N}$. Since ${}_ML^2(M_1)_M$ is isomorphic to $X\otimes_N X^*$, we have ${}_ML^2(M_1)_M\cong L^2(M)_{\iota_{M,N}\overline{\iota_{M,N}}}$. The endomorphism $\iota_{M,N}\overline{\iota_{M,N}}$ of M is called the canonical endomorphism for $N \subset M$. The endomorphism $\overline{\iota_{M,N}}\iota_{M,N}$ of N is called the dual canonical endomorphism, which is the canonical endomorphism for the inclusion $\overline{\iota_{M,N}}(M) \subset N$.

We often use diagrams for computation of intertwiners. An intertwiner $s \in (\rho, \sigma\tau)$ is expressed by the diagram



For $t \in (\sigma, \mu\nu)$ and $u \in (\tau, \xi\eta)$, the products $ts \in (\rho, \mu\nu\tau)$ and $\sigma(u)s \in (\rho, \sigma\xi\eta)$ are expressed as



For special operators $1 \in (\rho, \rho)$, $\sqrt{d(\rho)}r_\rho \in (\mathrm{id}, \bar{\rho}\rho)$, and $\sqrt{d(\rho)}\bar{r}_\rho^* \in (\mathrm{id}, \rho\bar{\rho})$, we use the following diagrammatic expressions:



Then the equations $d(\rho)\bar{r}_\rho^*\rho(r_\rho) = 1$ and $d(\rho)\bar{\rho}(\bar{r}_\rho^*)r_\rho = 1$ are expressed as



The following lemma holds (see [28, Section 3]):

Theorem 2.9. *Let $N \subset M$ be an irreducible inclusion of properly infinite factors with finite index and let*

$$[\overline{\iota_{M,N}}\iota_{M,N}] = \bigoplus_{i=0}^{m} n_i[\rho_i]$$

be the irreducible decomposition, where n_i is the multiplicity of ρ_i in $\overline{\iota_{M,N}}\iota_{M,N}$. Then

(1) *$n_i \leq d(\rho_i)$.*
(2) *Let $\mathcal{H}_i = (\iota_{M,N}, \iota_{M,N}\rho_i)$. For $s_1, s_2 \in \mathcal{H}_i$,*

$$E_N(s_1 s_2^*) = \frac{\langle s_1, s_2 \rangle}{d(\rho_i)}.$$

(3) $\dim \mathcal{H}_i = n_i$.

(4) *Let* $\{s(i)_j\}_{j=1}^{n_i}$ *be an orthonormal basis of* $\mathcal{H}_i$. *Then* $x \in M$ *has the following unique expansion*

$$x = \sum_{i=0}^{m} \sum_{j=1}^{n_i} s(i)_j^* x(i)_j$$

with coefficients $x(i)_j \in N$. *The coefficients* $x(i)_j$ *is given by*

$$x(i)_j = d(\rho_i) E_N(s(i)_j x).$$

The fourth statement above says that $\{\sqrt{d(\rho_i)}s(i)_j^*\}_{ij}$ is a Pimsner-Popa basis [51]. The advantage of working on properly infinite factors is that we can choose each element of the basis from an intertwiner space, and so (4) is considered as a crossed product type decomposition.

Let A_i be the linear span of $\{s^* e_N t;\ s, t \in \mathcal{H}_i\}$. Then [28, Section 3] shows that A_i is a simple component of $M \cap N'$ and

$$M_1 \cap N' = \bigoplus_{i=0}^{m} A_i.$$

We can introduce a right action of A_i on $\mathcal{H}_i$ by

$$v \cdot (s^* e_N t) = \frac{1}{d(\rho_i)} \langle v, s \rangle t,$$

which gives a $*$-isomorphism from the opposite algebra of A_i onto $\mathbb{B}(\mathcal{H}_i)$. Let $e(i)_{jk} = d(\rho_i) s(i)_j^* e_N s(i)_k$. Then $\{e(i)_{jk}\}_{jk}$ is a system of matrix unit of A_i. Let P be an intermediate subfactor and let $m_i = \dim(\iota_{P,N}, \iota_{P,N}\rho_i)$. Then we may and do assume that $\{s(i)_j\}_{j=1}^{m_i}$ is an orthonormal basis for $\mathcal{H}_i \cap P = (\iota_{P,N}, \iota_{P,N}\rho_i)$.

Lemma 2.10. *Let the notation be as above. Then*

(1) *The restriction* $E_P|_{\mathcal{H}_i}$ *of* E_P *to* $\mathcal{H}_i$ *is the projection onto* $\mathcal{H}_i \cap P = (\iota_{P,N}, \iota_{P,N}\rho_i)$.

(2) *The Jones projection* e_P *is expressed as*

$$e_P = \sum_{i=0}^{m} \sum_{j=1}^{m_i} e(i)_{jj}.$$

(3) *Let* z_i *be the unit of* A_i. *Then*

$$s \cdot z_i e_P = E_P(s), \quad \forall s \in \mathcal{H}_i.$$

Proof. (1) Note that $E_P(\mathcal{H}_i) = \mathcal{H}_i \cap P = (\iota_{P,N}, \iota_{P,N}\rho_i)$ holds and $E_P|_{\mathcal{H}_i}$ is an idempotent. Thus it suffices to show that $E_P|_{\mathcal{H}_i}$ is self-adjoint. For $s, t \in \mathcal{H}_i$, we have

$$\begin{aligned} \langle E_P(s), t \rangle &= d(\rho_i) E_N(E_P(s) t^*) = d(\rho_i) E_N(E_P(E_P(s) t^*)) \\ &= d(\rho_i) E_N(E_P(s) E_P(t)^*)) = \langle E_P(s), E_P(t) \rangle, \end{aligned}$$

which shows the statement.

(2) Let φ be a faithful normal state of M satisfying $\varphi \cdot E_N = \varphi$. Then (1) and Theorem 2.9,(4) imply the following for $x \in M$ and the GNS cyclic and separating

vector Ω for φ:

$$\begin{aligned} e_Px\Omega &= E_P(\sum_{i=0}^{m}\sum_{j=1}^{n_i} t(i)_j^*x(i)_j)\Omega = \sum_{i=0}^{m}\sum_{j=1}^{n_i} E_P(t(i)_j)^*x(i)_j\Omega \\ &= \sum_{i=0}^{m}\sum_{j=1}^{m_i} t(i)_j^*x(i)_j\Omega = \sum_{i=0}^{m} d(\rho_i)\sum_{j=1}^{m_i} t(i)_j^*E_N(t(i)_jx)\Omega \\ &= \sum_{i=0}^{m} d(\rho_i)\sum_{j=1}^{m_i} t(i)_j^*e_Nt(i)_jx\Omega. \end{aligned}$$

This shows (2). (3) follows from (2). □

Remark 2.11. The above lemma gives a powerful method to compute the angle between subfactors. Let Q be another intermediate subfactor of $N \subset M$. To compute the angle between P and Q, it suffices to compute the angle between two subspaces $(\iota_{P,N}, \iota_{P,N}\rho_i)$ and $(\iota_{Q,N}, \iota_{Q,N}\rho_i)$ of $(\iota_{M,N}, \iota_{M,N}\rho_i)$ for those ρ_i contained in $\overline{\iota_{P,N}}\iota_{P,N}$ and $\overline{\iota_{Q,N}}\iota_{Q,N}$. Equivalently, it suffices to compute the spectrum of the restriction of E_PE_Q to $(\iota_{P,N}, \iota_{P,N}\rho_i)$.

Remark 2.12. Even when the inclusion $N \subset M$ is reducible, the angle $\mathrm{Ang}(P,Q)$ can be computed from $E_PE_Q|_{(\iota_{P,N},\iota_{P,N}\rho_i)}$ as long as $N \subset P$ and $N \subset Q$ are irreducible. Indeed we have $M = P \oplus \ker E_P$ and

$$P = \bigoplus_{i=0}^{m} N(\iota_{P,N}, \iota_{P,N}\rho_i).$$

Since $E_PE_QE_P$ preserves $(\iota_{P,N}, \iota_{P,N}\rho_i)$, it suffice to compute $E_PE_Q|_{(\iota_{P,N},\iota_{P,N}\rho_i)}$ to obtain the eigenvalues of $E_PE_QE_P$.

3. Q-SYSTEMS AND ANGLES

When $\begin{matrix} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{matrix}$ is an irreducible noncommuting quadrilateral of subfactors and $N \subset P$ and $N \subset Q$ are 2-supertransitive, it is observed in [13, Lemma 4.2.1] that $L^2(P)$ and $L^2(Q)$ are equivalent as $N-N$ bimodules, or in other words, $\overline{\iota_{P,N}}\iota_{P,N}$ is equivalent to $\overline{\iota_{Q,N}}\iota_{Q,N}$. Consequently, $[P:N]=[Q:N]$ and so $[M:P]=[M:Q]$. This follows from the fact that the two projections e_P and e_Q are equivalent in $M_1 \cap N'$, which is identified with $\mathrm{End}_N L^2(M)_N$. Let $\check{P}$ and $\check{Q}$ be subfactors of N such that $\check{P} \subset N \subset P$ and $\check{Q} \subset N \subset P$ are towers. It is known that under a certain situation, such as 3-supertransitivity of $N \subset P$ and $N \subset Q$, the equivalence ${}_NL^2(P)_N \cong {}_NL^2(Q)_N$ implies inner conjugacy of $\check{P}$ and $\check{Q}$ in N [27], [35] though it is not necessarily the case in general.

To characterize the canonical endomorphism, Longo [43] introduced the notion of Q-systems.

Definition 3.1. Let $\mathcal{M}$ be a properly infinite factor and let $\gamma \in \mathrm{End}_0(\mathcal{M})$. We say that (γ, v, w) is a Q-system if the following three conditions are satisfied:

(1) $v \in (\mathrm{id}_{\mathcal{M}}, \gamma)$ and $w \in (\gamma, \gamma^2)$ are isometries.
(2) There exists a positive number d such that $v^*w = w^*\gamma(v) = 1/d$.
(3) $ww = \gamma(w)w$.

Two Q-systems (γ, v, w) and (γ', v', w') are said to be equivalent if there exists a unitary $u \in (\gamma, \gamma')$ such that $v' = uv$ and $w' = u\gamma(u)wu^*$.

It is known that (3) is equivalent to $ww^* = \gamma(w^*)w$ under the assumption (1) and (2) [45], [27]. If $\gamma = \sigma\bar{\sigma}$ with $\sigma \in \mathrm{Mor}_0(\mathcal{N}, \mathcal{M})$ and $v = \bar{r}_\sigma \in (\mathrm{id}_{\mathcal{M}}, \gamma)$, $w = \sigma(r_\sigma) \in \sigma((\mathrm{id}_{\mathcal{N}}, \bar{\sigma}\sigma)) \subset (\gamma, \gamma^2)$, then (γ, v, w) is a Q-system with $d = d(\sigma)$. The equivalence class of this Q-system only depends on the inner conjugacy class of $\sigma(\mathcal{N})$ in $\mathcal{M}$. Thus we say that (γ, v, w) *arises from* $\sigma(\mathcal{N}) \subset \mathcal{M}$. On the other hand, Longo [43] showed that any Q-system (γ, v, w) arises from a subfactor $\mathcal{N} \subset \mathcal{M}$, determined by $E_{\mathcal{N}}(x) = w^*\gamma(x)w$, and equivalence of two Q-systems exactly corresponds to inner conjugacy of the corresponding two subfactors.

Let $\mathcal{N} \subset \mathcal{M}$ be an inclusion of properly infinite factors of finite index and let (γ, v, w) be the Q-system arising from the inclusion. The second-named author and Kosaki [27] introduced the “second cohomology" $H^2(\mathcal{N} \subset \mathcal{M})$, which is always a finite set, by the equivalence classes of Q-systems (γ_1, v_1, w_2) such that γ and γ_1 are equivalent. When $\mathcal{N}$ is the fixed point algebra of $\mathcal{M}$ by an outer action of a finite group G, then we can identify $H^2(\mathcal{N} \subset \mathcal{M})$ with the second cohomology group $H^2(G, \mathbb{T})$.

The following is [27, Lemma 6.1].

Lemma 3.2. *Let $\mathcal{N} \subset \mathcal{M}$ be an irreducible inclusion of factors of finite index. If $\mathcal{N} \subset \mathcal{M}$ is 3-supertransitive, then $H^2(\mathcal{N} \subset \mathcal{M})$ is trivial.*

When $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ is a noncommuting quadrilateral of subfactors and $N \subset P$ and $N \subset Q$ are 2-supertransitive, the two inclusions $\check{P} \subset N$ and $\check{Q} \subset N$ give equivalent canonical endomorphisms as we observed above.

Definition 3.3. Let $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ be an irreducible noncommuting quadrilateral of factors with 2-supertransitive $N \subset P$ and $N \subset Q$. We denote by $c(\mathfrak{Q})$ the class of the Q-system arising from $\check{Q} \subset N$ in $H^2(\widehat{N \subset P})$.

The class $c(\mathfrak{Q})$ is very much related to the angle $\Theta(P, Q)$ and there are at most two possibilities for $\Theta(P, Q)$ for a given class $c(\mathfrak{Q})$ as we will see below.

The following lemma is an easy consequence of the definition above:

Lemma 3.4. *For an irreducible noncommuting quadrilateral $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ of factors with 2-supertransitive $N \subset P$ and $N \subset Q$, the following three conditions are equivalent:*

(1) *The class* $c(\mathfrak{Q}) \in H^2(\widehat{N \subset P})$ *is trivial.*
(2) *The two subfactors* $\check{P}$ *and* $\check{Q}$ *are inner conjugate in* N*.*
(3) *There exists an isomorphism* $\pi : P \to Q$ *of* P *onto* Q *such that the restriction* $\pi|_N$ *of* π *to* N *is the identity map.*

Since the Q-systems we encounter in this paper always have γ decomposed into two irreducible components, we give a detailed description of the condition (3) of Definition 3.1 in this case. Let $\sigma \in \mathrm{End}_0(\mathcal{M})$ be an irreducible endomorphism satisfying $[\mathrm{id}_{\mathcal{M}}] \neq [\sigma]$ and let $\gamma = \mathrm{id}_{\mathcal{M}} \oplus \sigma$. Note that for γ to have a Q-system, the endomorphism σ must be self-conjugate. We fix isometries $v \in (\mathrm{id}_{\mathcal{M}}, \gamma)$ and $v_1 \in (\sigma, \gamma)$, which means $\gamma(x) = vxv^* + v_1\sigma(x)v_1^*$. We give a description of a Q-system of the form (γ, v, w) in terms of an isometry in (σ, σ^2). It is easy to show that an isometry $w \in (\gamma, \gamma^2)$ satisfying (2) of Definition 3.1 is of the following form and vice versa:

$$w = \frac{v}{\sqrt{d(\sigma)+1}} + \frac{v_1\sigma(v)v_1^*}{\sqrt{d(\sigma)+1}} + \sqrt{\frac{d(\sigma)}{d(\sigma)+1}}cv_1\sigma(v_1)r_\sigma v^* + \sqrt{\frac{d(\sigma)-1}{d(\sigma)+1}}v_1\sigma(v_1)sv_1^*,$$

where c is a complex number with $|c| = 1$ and $s \in (\sigma, \sigma^2)$ is an isometry. Passing from w to $u\gamma(u)wu^*$ with $u = vv^* + \bar{c}v_1v_1^*$ (or redefining r_σ) if necessary, we may always assume $c = 1$ for a Q-system (γ, v, w). We say that such a Q-system is normalized (with respect to (v, v_1, r_σ)). It is straightforward to show the following:

Lemma 3.5. *Let* σ*,* γ*,* v*, and* v_1 *be as above. If* $s \in (\sigma, \sigma^2)$ *is an isometry satisfying*

$$\sigma(s)r_\sigma = sr_\sigma, \tag{3.1}$$

$$\sqrt{d(\sigma)}r_\sigma + (d(\sigma)-1)s^2 = \sqrt{d(\sigma)}\sigma(r_\sigma) + (d(\sigma)-1)\sigma(s)s, \tag{3.2}$$

and w *is given by*

$$w = \frac{v}{\sqrt{d(\sigma)+1}} + \frac{v_1\sigma(v)v_1^*}{\sqrt{d(\sigma)+1}} + \sqrt{\frac{d(\sigma)}{d(\sigma)+1}}v_1\sigma(v_1)r_\sigma v^* + \sqrt{\frac{d(\sigma)-1}{d(\sigma)+1}}v_1\sigma(v_1)sv_1^*, \tag{3.3}$$

then (γ, v, w) *is a* Q*-system. Conversely, every normalized* Q*-system is of this form with an isometry* $s \in (\sigma, \sigma^2)$ *satisfying* (3.1) *and* (3.2)*. If* s *and* s' *are isometries in* (σ, σ^2) *satisfyingly* (3.1) *and* (3.2)*, the corresponding* Q*-systems are equivalent if and only if* $s = \pm s'$*.*

Remark 3.6. When the Q-system (γ, v, w) as above arises from a 3-supertransitive subfactor, we have $\dim(\sigma, \sigma^2) = 1$. If $s, s' \in (\sigma, \sigma^2)$ are isometries satisfying (3.1) and (3.2), there exists a complex number ω of modulus 1 such that $s' = \omega s$ and (3.2) implies $\omega^2 = 1$. This proves Lemma 3.2 again.

We keep using the same notation as above and consider the case with an inclusion $\mathcal{M} \subset \mathcal{M}_1$ such that $\gamma = \bar{\iota}\iota$, $v = r_\iota$, $w = \bar{\iota}(\bar{r}_\iota)$ where $\iota = \iota_{\mathcal{M}_1,\mathcal{M}}$. We set

$$t = \frac{d(\iota)}{\sqrt{d(\sigma)}}\iota(v_1^*)\bar{r}_\iota,$$

which is an isometry in $(\iota, \iota\sigma)$ thanks to Frobenius reciprocity [20, Proposition 2.2]. Then Theorem 2.9,(4) shows that for every $x \in \mathcal{M}_1$, we have

$$x = E_{\mathcal{M}}(x) + d(\sigma)E_{\mathcal{M}}(xt^*)t.$$

In particular,

$$t^2 = E_{\mathcal{M}}(t^2) + d(\sigma)E_{\mathcal{M}}(t^2t^*)t,$$
$$t^* = E_{\mathcal{M}}(t^*) + d(\sigma)E_{\mathcal{M}}(t^{*2})t = d(\sigma)E_{\mathcal{M}}(t^{*2})t,$$

where we use the fact $E_{\mathcal{M}}(t) \in (\mathrm{id}_{\mathcal{M}}, \sigma) = \{0\}$. Therefore the $*$-algebra structure of $\mathcal{M}_1$ is completely determined by $E_{\mathcal{M}}(t^2)$ and $E_{\mathcal{M}}(t^2t^*)$, which can be computed in terms of the Q-system as follows:

$$\begin{aligned} E_{\mathcal{M}}(t^2) &= \frac{d(\iota)^2}{d(\sigma)} r_\iota^* \bar{\iota}(\iota(v_1^*)\bar{r}_\iota \iota(v_1^*)\bar{r}_\iota) r_\iota = \frac{d(\iota)}{d(\sigma)} v_1^* \bar{\iota}(\iota(v_1^*)\bar{r}_\iota) r_\iota \\ &= \frac{d(\iota)}{d(\sigma)} \sigma(v_1^*) v_1^* w v = \frac{1}{\sqrt{d(\sigma)}} r_\sigma, \end{aligned}$$

$$\begin{aligned} E_{\mathcal{M}}(t^2t^*) &= \frac{d(\iota)^3}{\sqrt{d(\sigma)}^3} r_\iota^* \bar{\iota}(\iota(v_1^*)\bar{r}_\iota \iota(v_1^*)\bar{r}_\iota \bar{r}_\iota^* \iota(v_1)) r_\iota = \frac{d(\iota)}{\sqrt{d(\sigma)}^3} v_1^* \bar{\iota}(\iota(v_1^*)\bar{r}_\iota) v_1 \\ &= \frac{d(\iota)}{\sqrt{d(\sigma)}^3} \sigma(v_1^*) v_1^* w v_1 = \sqrt{\frac{d(\sigma)-1}{d(\sigma)^3}} s. \end{aligned}$$

Lemma 3.7. *Let the notation be as above. Then the following hold:*

$$t^2 = \frac{1}{\sqrt{d(\sigma)}} r_\sigma + \sqrt{\frac{d(\sigma)-1}{d(\sigma)}} st,$$
$$t^* = \sqrt{d(\sigma)} r_\sigma^* t.$$

Corollary 3.8. *Let $\mathcal{M}$ be a properly infinite factor and let $\sigma \in \mathrm{End}_0(\mathcal{M})$ be an irreducible self-conjugate endomorphism. If there exists a Q-system for $\mathrm{id}_{\mathcal{M}} \oplus \sigma$, then $[\sigma]$ is a real sector, that is, $r_\sigma = \bar{r}_\sigma$.*

Proof. Using one of the above formulae, we get

$$t = (t^*)^* = \sqrt{d(\sigma)}(r_\sigma^* t)^* = \sqrt{d(\sigma)} t^* r_\sigma = d(\sigma) r_\sigma^* t r_\sigma = d(\sigma) r_\sigma^* \sigma(r_\sigma) t,$$

which shows $r_\sigma^* \sigma(r_\sigma) = 1/d(\sigma)$. ☐

Now we show how the class $c(\mathfrak{Q})$ determines the angle between P and Q. Let $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ be an irreducible quadrilateral of factors such that $N \subset P$ and $N \subset Q$ are 2-supertransitive. We assume that ${}_N L^2(P)_N$ and ${}_N L^2(Q)_N$ are equivalent. We apply the above argument to the case where $\mathcal{M}_1 = P$, $\mathcal{M} = N$, $\iota = \iota_{P,N}$ and $\gamma = \bar{\iota}\iota$. We set $\check{P} = \bar{\iota}(P)$. We choose two isometries $v = r_\iota \in (\mathrm{id}, \gamma)$ and $v_1 \in (\sigma, \gamma)$. Then we may assume that the Q-system $(\gamma, v, w_P = \bar{\iota}(\bar{r}_\iota))$ is normalized with respect to (v, v_1, r_σ). By choosing an appropriate representative of the conjugate sector of $[\iota_{Q,N}]$, we may also assume $\gamma = \overline{\iota_{Q,N}}\iota_{Q,N}$, $v = r_{\iota_{Q,N}}$, and that the Q-system $(\gamma, v, w_Q = \overline{\iota_{Q,N}}(\bar{r}_{\iota_{Q,N}}))$ is normalized with respect to (v, v_1, r_σ). By definition, the class $c(\mathfrak{Q})$ is given by (γ, v, w_Q) in $H^2(\check{P} \subset N)$. Let s_P and s_Q be isometries in (σ, σ^2) corresponding to w_P and w_Q through (3.3) respectively. We set

$$t_P = \sqrt{\frac{d(\sigma)+1}{d(\sigma)}}\iota_{P,N}(v_1^*)\bar{r}_{\iota_{P,N}} \in (\iota_{P,N}, \iota_{P,N}\sigma),$$

$$t_Q = \sqrt{\frac{d(\sigma)+1}{d(\sigma)}}\iota_{Q,N}(v_1^*)\bar{r}_{\iota_{Q,N}} \in (\iota_{Q,N}, \iota_{Q,N}\sigma),$$

which are isometries.

Lemma 3.9. *Let* $\mathfrak{Q} = \begin{matrix} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{matrix}$ *be an irreducible quadrilateral of factors such that* $N \subset P$ *and* $N \subset Q$ *are 2-supertransitive. If* $L^2(P)$ *and* $L^2(Q)$ *are equivalent as* $N-N$ *bimodules, then*

(1) $\langle t_P, t_Q\rangle$ *satisfies the following quadratic equation:*

$$\langle t_P, t_Q\rangle^2 - \frac{d(\sigma)-1}{d(\sigma)}\langle s_P, s_Q\rangle\langle t_P, t_Q\rangle - \frac{1}{d(\sigma)} = 0.$$

(2) $\langle s_P, s_Q\rangle$ *and* $\langle t_P, t_Q\rangle$ *are real numbers.*

Proof. (1) Lemma 3.7 implies

$$\langle t_P, t_Q\rangle^2 = {t_Q^2}^* t_P^2 = \frac{1}{d(\sigma)} + \frac{d(\sigma)-1}{d(\sigma)}\langle s_P, s_Q\rangle\langle t_P, t_Q\rangle.$$

(2) Thanks to Lemma 2.10, we have $E_P(t_Q) = \langle t_Q, t_P\rangle t_P$ and $E_Q(t_P) = \langle t_P, t_Q\rangle t_Q$. Thus

$$E_N(t_P t_Q) = E_N(E_P(t_P t_Q)) = \langle t_Q, t_P\rangle E_N(t_P^2) = \frac{\langle t_Q, t_P\rangle}{\sqrt{d(\sigma)}}r_\sigma,$$

$$E_N(t_P t_Q) = E_N(E_Q(t_P t_Q)) = \langle t_P, t_Q\rangle E_N(t_Q^2) = \frac{\langle t_P, t_Q\rangle}{\sqrt{d(\sigma)}}r_\sigma,$$

which shows that $\langle t_P, t_Q\rangle$ is real. This and (1) show that $\langle s_P, s_Q\rangle$ is also real. □

Note that $\dim(\iota_{P,N}, \iota_{P,N}\sigma) = \dim(\iota_{Q,N}, \iota_{Q,N}\sigma) = 1$ by Frobenius reciprocity. Thus Remark 2.11 shows $\cos\Theta(P,Q) = |\langle t_P, t_Q\rangle|$.

Theorem 3.10. *Let* $\mathfrak{Q} = \begin{matrix} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{matrix}$ *be an irreducible quadrilateral of factors such that* $N \subset P$ *and* $N \subset Q$ *are 2-supertransitive. If* $L^2(P)$ *and* $L^2(Q)$ *are equivalent as* $N-N$ *bimodules, then*

(1) $\mathfrak{Q}$ *is not commuting.*

(2) *The angle* $\Theta(P,Q)$ *is given by*

$$\begin{aligned}\cos\Theta(P,Q) &= \frac{\sqrt{(d(\sigma)-1)^2\langle s_P,s_Q\rangle^2+4d(\sigma)}+(d(\sigma)-1)|\langle s_P,s_Q\rangle|}{2d(\sigma)}\\ \text{or} &= \frac{\sqrt{(d(\sigma)-1)^2\langle s_P,s_Q\rangle^2+4d(\sigma)}-(d(\sigma)-1)|\langle s_P,s_Q\rangle|}{2d(\sigma)}.\end{aligned}$$

Proof. (1) If $\mathfrak{Q}$ were commuting, we would have $\langle t_P,t_Q\rangle=0$, which never occurs due to Lemma 3.9,(1). (2) follows from Lemma 3.9 too. □

Note that $|\langle s_P,s_Q\rangle|$ is a numerical invariant of the class $c(\mathfrak{Q})$. Therefore the above theorem says that there are only two possibilities of the angle $\Theta(P,Q)$ given $c(\mathfrak{Q})$.

Corollary 3.11. *Let* $\mathfrak{Q}=\begin{array}{ccc}P&\subset&M\\ \cup&&\cup\\ N&\subset&Q\end{array}$ *be an irreducible noncommuting quadrilateral of factors such that* $N\subset P$ *and* $N\subset Q$ *are 2-supertransitive. Then*

$$0<\Theta(P,Q)\leq\cos^{-1}\frac{1}{[P:N]-1}.$$

The equality holds if and only if the class $c(\mathfrak{Q})\in H^2(\check{P}\subset N)$ *is trivial. In particular, if* $N\subset P$ *is 3-supertransitive,*

$$\Theta(P,Q)=\cos^{-1}\frac{1}{[P:N]-1}.$$

Proof. Note that $|\langle t_P,t_Q\rangle|=1$ never occurs since $|\langle t_P,t_Q\rangle|=1$ would imply $P=Q$. The statement follows from

$$\frac{1}{d(\sigma)}\leq\frac{\sqrt{(d(\sigma)-1)^2\langle s_P,s_Q\rangle^2+4d(\sigma)}-(d(\sigma)-1)|\langle s_P,s_Q\rangle|}{2d(\sigma)},$$

where the quality holds if and only if $\langle s_P,s_Q\rangle=\pm1$. □

Corollary 3.12. *Let* $\mathfrak{Q}=\begin{array}{ccc}P&\subset&M\\ \cup&&\cup\\ N&\subset&Q\end{array}$ *be an irreducible noncommuting quadrilateral of factors such that* $N\subset P$ *and* $N\subset Q$ *are 2-supertransitive. If* $\mathfrak{Q}$ *is cocommuting, then*

$$[M:P]\leq[P:N]-1.$$

The equality holds if and only if the class $c(\mathfrak{Q})\in H^2(\check{P}\subset N)$ *is trivial. In particular, if* $\mathfrak{Q}$ *is cocommuting and* $N\subset P$ *is 3-supertransitive,*

$$[M:P]=[P:N]-1.$$

Proof. This follows from Lemma 2.4 and Lemma 3.11. □

Remark 3.13. There is an example of a quadrilateral $\mathfrak{Q}$ fulfilling the assumption of Corollary 3.12 with non-trivial $c(\mathfrak{Q})$. Let q be a prime power and let F be a finite field with q elements. We set $G=PSL(n,q)$ with $n\geq3$, which naturally acts on the projective space PF^{n-1}. It is known that this action is 2-transitive. We denote by

F^{n*} the dual space of F^n, on which G naturally acts too. Let $\{e_i\}_{i=1}^n$ be the canonical basis of F^n and let $\{e_i^*\}_{i=1}^n$ be the dual basis of $\{e_i\}_{i=1}^n$. Let H be the stabilizer of Fe_1 and let K be the stabilizer of Fe_1^*. Then we have $[G : H] = [G : K] = (q^n - 1)/(q - 1)$ and $[H : H \cap K] = [K : H \cap K] = q^{n-1}$. Let $\mathfrak{Q}$ be the quadrilateral constructed by Lemma 2.5 with these G, H, and K. Then $[M : P]$ is strictly smaller than $[P : N] - 1$. In this case we have $\cos\Theta(P, Q) = q^{-n/2}$.

Lemma 3.14. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *be an irreducible noncommuting quadrilateral of factors such that* $N \subset P$ *and* $N \subset Q$ *are 2-supertransitive and* $H^2(\widehat{N \subset P})$ *is trivial. Let* $[\overline{\iota_{P,N}}\iota_{P,N}] = [\mathrm{id}_N] \oplus [\sigma]$. *If* $\dim(\iota_{M,N}, \iota_{M,N}\sigma) = 2$ *and there is an intermediate subfactor* $N \subset R \subset M$ *other than* P *and* Q *such that* $L^2(R)$ *is equivalent to* $L^2(P)$ *as* $N - N$ *bimodules, then* $[P : N] = 3$.

Proof. Let t_P and t_Q be as in Lemma 3.9. We may and do assume $\langle t_P, t_Q\rangle = 1/d(\sigma)$ by replacing t_Q with $-t_Q$ if necessary. Assume that there exists an intermediate subfactor $N \subset R \subset M$ other than P and Q such that $L^2(R)$ is equivalent to $L^2(P)$ as $N - N$ bimodules. Then we may choose an isometry $t \in (\iota_{R,N}, \iota_{R,N}\sigma)$ such that $\langle t, t_P\rangle = 1/d(\sigma)$, $\langle t, t_Q\rangle = \epsilon/d(\sigma)$ with $\epsilon \in \{1, -1\}$, and $E_N(t^2) = 1/\sqrt{d(\sigma)}r_\sigma$. Since $\dim(\iota_{M,N}, \iota_{M,N}\sigma) = 2$, the intertwiner space $(\iota_{M,N}, \iota_{M,N}\sigma)$ is spanned by t_P and t_Q. Therefore, the first two equalities imply

$$t = \frac{d(\sigma) - \epsilon}{d(\sigma)^2 - 1}(t_P + \epsilon t_Q).$$

Since t is an isometry, we get

$$\begin{aligned} 1 &= \langle t, t\rangle = \frac{(d(\sigma) - \epsilon)^2}{(d(\sigma)^2 - 1)^2}\langle t_P + \epsilon t_Q, t_P + \epsilon t_Q\rangle = \frac{(d(\sigma) - \epsilon)^2}{(d(\sigma)^2 - 1)^2}\frac{2(d(\sigma) + \epsilon)}{d(\sigma)} \\ &= \frac{2(d(\sigma) - \epsilon)}{d(\sigma)(d(\sigma)^2 - 1)} \end{aligned}$$

If $\epsilon = 1$, we would have $d(\sigma) = 1$, which contradicts $\dim(\iota_{M,N}, \iota_{M,N}\sigma) = 2$ and Theorem 2.9,(1). Therefore we get $\epsilon = -1$ and $d(\sigma) = 2$, which means $[P : N] = 3$. □

Remark 3.15. Let $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ be an irreducible noncommuting quadrilateral of factors. Then there exists σ contained both in $\overline{\iota_{P,N}}\iota_{P,N}$ and in $\overline{\iota_{Q,N}}\iota_{Q,N}$ so that the multiplicity of σ in $\overline{\iota_{M,N}}\iota_{M,N}$ is at least 2. Therefore σ is not an automorphism due to Theorem 2.9,(1). In particular, the subfactor $N \subset P$ is not the crossed product by a finite group action.

4. (2,2)-SUPERTRANSITIVE QUADRILATERALS

In what follows, for a non-commuting quadrilateral $\begin{matrix} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{matrix}$ of properly infinite factors with 2-supertransitive elementary subfactors, we use the following notation: the symbols ι, κ, ι_Q, and κ_Q denote the inclusion maps

$$\iota : N \hookrightarrow P, \quad \kappa : P \hookrightarrow M, \quad \iota_Q : N \hookrightarrow Q, \quad \kappa_Q : Q \hookrightarrow M.$$

Since $\bar{\iota}\iota$, $\iota\bar{\iota}$, etc are decomposed into two irreducible sectors, one of which is the identity sector, we choose representatives $\xi, \eta \in \mathrm{End}_0(P)$, $\xi_Q, \eta_Q \in \mathrm{End}_0(Q)$, $\hat{\xi}, \hat{\xi}_Q \in \mathrm{End}_0(N)$, and $\hat{\eta}, \hat{\eta}_Q \in \mathrm{End}_0(M)$ satisfying

$$[\iota\bar{\iota}] = [\mathrm{id}_P] \oplus [\xi], \quad [\bar{\kappa}\kappa] = [\mathrm{id}_P] \oplus [\eta],$$

$$[\iota_Q\bar{\iota}_Q] = [\mathrm{id}_Q] \oplus [\xi_Q], \quad [\bar{\kappa}_Q\kappa_Q] = [\mathrm{id}_Q] \oplus [\eta_Q],$$

$$[\bar{\iota}\iota] = [\bar{\iota}_Q\iota_Q] = [\mathrm{id}_N] \oplus [\hat{\xi}],$$

$$[\kappa\bar{\kappa}] = [\mathrm{id}_M] \oplus [\hat{\eta}], \quad [\kappa_Q\bar{\kappa}_Q] = [\mathrm{id}_M] \oplus [\hat{\eta}_Q].$$

Note that neither ξ nor $\hat{\xi}$ is an automorphism (see Remark 3.15). When $N \subset P$ and $N \subset Q$ are 3-supertransitive, the sectors $\xi\iota$ and $\xi_Q\iota_Q$ are decomposed into two irreducible components and we use the following notation:

$$[\xi\iota] = [\iota] \oplus [\iota'], \quad [\xi_Q\iota_Q] = [\iota_Q] \oplus [\iota'_Q].$$

In this case, we also have

$$[\iota\hat{\xi}] = [\iota] \oplus [\iota'], \quad [\iota_Q\hat{\xi}_Q] = [\iota_Q] \oplus [\iota'_Q].$$

In the same way, when $P \subset M$ and $Q \subset M$ are 3-supertransitive, we use the following notation:

$$[\kappa\eta] = [\kappa] \oplus [\kappa'], \quad [\kappa_Q\eta_Q] = [\kappa_Q] \oplus [\kappa'_Q],$$

$$[\hat{\eta}\kappa] = [\kappa] \oplus [\kappa'], \quad [\hat{\eta}_Q\kappa_Q] = [\kappa_Q] \oplus [\kappa'_Q].$$

In the sequel we often use the techniques developed in [17]. For example, let ρ, σ be irreducible endomorphisms of P associated with the inclusion $N \subset P$. Then $\rho\xi$ contains σ if and only if either $[\rho] = [\sigma]$ or the distance between ρ and σ in the dual principal graph of $N \subset P$ is two. When $[\rho] \neq [\sigma]$, the multiplicity of σ in $\rho\xi$ is the number of the paths from ρ to σ. The multiplicity of ρ in $\rho\xi$ is $\sum_{i=1}^m n_i^2 - 1$ where $[\rho\iota] = \oplus_i^m n_i[\tau_i]$, is the irreducible decomposition of $\rho\iota$. In particular, the inclusion $N \subset P$ is 3-supertransitive if and only if the multiplicity of ξ in ξ^2 is one.

Lemma 4.1. *Let* $\mathfrak{Q} = \begin{matrix} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{matrix}$ *be an irreducible noncommuting quadrilateral of factors. We assume that* $\mathfrak{Q}$ *is (2,2)-supertransitive. Then*

(1) $[\xi] \neq [\eta]$.

(2) ξ^2 *contains* η. *Consequently, the depth of* $N \subset P$ *is at least 4.*

Proof. (1) Since $N \subset M$ is irreducible, we have $\dim(\kappa\iota, \kappa\iota) = 1$. Thus Frobenius reciprocity implies

$$1 = \dim(\kappa\iota, \kappa\iota) = (\bar{\kappa}\kappa, \iota\bar{\iota}) = (\mathrm{id}_P \oplus \eta, \mathrm{id}_P \oplus \xi),$$

which shows $[\xi] \neq [\eta]$.

(2) Since $\mathfrak{Q}$ is noncommuting and

$$L^2(P) \cong L^2(Q) \cong L^2(N) \oplus L^2(N)_{\hat{\xi}},$$

as $N - N$ bimodules, the $N - N$ bimodule $L^2(M)$ contains $L^2(N)_{\hat{\xi}}$ with multiplicity at least two. Thus

$$\begin{aligned} 2 &\leq \dim(\bar{\iota}\bar{\kappa}\kappa\iota, \hat{\xi}) = \dim(\bar{\kappa}\kappa, \iota\hat{\xi}\bar{\iota}) = \dim(\bar{\kappa}\kappa, \iota\bar{\iota}\iota\bar{\iota}) - \dim(\bar{\kappa}\kappa, \iota\bar{\iota}) \\ &= \dim(\mathrm{id}_P \oplus \eta, \xi \oplus \xi^2) = 1 + \dim(\eta, \xi^2), \end{aligned}$$

which shows the statement. □

Remark 4.2. Using the above lemma, we give a short proof of Theorem 2.7 only assuming (4,4)-supertransitivity of $\mathfrak{Q}$ (see Corollary 4.9 and Corollary 5.18 below for sharper results in this direction). Since $N \subset P$ is 4-supertransitive, Lemma 4.1 shows

$$[\xi^2] = [\mathrm{id}_P] \oplus [\xi] \oplus [\eta].$$

If the principal graph is A_5, η is an automorphism and M is the crossed product $P \rtimes_\eta \mathbb{Z}/2\mathbb{Z}$. Therefore the proof of [18, Theorem 3.1] implies that there exists an outer action of $\mathfrak{S}_3$ on M such that N is the fixed point algebra of the action. Assume now that the principal graph of $N \subset M$ is not A_5. Then we have $\dim(\eta\iota, \eta\iota) \geq 2$. Frobenius reciprocity implies

$$\begin{aligned} \dim(\eta^2, \xi) &= \dim(\eta, \eta\xi) = \dim(\eta, \eta\iota\bar{\iota}) - \dim(\eta, \eta) = \dim(\eta\iota, \eta\iota) - 1 \\ &\geq 1. \end{aligned}$$

Since $P \subset M$ is 4-supertransitive, we get $[\eta^2] = [\mathrm{id}_P] \oplus [\eta] \oplus [\xi]$. Therefore $d(\xi) = d(\eta)$ and $d(\xi)^2 = 1 + 2d(\xi)$, which shows that $[M : P] = [P : N] = 2 + \sqrt{2} = 4\cos^2(\pi/8)$ and all the elementary subfactors are the A_7 subfactor. Uniqueness of such a quadrilateral follows from Lemma 3.2 and Lemma 3.14 as η is uniquely determined by $N \subset P$ (see Theorem 4.7 below for details).

The following is one of the main results in this paper, which says that irreducible noncommuting (2,2)-supertransitive quadrilaterals with trivial cohomological obstruction are classified into four classes and one exceptional case.

Theorem 4.3. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *be an irreducible noncommuting quadrilateral of factors. We assume that* $\mathfrak{Q}$ *is (2,2)-supertransitive and the class* $c(\mathfrak{Q}) \in H^2(\widehat{N \subset P})$ *is trivial. Then*

$$\Theta(M, N) = \cos^{-1} \frac{1}{[P : N] - 1}.$$

Moreover,

(1) *If* $\mathfrak{Q}$ *is not cocommuting and the class* $c(\hat{\mathfrak{Q}})$ *in* $H^2(P \subset M)$ *is trivial, then* $[M : P] = [P : N]$. *In this case, there exists an outer automorphism* α *of* P *such that* $[\eta] = [\alpha\xi] = [\xi\alpha^{-1}]$ *and* ξ^2 *contains* η. (*In consequence, the automorphism* α *is contained in* ξ^3.)

(2) *If* $\mathfrak{Q}$ *is cocommuting, then* $[M : P] = [P : N] - 1$ *and* $[\xi] = [\bar{\kappa}\kappa_Q\pi]$, *where* π *is an isomorphism from* P *to* Q *such that the restriction of* π *to* N *is the identity map. In this case, the Galois group*

$$\mathrm{Gal}(M/N) = \{\theta \in \mathrm{Aut}(M);\ \theta(x) = x,\ \forall x \in N\}$$

is either trivial or isomorphic to $\mathbb{Z}/2\mathbb{Z}$,$\mathbb{Z}/3\mathbb{Z}$, *or the symmetric group* $\mathfrak{S}_3$ *of degree 3.*

If $\mathrm{Gal}(M/N) = \{\mathrm{id}_M, \theta\} \cong \mathbb{Z}/2\mathbb{Z}$, *then* θ *switches* P *and* Q *and* $[\xi] = [\bar{\kappa}\theta\kappa]$.

If $\mathrm{Gal}(M/N) = \{\mathrm{id}_M, \theta, \theta^2\} \cong \mathbb{Z}/3\mathbb{Z}$, *then either* $\theta(P) = Q$ *or* $\theta(Q) = P$ *holds and* $[\xi] = [\bar{\kappa}\theta\kappa] = [\bar{\kappa}\theta^2\kappa]$.

If $\mathrm{Gal}(M/N) \cong \mathfrak{S}_3$, *then* N *is the fixed point algebra of an outer action of* $\mathfrak{S}_3$ *on* M *and* P *and* Q *are the fixed point algebras of two different order two elements in* $\mathfrak{S}_3$.

Proof. The angle between P and Q was already computed in Theorem 3.10. Lemma 3.4 shows that since $c(\mathfrak{Q})$ is trivial there exists an isomorphism π from P onto Q such that the restriction of π to N is trivial. This means $\iota_Q = \pi\iota$. Thus $\kappa\iota = \kappa_Q\iota_Q$ implies $\kappa\iota = \kappa_Q\pi\iota$ and

$$\begin{aligned} 1 &= \dim(\kappa\iota, \kappa_Q\pi\iota) = \dim(\kappa\iota\bar{\iota}, \kappa_Q\pi) = \dim(\kappa, \kappa_Q\pi) + \dim(\kappa\xi, \kappa_Q\pi) \\ &= \dim(\kappa, \kappa_Q\pi) + \dim(\xi, \bar{\kappa}\kappa_Q\pi). \end{aligned}$$

We claim $[\kappa] \neq [\kappa_Q\pi]$, and in consequence, claim that ξ is contained in $\bar{\kappa}\kappa_Q\pi$. Indeed, if $[\kappa] = [\kappa_Q\pi]$ were the case, there would exist a unitary u in M such that $x = u\pi(x)u^*$ holds for all $x \in P$. In particular, the unitary u commutes with all $x \in N$ and u is a scalar. However, this means $P = Q$, which is a contradiction. Therefore the claim holds.

(1) Assume that $\mathfrak{Q}$ is not cocommuting and the class $c(\hat{\mathfrak{Q}})$ in $H^2(P \subset M)$ is trivial. Then Lemma 3.4 applied to the dual quadrilateral $\hat{\mathfrak{Q}}$ implies that P and Q are inner conjugate in M and there exists a unitary $v \in M$ satisfying $vQv^* = P$. We introduce an automorphism α of P by setting $\alpha(x) = v\pi(x)v^*$ for $x \in P$. This implies $[\kappa\alpha] = [\kappa_Q\pi]$ and so

$$[\bar{\kappa}\kappa_Q\pi] = [\bar{\kappa}\kappa\alpha] = [\alpha] \oplus [\eta\alpha].$$

Since ξ is contained in $\bar{\kappa}\kappa_Q\pi$, either $[\xi] = [\alpha]$ or $[\xi] = [\alpha\eta]$ holds. Note that $[\xi] = [\alpha]$ never occurs thanks to Remark 3.15. Thus we have $[\xi] = [\eta\alpha]$ and so $[\eta] = [\xi\alpha^{-1}]$, which is equal to $[\alpha\xi]$ for η is self-conjugate. Since ξ^2 contains η and id_P, we conclude that ξ^3 contains α.

(2) Assume that $\mathfrak{Q}$ is cocommuting. Then Theorem 3.10,(1) applied to the dual quadrilateral $\hat{\mathfrak{Q}}$ implies that $[\kappa\bar{\kappa}] \neq [\kappa_Q\bar{\kappa}_Q]$ and so

$$\dim(\bar{\kappa}\kappa_Q\pi, \bar{\kappa}\kappa_Q\pi) = \dim(\kappa\bar{\kappa}, \kappa_Q\bar{\kappa}_Q) = 1.$$

This shows that $\bar{\kappa}\kappa_Q\pi$ is irreducible and $[\xi] = [\bar{\kappa}\kappa_Q\pi]$, which implies

$$[P : N] - 1 = d(\xi) = d(\kappa)d(\kappa_Q) = [M : P].$$

(This is also shown in Corollary 3.12 with help of Lemma 2.4 though we don't need to use it here.)

Note that the Galois group of the inclusion $N \subset M$ is isomorphic to the set of 1-dimensional sectors G contained in $\kappa\iota\bar{\iota}\bar{\kappa}$. (Note that since $N \subset M$ is irreducible, the multiplicity of any automorphism in $\kappa\iota\bar{\iota}\bar{\kappa}$ is at most one.) We assume that G is non-trivial now. Using $[\xi] = [\bar{\kappa}\kappa_Q\pi]$, we get

$$\begin{aligned}[\kappa\iota\bar{\iota}\bar{\kappa}] &= [\kappa\bar{\kappa}] \oplus [\kappa\xi\bar{\kappa}] = [\mathrm{id}_M] \oplus [\hat{\eta}] \oplus [\kappa\bar{\kappa}\kappa_Q\pi\bar{\kappa}] \\ &= [\mathrm{id}_M] \oplus [\hat{\eta}] \oplus [\kappa_Q\pi\bar{\kappa}] \oplus [\hat{\eta}\kappa_Q\pi\bar{\kappa}].\end{aligned}$$

Assume that $\hat{\eta}$ is an automorphism. Then $[M : P] = 2$ and so $[P : N] = 3$. Since $N \subset P$ is 2-supertransitive, it is the A_5 subfactor. Therefore, the proof of [18, Theorem 3.1] shows that there exists a unique outer action of $\mathfrak{S}_3$ on M such that N is the fixed point algebra of the action.

Assume that $\hat{\eta}$ is not an automorphism. We claim that $\kappa_Q\pi\bar{\kappa}$ contains an automorphism. By the assumption of non-triviality of $\mathrm{Gal}(M/N)$, either $\kappa_Q\pi\bar{\kappa}$ or $\hat{\eta}\kappa_Q\pi\bar{\kappa}$ contains an automorphism. Assume that the latter contains an automorphism, say α. Then Frobenius reciprocity implies

$$1 = \dim(\hat{\eta}\kappa_Q\pi\bar{\kappa}, \alpha) = \dim(\kappa_Q\pi\bar{\kappa}, \hat{\eta}\alpha),$$

and $\kappa_Q\pi\bar{\kappa}$ contains $\hat{\eta}\alpha$. Since $d(\kappa_Q\pi\bar{\kappa}) = d(\hat{\eta}\alpha) + 1$, we conclude that $\kappa_Q\pi\bar{\kappa}$ contains an automorphism.

Let θ be an automorphism contained in $\kappa_Q\pi\bar{\kappa}$. Then Frobenius reciprocity implies $[\theta\kappa] = [\kappa_Q\pi]$. We may assume that $\theta\kappa = \kappa_Q\pi$ holds by choosing an appropriate representative of $[\theta]$. Then $\theta(P) = Q$ and $\theta\kappa\iota = \kappa_Q\pi\iota = \kappa_Q\iota_Q = \kappa\iota$, which shows $\theta \in \mathrm{Gal}(M, N)$. Using $[\theta\kappa] = [\kappa_Q\pi]$, we get

$$\begin{aligned}[\kappa\iota\bar{\iota}\bar{\kappa}] &= [\mathrm{id}_M] \oplus [\hat{\eta}] \oplus [\theta\kappa\bar{\kappa}] \oplus [\hat{\eta}\theta\kappa\bar{\kappa}] \\ &= [\mathrm{id}_M] \oplus [\theta] \oplus [\hat{\eta}] \oplus [\theta\hat{\eta}] \oplus [\hat{\eta}\theta] \oplus [\hat{\eta}\theta\hat{\eta}].\end{aligned}$$

If $\hat{\eta}\theta\hat{\eta}$ does not contain any automorphism, then $\mathrm{Gal}(M, N) = \{\mathrm{id}_M, \theta\} \cong \mathbb{Z}/2\mathbb{Z}$.

Assume that $\hat{\eta}\theta\hat{\eta}$ does contain an automorphism, say, β. Then Frobenius reciprocity again implies $[\hat{\eta}\theta] = [\beta\hat{\eta}]$ (and $[\theta\hat{\eta}] = [\hat{\eta}\beta]$), and

$$[\kappa\iota\bar{\iota}\bar{\kappa}] = [\mathrm{id}_M] \oplus [\theta] \oplus [\hat{\eta}] \oplus [\theta\hat{\eta}] \oplus [\hat{\eta}\theta] \oplus [\beta\hat{\eta}^2].$$

Therefore

$$G = \{[\mathrm{id}_M], [\theta]\} \cup \{[\beta\gamma]\}_{\gamma\in\mathrm{Gal}(M/\hat{\eta}(M))}.$$

Since $[\beta] \in G$, the group $G_1 = \{[\gamma]\}_{\gamma\in\mathrm{Gal}(M/\hat{\eta}(M))}$ is a subgroup of G such that $\#G = 2 + \#G_1$. Since $\#G_1$ divides $\#G$, either G_1 is trivial or $\#G_1 = 2$. If G_1 is trivial, we get $\mathrm{Gal}(M/N) \cong \{[\mathrm{id}_M], [\theta], [\beta]\} \cong \mathbb{Z}/3\mathbb{Z}$.

Suppose that $G_1 = \{[\mathrm{id}_M], [\gamma]\}$. Then $G = \{[\mathrm{id}_M], [\theta], [\beta], [\beta\gamma]\}$ and $[\theta] = [\gamma]$. This would imply $[\theta\hat{\eta}] = [\gamma\hat{\eta}] = [\hat{\eta}]$ and $[\hat{\eta}\theta\hat{\eta}] = [\hat{\eta}^2]$ would contain $[\mathrm{id}_M]$. However, this means that $\kappa\iota\bar{\iota}\bar{\kappa}$ contain id_M with multiplicity two, which is a contradiction. Therefore G_1 is trivial. □

Corollary 4.4. *Let* $\mathfrak{Q} = \begin{matrix} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{matrix}$ *be a (3,3)-supertransitive irreducible quadrilateral of factors such that* $\mathfrak{Q}$ *is neither commuting nor cocommuting. Then* $\Theta(P,Q) = \cos^{-1} 1/([P:N]-1)$, $[M:P]=[P:N]$ *and there exists an outer automorphism* α *of* P *such that* $[\eta]=[\alpha\xi]=[\xi\alpha^{-1}]$. *Moreover,*

1. *The automorphism* α *satisfies* $[\alpha\iota']=[\iota']$.
2. $\dim(\kappa\iota, \kappa\iota\hat{\xi}) = 2$.
3. *The Galois group* $\mathrm{Gal}(M/N)$ *is either trivial or isomorphic to* $\mathbb{Z}/2\mathbb{Z}$. *It is isomorphic to* $\mathbb{Z}/2\mathbb{Z}$ *if and only if the order of* $[\alpha]$ *is two. When* $\mathrm{Gal}(M/N) = \{\mathrm{id}, \theta\}$, *the automorphism* θ *switches* P *and* Q.

Proof. The first part follows from Lemma 3.2 and Theorem 4.3.

(1) Since $\alpha\xi$ is contained in $\iota'\bar{\iota}$, by Frobenius reciprocity ι' is contained in $\alpha\xi\iota = \alpha\iota\oplus\alpha\iota'$. Since $d(\iota') = d(\iota)(d(\iota)^2-2)$, the equality $[\iota']=[\alpha\iota]$ would imply that $N\subset P$ is the A_5 subfactor, which does not allow ξ^2 to contain $\alpha\xi$, and so $[\alpha\iota']=[\iota']$.

(2) Thanks to Lemma 4.1, we have

$$\begin{aligned}
[\bar{\iota}\bar{\kappa}\kappa\iota] &= [\bar{\iota}\iota]\oplus[\bar{\iota}\eta\iota] = [\mathrm{id}_N]\oplus[\hat{\xi}]\oplus[\bar{\iota}\alpha\xi\iota] = [\mathrm{id}_N]\oplus[\hat{\xi}]\oplus[\bar{\iota}\alpha\iota]\oplus[\bar{\iota}\alpha\iota'] \\
&= [\mathrm{id}_N]\oplus[\hat{\xi}]\oplus[\bar{\iota}\alpha\iota]\oplus[\bar{\iota}\iota'].
\end{aligned}$$

Frobenius reciprocity implies

$$\dim(\bar{\iota}\alpha\iota, \hat{\xi}) = \dim(\alpha\iota, \iota\hat{\xi}) = \dim(\alpha\iota,\iota)+\dim(\alpha\iota,\iota') = \dim(\alpha\iota,\iota),$$

where we use $[\alpha\iota]\neq[\iota']$ again. Since the Galois group of $N\subset P$ is trivial, we have $\dim(\alpha\iota,\iota)=0$ and $\dim(\bar{\iota}\alpha\iota,\hat{\xi})=0$. Since $N\subset P$ is 3-supertransitive, the endomorphism $\bar{\iota}\iota'$ contains $\hat{\xi}$ with multiplicity one and we get the statement.

(3) The Galois group $\mathrm{Gal}(M/N)$ is isomorphic to the set of 1 dimensional sectors contained in $\kappa\iota\bar{\iota}\bar{\kappa}$. By a similar computation as above,

$$[\kappa\iota\bar{\iota}\bar{\kappa}] = [\kappa\bar{\kappa}]\oplus[\kappa\xi\bar{\kappa}] = [\mathrm{id}_M]\oplus[\hat{\eta}]\oplus[\kappa\eta\alpha\bar{\kappa}] = [\mathrm{id}_M]\oplus[\hat{\eta}]\oplus[\kappa\alpha\bar{\kappa}]\oplus[\kappa'\alpha\bar{\kappa}].$$

Since the depth of $P\subset M$ is at least 4 due to Lemma 4.1, the endomorphism $\hat{\eta}$ is not an automorphism. If $\kappa'\alpha\bar{\kappa}$ contained an automorphism θ, Frobenius reciprocity would imply $[\kappa'\alpha]=[\theta\kappa]$ and $d(\kappa)=d(\kappa')$, which is a contradiction.

Assume that $\mathrm{Gal}(M/N)$ is not trivial. Then $\kappa\alpha\bar{\kappa}$ contains an automorphism, say, θ. Then Frobenius reciprocity implies $[\theta\kappa]=[\kappa\alpha]$ and

$$[\mathrm{id}_P]\oplus[\alpha\xi] = [\bar{\kappa}\kappa] = [\alpha^{-1}\bar{\kappa}\kappa\alpha] = [\mathrm{id}_P]\oplus[\alpha^{-1}\alpha\xi\alpha] = [\mathrm{id}_P]\oplus[\xi\alpha].$$

Since $[\alpha\xi]=[\xi\alpha^{-1}]$, this shows that ξ^2 contains α^2. If α^2 were outer, we would have $[\alpha^2\iota]=[\iota']$ as $\iota'\bar{\iota}$ would contain α^2. However this contradicts $d(\iota)\neq d(\iota')$. Thus the order of $[\alpha]$ is two. This implies $[\theta^2\kappa]=[\kappa\alpha^2]=[\kappa]$. Since the Galois group of $P\subset M$ is trivial, we get $[\theta^2]=[\mathrm{id}_M]$. Let β be another automorphism contained in $\kappa\alpha\bar{\kappa}$. Then the above argument shows that $[\theta\kappa]=[\beta\kappa]$ and $[\beta^{-1}\theta\kappa]=[\kappa]$, which implies $[\beta^{-1}\theta]=[\mathrm{id}_M]$. Therefore $\mathrm{Gal}(M/N)\cong\mathbb{Z}/2\mathbb{Z}$.

Assume now that the order of $[\alpha]$ is two. Then

$$[\overline{(\kappa\alpha)}\kappa\alpha] = [\alpha^{-1}(\mathrm{id}_P\oplus\eta)\alpha] = [\mathrm{id}_P]\oplus[\alpha^{-1}\eta\alpha] = [\mathrm{id}_P]\oplus[\eta] = [\bar{\kappa}\kappa].$$

Since $H^2(\widehat{P \subset M})$ is trivial, the subfactors $\alpha^{-1}\bar{\kappa}(M)$ and $\bar{\kappa}(M)$ are inner conjugate in P and so there exists an automorphism θ of M such that $[\theta\kappa] = [\kappa\alpha]$. Since the Galois group of $\bar{\kappa}(M) \subset P$ is trivial, we have $[\kappa\alpha] \neq [\kappa]$ and θ is outer. The same computation as above implies that $[\theta]$ is contained in $[\kappa\iota\bar{\iota}\bar{\kappa}]$ and so $\mathrm{Gal}(M/N)$ is not trivial.

The above argument shows that when $\mathrm{Gal}(M/N)$ is isomorphic to $\mathbb{Z}/2\mathbb{Z}$, we have $[\theta\kappa] = [\kappa\alpha] = [\kappa_Q\pi]$ and we can choose a representative θ of $[\theta]$ so that $\theta\kappa = \kappa_Q\pi$ holds. Then $\theta(P) = Q$ and $\theta\kappa\iota = \kappa_Q\pi\iota = \kappa_Q\iota_Q = \kappa\iota$, and so $\mathrm{Gal}(M/N) = \{\mathrm{id}, \theta\}$. □

Corollary 4.5. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *be an irreducible noncommuting quadrilateral of factors such that* $\mathfrak{Q}$ *is (2,2)-supertransitive and the class* $c(\mathfrak{Q})$ *is trivial. Assume that* $\mathfrak{Q}$ *is cocommuting and the Galois group* $\mathrm{Gal}(M/N)$ *is isomorphic to* $\mathbb{Z}/3\mathbb{Z}$*. Then* $[M:P] \geq (5+\sqrt{13})/2$ *and* $\mathfrak{Q}$ *is neither (4,2)-supertransitive nor (2,3)-supertransitive. If* $\mathfrak{Q}$ *is (3,2)-supertransitive, the dual principal graph (to be more precise, the induction-reduction graph for* $M-M$ *and* $M-P$ *bimodules) contains one of the graphs of the Haagerup subfactor.*

Proof. Let $\mathrm{Gal}(M/N) = \{\mathrm{id}_M, \theta, \theta^2 = \theta^{-1}\}$. The proof of Theorem 4.3 shows that $[\theta\hat{\eta}] = [\hat{\eta}\theta^{-1}]$, the endomorphism $\hat{\eta}^2$ does not contain any non-trivial automorphism and

$$[\kappa\iota\bar{\iota}\bar{\kappa}] = [\mathrm{id}_M] \oplus [\theta] \oplus [\hat{\eta}] \oplus [\theta\hat{\eta}] \oplus [\theta^2\hat{\eta}] \oplus [\theta^2\hat{\eta}^2].$$

Note that $[\hat{\eta}]$, $[\theta\hat{\eta}]$, $[\theta^2\hat{\eta}]$ are distinct sectors. Since $\hat{\eta}^2$ contains $\hat{\eta}$, the endomorphism $\kappa\iota\bar{\iota}\bar{\kappa}$ contains $\theta^2\hat{\eta}$ with multiplicity at least 2. Since $[\theta\kappa\iota] = [\kappa\iota]$, the multiplicities of $\hat{\eta}$ and $\theta\hat{\eta}$ in $\kappa\iota\bar{\iota}\bar{\kappa}$ are also at least 2, which implies $\hat{\eta}^2$ contains $\mathrm{id}_M \oplus \hat{\eta} \oplus \theta\hat{\eta} \oplus \theta^2\hat{\eta}$. This shows that $P \subset M$ is not 4-supertransitive and $d(\hat{\eta})^2 \geq 1 + 3d(\hat{\eta})$, which implies $[M:P] = 1 + d(\hat{\eta}) \geq (5+\sqrt{13})/2$. Note that $N \subset P$ is 3-supertransitive if and only if $\dim(\xi, \xi^2) = 1$. Using $[\xi] = [\bar{\kappa}\theta\kappa]$, we get

$$\begin{aligned} \dim(\xi, \xi^2) &= \dim(\bar{\kappa}\theta\kappa, \bar{\kappa}\theta\kappa\bar{\kappa}\theta\kappa) = \dim(\kappa\bar{\kappa}\theta\kappa\bar{\kappa}, \theta\kappa\bar{\kappa}\theta) \\ &= \dim((\mathrm{id}_M \oplus \hat{\eta})\theta(\mathrm{id}_M \oplus \hat{\eta}), \theta(\mathrm{id}_M \oplus \hat{\eta})\theta) \\ &= \dim(\theta \oplus \theta\hat{\eta} \oplus \theta^2\hat{\eta} \oplus \theta^2\hat{\eta}^2, \theta^2 \oplus \hat{\eta}) \\ &= \dim(\theta^2\hat{\eta}^2, \theta^2 \oplus \hat{\eta}) \geq 2, \end{aligned}$$

which shows that $N \subset P$ is not 3-supertransitive.

Assume that $\mathfrak{Q}$ is (3,2)-supertransitive. Then $\kappa'\bar{\kappa}$ contains $\hat{\eta} \oplus \theta\hat{\eta} \oplus \theta^2\hat{\eta}$. and the dual principal graph of $P \subset M$ contains the following graph,



which is one the principal graphs of the Haagerup subfactor [1]. □

In Section 5 and in the Appendix, we will show that there exists a (3,2)-supertransitive quadrilateral of factors in the above class such that $P \subset M$ is the Haagerup subfactor and that such a quadrilateral is unique up to flip conjugacy.

Corollary 4.6. *Let* $\mathfrak{Q} = \begin{matrix} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{matrix}$ *be an irreducible noncommuting quadrilateral of factors. Assume that* $\mathfrak{Q}$ *is cocommuting and (2,3)-supertransitive. Then* $\Theta(P,Q) = \cos^{-1} 1/([P:N]-1)$, $[M:P] = [P:N]-1$, *and the Galois group* $\mathrm{Gal}(M/N)$ *is either trivial or isomorphic to* $\mathbb{Z}/2\mathbb{Z}$ *or* $\mathfrak{S}_3$. *Moreover,*

1. *Two equalities* $[\eta\iota] = [\iota']$ *and* $[\bar{\iota}\bar{\kappa}\kappa\iota] = [\mathrm{id}_M] \oplus [\hat{\xi}] \oplus [\bar{\iota}\iota']$ *hold. In consequence, the endomorphism* $\iota'\bar{\iota}$ *contains* η *with multiplicity one and* $[\iota'\bar{\iota}] = [\eta] \oplus [\eta\xi]$.
2. $\dim(\kappa\iota, \kappa\iota\hat{\xi}) = 2$.
3. *Assume that* $\mathrm{Gal}(M/N)$ *is isomorphic to* $\mathbb{Z}/2\mathbb{Z}$ *and* $\mathrm{Gal}(M/N) = \{\mathrm{id}_M, \theta\}$. *Then*
$$[\kappa\iota\bar{\iota}\bar{\kappa}] = [\mathrm{id}_M] \oplus [\theta] \oplus [\hat{\eta}] \oplus [\theta\hat{\eta}] \oplus [\hat{\eta}\theta] \oplus [\hat{\eta}\theta\hat{\eta}],$$
and $\dim(\kappa\iota\bar{\iota}\bar{\kappa}, \kappa\iota\bar{\iota}\bar{\kappa}) \geq 7$. *The sectors* $[\hat{\eta}]$, $[\theta\hat{\eta}]$, $[\hat{\eta}\theta]$, *and* $[\theta\hat{\eta}\theta]$ *are distinct. The endomorphism* $\hat{\eta}\theta\hat{\eta}$ *contains* $\theta\hat{\eta}\theta$ *with multiplicity one and it does not contain either any automorphism or any of* $\hat{\eta}$, $\theta\hat{\eta}$, $\hat{\eta}\theta$. *The endomorphism* $\hat{\eta}^2$ *does not contain any of* $\theta\hat{\eta}$, $\hat{\eta}\theta$, $\theta\hat{\eta}\theta$.
4. *The endomorphism* ξ^2 *contains* $\bar{\pi}\eta_Q\pi$ *and* $[\bar{\pi}\eta_Q\pi\iota] = [\iota']$ *holds. Assume that* $H^2(\widehat{P \subset M})$ *is trivial. Then the Galois group* $\mathrm{Gal}(M/N)$ *is trivial if and only if* $[\eta] \neq [\bar{\pi}\eta_Q\pi]$. *Assume further that* $\mathrm{Gal}(M/N)$ *is trivial. Then the dual principal graph of* $N \subset P$ *contains* D_6 *in such a way that every endpoint of* D_6 *is also an endpoint of the dual principal graph.*
5. *Assume that* $P \subset M$ *is 3-supertransitive and* $\mathrm{Gal}(M/N)$ *is trivial. Then* $Q \subset M$ *is 3-supertransitive as well and*
$$[\kappa\iota\bar{\iota}\bar{\kappa}] = [\mathrm{id}_M] \oplus [\hat{\eta}] \oplus 2[\kappa_Q\pi\bar{\kappa}] \oplus [\kappa'_Q\bar{\kappa}_Q] = [\mathrm{id}_M] \oplus [\hat{\eta}_Q] \oplus 2[\kappa_Q\pi\bar{\kappa}] \oplus [\kappa'\bar{\kappa}].$$
The sector $[\kappa_Q\pi\bar{\kappa}]$ *is irreducible and self-conjugate and* $[\kappa'] = [\kappa'_Q\pi]$ *holds. Unless the principal graph of* $P \subset M$ *is* A_4, *the inequality* $\dim(\kappa\iota\bar{\iota}\bar{\kappa}, \kappa\iota\bar{\iota}\bar{\kappa}) \geq 8$ *holds.*

Proof. The first part follows from Lemma 3.4, Theorem 4.3, and Corollary 4.5.

(1) Let $d = d(\iota) = [P:N]-1$. Then we have $d(\xi) = d^2-1$ and $d(\iota') = d(d^2-2)$. On the other hand, we have $d(\eta) = [M:P]-1 = [P:N]-2 = d^2-2$. Since $\eta\iota$ contains ι' and $d(\eta\iota) = d(\iota')$, we get $[\eta\iota] = [\iota']$.

(2) Using (1), we get
$$[\bar{\iota}\bar{\kappa}\kappa\iota] = [\bar{\iota}\iota] \oplus [\bar{\iota}\eta\iota] = [\mathrm{id}_N] \oplus [\hat{\xi}] \oplus [\bar{\iota}\iota'].$$
Since $N \subset P$ is 3-supertransitive, $\bar{\iota}\iota'$ contains $\hat{\xi}$ with multiplicity one and $\dim(\kappa\iota, \kappa\iota\hat{\xi}) = \dim(\bar{\iota}\bar{\kappa}\kappa\iota, \hat{\xi}) = 2$.

(3) The proof of Theorem 4.3 shows that $\hat{\eta}\theta\hat{\eta}$ contain no automorphism, $[\theta\kappa] = [\kappa_Q\pi]$, $[\xi] = [\bar{\kappa}\theta\kappa]$, and
$$[\kappa\iota\bar{\iota}\bar{\kappa}] = [\mathrm{id}_M] \oplus [\theta] \oplus [\hat{\eta}] \oplus [\theta\hat{\eta}] \oplus [\hat{\eta}\theta] \oplus [\hat{\eta}\theta\hat{\eta}].$$

If $[\theta\hat{\eta}] = [\hat{\eta}]$, we would have $[\hat{\eta}\theta\hat{\eta}] = [\hat{\eta}^2]$, which contains id_M. Thus $[\theta\hat{\eta}] \neq [\hat{\eta}]$. In the same way, we get $[\hat{\eta}\theta] \neq [\hat{\eta}]$ and $[\theta\hat{\eta}] \neq [\theta\hat{\eta}\theta] \neq [\hat{\eta}\theta]$. If $[\hat{\eta}] = [\theta\hat{\eta}\theta]$, we would have

$$[\kappa\bar{\kappa}] = [\mathrm{id}_M] \oplus [\hat{\eta}] = [\mathrm{id}_M] \oplus [\theta\hat{\eta}\theta] = [\theta\kappa\bar{\kappa}\theta] = [\kappa_Q\pi\bar{\pi}\bar{\kappa}_Q] = [\kappa_Q\bar{\kappa}_Q],$$

which contradicts the assumption that $\mathfrak{Q}$ is cocommuting thanks to Theorem 3.10,(1). Thus $[\hat{\eta}] \neq [\theta\hat{\eta}\theta]$ and $[\theta\hat{\eta}] \neq [\hat{\eta}\theta]$. Since $N \subset P$ is 3-supertransitive, we have $\dim(\xi, \xi^2) = 1$, and

$$\begin{aligned} 1 &= \dim(\xi,\xi^2) = \dim(\bar{\kappa}\theta\kappa, \bar{\kappa}\theta\kappa\bar{\kappa}\theta\kappa) = \dim(\kappa\bar{\kappa}\theta\kappa\bar{\kappa}, \theta\kappa\bar{\kappa}\theta) \\ &= \dim((\mathrm{id}_M \oplus \hat{\eta})\theta(\mathrm{id}_M \oplus \hat{\eta}), \theta(\mathrm{id}_M \oplus \hat{\eta})\theta) \\ &= \dim(\theta \oplus \theta\hat{\eta} \oplus \hat{\eta}\theta \oplus \hat{\eta}\theta\hat{\eta}, \mathrm{id}_M \oplus \theta\hat{\eta}\theta) \\ &= \dim(\hat{\eta}\theta\hat{\eta}, \theta\hat{\eta}\theta), \end{aligned}$$

which means that $\hat{\eta}\theta\hat{\eta}$ contains $\theta\hat{\eta}\theta$ with multiplicity one. Since $[\theta\kappa\iota] = [\kappa\iota]$, the multiplicities of $\hat{\eta}$, $\theta\hat{\eta}$, $\hat{\eta}\theta$ and $\theta\hat{\eta}\theta$ in $\kappa\iota\bar{\iota}\bar{\kappa}$ should be the same, which is the same as $\hat{\eta}\theta\hat{\eta}$ contains $\theta\hat{\eta}\theta$ with multiplicity one. Thus $\hat{\eta}\theta\hat{\eta}$ does not contain any of $\hat{\eta}$, $\theta\hat{\eta}$, $\hat{\eta}\theta$. Thanks to Frobenius reciprocity we conclude that $\hat{\eta}^2$ does not contain any of $\theta\hat{\eta}$, $\hat{\eta}\theta$, $\theta\hat{\eta}\theta$.

(4) Since $[\xi] = [\bar{\kappa}\kappa_Q\pi] = [\bar{\pi}\bar{\kappa}_Q\kappa]$, we have

$$[\xi^2] = [\bar{\pi}\bar{\kappa}_Q\kappa\bar{\kappa}\kappa_Q\pi] = [\bar{\pi}\bar{\kappa}_Q\kappa_Q\pi] \oplus [\bar{\pi}\bar{\kappa}_Q\hat{\eta}\kappa_Q\pi],$$

which contains $\bar{\pi}\eta_Q\pi$. Since $N \subset P$ is 3-supertransitive and $\bar{\pi}\eta_Q\pi$ (which is not equivalent to id) is contained in ξ^2, the morphism $\iota'\bar{\iota}$ contains $\bar{\pi}\eta_Q\pi$ and so $\bar{\pi}\eta_Q\pi\iota$ contains ι'. Since $d(\iota') = d(\bar{\pi}\eta_Q\pi\iota)$, we conclude $[\bar{\pi}\eta_Q\pi\iota] = [\iota']$.

The proof of Theorem 4.3 shows that $\mathrm{Gal}(M/N)$ is non-trivial if and only if there exists and automorphism θ of M such that $[\theta\kappa] = [\kappa_Q\pi]$. Assume first that such an automorphism θ exists. Then $[\bar{\kappa}\kappa] = [\bar{\pi}\bar{\kappa}_Q\kappa_Q\pi]$ and $[\eta] = [\bar{\pi}\eta_Q\pi]$ holds. Assume conversely that $[\eta] = [\bar{\pi}\eta_Q\pi]$ holds. Then

$$[\bar{\kappa}\kappa] = [\mathrm{id}_P] \oplus [\eta] = [\bar{\pi}(\mathrm{id}_Q \oplus \eta_Q)\pi] = [\bar{\pi}\bar{\kappa}_Q\kappa_Q\pi].$$

Since $H^2(\widehat{P \subset M})$ is trivial, the subfactors $\bar{\kappa}(M)$ and $\bar{\pi}\bar{\kappa}_Q(M)$ are inner conjugate in P and there exists an automorphism θ of M such that $[\theta\kappa] = [\kappa_Q\pi]$. Therefore the Galois group $\mathrm{Gal}(M/N)$ is trivial if and only if $[\eta] \neq [\bar{\pi}\eta_Q\pi]$.

Now we assume that $\mathrm{Gal}(M/N)$ is trivial. Then the dual principal graph of $N \subset P$ is as follows:



which contains D_6.

(5) Since $[\eta] \neq [\bar{\pi}\eta_Q\pi]$,

$$\dim(\kappa_Q\pi\bar{\kappa}, \kappa_Q\pi\bar{\kappa}) = \dim(\bar{\pi}\bar{\kappa}_Q\kappa_Q\pi, \bar{\kappa}\kappa) = \dim(\mathrm{id}_P \oplus \bar{\pi}\eta_Q\pi, \mathrm{id}_P \oplus \eta) = 1,$$

and $\kappa_Q\pi\bar{\kappa}$ is irreducible. Since $[\xi] = [\bar{\kappa}\kappa_Q\pi]$ is self-conjugate,

$$1 = \dim(\bar{\kappa}\kappa_Q\pi, \bar{\pi}\bar{\kappa}_Q\kappa) = \dim(\kappa_Q\pi\bar{\kappa}, \kappa\bar{\pi}\bar{\kappa}_Q),$$

and $\kappa_Q\pi\bar{\kappa}$ is self-conjugate. This and the proof of Theorem 4.3 imply

$$\begin{aligned}[][\kappa\iota\bar{\iota}\bar{\kappa}] &= [\mathrm{id}_M] \oplus [\hat{\eta}] \oplus [\kappa_Q\pi\bar{\kappa}] \oplus [\hat{\eta}\kappa_Q\pi\bar{\kappa}] = [\mathrm{id}_M] \oplus [\hat{\eta}] \oplus [\kappa_Q\pi\bar{\kappa}] \oplus [\hat{\eta}\kappa\bar{\pi}\bar{\kappa}_Q] \\ &= [\mathrm{id}_M] \oplus [\hat{\eta}] \oplus 2[\kappa_Q\pi\bar{\kappa}] \oplus [\kappa'\bar{\pi}\bar{\kappa}_Q].\end{aligned}$$

On the other hand, replacing P by Q, we get

$$[\kappa\iota\bar{\iota}\bar{\kappa}] = [\mathrm{id}_M] \oplus [\hat{\eta}_Q] \oplus [\kappa_Q\pi\bar{\kappa}] \oplus [\hat{\eta}_Q\kappa_Q\pi\bar{\kappa}].$$

Since $[\hat{\eta}] \neq [\hat{\eta}_Q]$, the endomorphism $\hat{\eta}_Q\kappa_Q\pi\bar{\kappa}$ contains $\hat{\eta}$ and

$$1 \leq \dim(\hat{\eta}_Q\kappa_Q\pi\bar{\kappa}, \hat{\eta}) = \dim(\hat{\eta}_Q\kappa_Q\pi, \hat{\eta}\kappa) = \dim(\hat{\eta}_Q\kappa_Q\pi, \kappa \oplus \kappa').$$

If $\hat{\eta}_Q\kappa_Q\pi$ contained κ, we would have $[\hat{\eta}_Q] = [\kappa\bar{\pi}\bar{\kappa}_Q]$, which contradicts $d(\hat{\eta}_Q) \neq d(\kappa\bar{\pi}\bar{\kappa}_Q)$. Therefore $\hat{\eta}_Q\kappa_Q\pi$ contains $\kappa_Q\pi$ and κ', and so $[\hat{\eta}_Q\kappa_Q\pi] = [\kappa_Q\pi] \oplus [\kappa']$ as we have $d(\hat{\eta}_Q\kappa_Q\pi) = d(\kappa_Q\pi) + d(\kappa')$. This shows that $Q \subset M$ is 3-supertransitive and $[\kappa'_Q\pi] = [\kappa']$, which finishes the proof. ☐

About uniqueness, we have the following:

Theorem 4.7. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *and* $\tilde{\mathfrak{Q}} = \begin{array}{ccc} \tilde{P} & \subset & \tilde{M} \\ \cup & & \cup \\ \tilde{N} & \subset & \tilde{Q} \end{array}$ *be irreducible noncommuting and noncocommuting quadrilaterals of factors such that* $\mathfrak{Q}$ *and* $\tilde{\mathfrak{Q}}$ *are (3,3)-supertransitive. Assume that there exists an isomorphism* Φ *from* P *onto* $\tilde{P}$ *such that* $\Phi(N) = \tilde{N}$. *If there exists no sector* σ *contained in* ξ^2 *such that* $[\xi] \neq [\sigma]$, $[\eta] \neq [\sigma]$, *and* $d(\sigma) = d(\eta)$, *then* $\mathfrak{Q}$ *and* $\tilde{\mathfrak{Q}}$ *are conjugate.*

Proof. Since $P \subset M$ and $\tilde{P} \subset \tilde{M}$ are 3-supertransitive, $H^2(\widehat{P \subset M})$ and $H^2(\widehat{\tilde{P} \subset \tilde{M}})$ are trivial thanks to Lemma 3.2. Note that we have $[\iota_{\tilde{P},\tilde{N}}\overline{\iota_{\tilde{P},\tilde{N}}}] = [\Phi\iota_{P,N}\overline{\iota_{P,N}}\Phi^{-1}]$. Therefore Lemma 4.1 implies $[\overline{\iota_{\tilde{M},\tilde{P}}}\iota_{\tilde{M},\tilde{P}}] = [\Phi\overline{\iota_{M,P}}\iota_{M,P}\Phi^{-1}]$, which shows that Φ extends to an isomorphism Ψ from M onto $\tilde{M}$ as $H^2(\widehat{P \subset M})$ and $H^2(\widehat{\tilde{P} \subset \tilde{M}})$ are trivial. Lemma 3.14 and Corollary 4.4 show $\Psi(Q) = \tilde{Q}$. ☐

Theorem 4.8. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *and* $\tilde{\mathfrak{Q}} = \begin{array}{ccc} \tilde{P} & \subset & \tilde{M} \\ \cup & & \cup \\ \tilde{N} & \subset & \tilde{Q} \end{array}$ *be irreducible noncommuting and cocommuting quadrilaterals of factors such that* $H^2(\widehat{P \subset M})$ *and* $H^2(\widehat{\tilde{P} \subset \tilde{M}})$ *are trivial and* $\mathfrak{Q}$ *and* $\tilde{\mathfrak{Q}}$ *are (2,3)-supertransitive. Assume that there exists an isomorphism* Φ *from* P *onto* $\tilde{P}$ *such that* $\Phi(N) = \tilde{N}$. *Then*

1. *If* $\mathrm{Gal}(M/N) \cong \mathbb{Z}/2\mathbb{Z}$ *and* η *is the only sector* σ *contained in* ξ^2 *such that* $[\sigma\iota] = [\iota']$, *then* $\mathfrak{Q}$ *and* $\tilde{\mathfrak{Q}}$ *are conjugate.*
2. *If* $\mathrm{Gal}(M/N)$ *and* $\mathrm{Gal}(\tilde{M}/\tilde{N})$ *are trivial and there exists only two sectors* σ *contained in* ξ^2 *such that* $[\sigma\iota] = [\iota']$, *then* $\mathfrak{Q}$ *and* $\tilde{\mathfrak{Q}}$ *are flip conjugate.*

Proof. The proof of (1) is the same as that of Theorem 4.7. In (2),

$$[\overline{\iota_{\tilde{M},\tilde{P}}}\iota_{\tilde{M},\tilde{P}}] = [\Phi(\mathrm{id}_P \oplus \bar{\pi}\eta_Q\pi)\Phi^{-1}] = [\Phi\pi^{-1}\overline{\iota_{M,Q}}\iota_{M,Q}\pi\Phi^{-1}]$$

may occur instead of $[\overline{\iota_{\tilde{M},\tilde{P}}}\iota_{\tilde{M},\tilde{P}}] = [\Phi\overline{\iota_{M,P}}\iota_{M,P}\Phi^{-1}]$. In this case, the two quadrilaterals are flip conjugate. ☐

We can improve the assumption of Theorem 2.7.

Corollary 4.9. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *be an irreducible noncommuting quadrilateral of factors. Then*

1. *If $\mathfrak{Q}$ is not cocommuting and (3,4)-supertransitive, then all the elementary subfactors are isomorphic to A_7 subfactors. There exists a unique such quadrilateral up to isomorphism when the factors are hyperfinite II_1 factors.*
2. *If $\mathfrak{Q}$ is cocommuting and (2,4)-supertransitive, then there exists an outer action of $\mathfrak{S}_3$ on M such that N is the fixed point algebra of the action. There exists a unique such quadrilateral up to isomorphism when the factors are hyperfinite II_1 factors.*

Proof. Since $N \subset P$ is 4-supertransitive, we have $[\xi^2] = [\mathrm{id}_P] \oplus [\xi] \oplus [\eta]$.

(1) Since $\mathfrak{Q}$ is not cocommuting, there exists an automorphism θ of P such that $[\eta] = [\theta\xi]$ and so we get $d(\xi)^2 = 1 + 2d(\xi)$. This shows that the principal graphs of $N \subset P$ and $P \subset Q$ are A_7. The uniqueness follows from Theorem 4.7.

(2) Since $\mathfrak{Q}$ is cocommuting, we have $[\eta\iota] = [\iota']$ and so the principal graph of $N \subset P$ is A_5. The rest of the proof has already been stated in Remark 4.2 except for uniqueness, which follows from the uniqueness of outer actions of finite groups by Jones [29] and the Galois correspondence. ☐

5. CLASSIFICATION I

According to our discussions in the last section, we consider the following four classes of quadrilaterals $\mathfrak{Q}$ of factors. We assume that every quadrilateral $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ of factors appearing in this section is irreducible and noncommuting.

Class I: $\mathfrak{Q}$ is noncocommuting and (3,3)-supertransitive.

Class II: $\mathfrak{Q}$ is cocommuting and (2,3)-supertransitive. The Galois group $\mathrm{Gal}(M/N)$ is trivial.

Class III: $\mathfrak{Q}$ is cocommuting and (2,3)-supertransitive. The Galois group $\mathrm{Gal}(M/N)$ is isomorphic to $\mathbb{Z}/2\mathbb{Z}$.

Class IV: $\mathfrak{Q}$ is cocommuting and (3,2)-supertransitive. The Class $c(\mathfrak{Q})$ is trivial. The Galois group $\mathrm{Gal}(M/N)$ is isomorphic to $\mathbb{Z}/3\mathbb{Z}$.

For each class, we show that there exists an example of a quadrilateral and we seek an example with maximal supertransitivity.

We keep using the notation of sectors such as ι, κ, etc. as in the previous section, We often use explicit formulae of fusion rules of subfactors and the reader is refered to [17] for them.

5.1. **Class I.** The structure of quadrilaterals in Class I is relatively easy to describe.

Theorem 5.1. *Let* $N \subset P$ *be an irreducible 3-supertransitive inclusion of factors. Let* $\iota = \iota_{P,N}$ *and let* $[\iota_{P,N}\overline{\iota_{P,N}}] = [\mathrm{id}_P] \oplus [\xi]$ *and* $[\xi\iota] = [\iota] \oplus [\iota']$ *be the irreducible decomposition. Assume that there exists an outer automorphism* $\alpha \in \mathrm{Aut}(P)$ *such that* $[\xi] \neq [\alpha\xi] = [\xi\alpha^{-1}]$ *and* ξ^2 *contains* $\alpha\xi$*. If* $\mathrm{id}_P \oplus \alpha\xi$ *has a* Q*-system, then there exists a unique irreducible noncommuting quadrilateral* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *in Class I such that* $[\eta] = [\alpha\xi]$*.*

Proof. Assume that there exists a Q-system for $\mathrm{id}_P \oplus \alpha\xi$. We claim that a Q-system with $\mathrm{id}_P \oplus \alpha\xi$ is unique up to equivalence. For this, it suffices to show $\dim(\alpha\xi, (\alpha\xi)^2) = 1$ thanks to Lemma 3.5. As in the proof of Corollary 4.5, we have $[\alpha\iota'] = [\iota']$, $[\alpha\iota] \neq [\iota]$, and $d(\iota) \neq d(\iota')$. Thus

$$\begin{aligned} \dim(\alpha\xi, (\alpha\xi)^2) &= \dim(\alpha\xi, \xi^2) = \dim(\alpha\xi, \xi\iota\bar{\iota}) - \dim(\alpha\xi, \xi) = \dim(\alpha\xi\iota, \xi\iota) \\ &= \dim(\alpha\iota \oplus \alpha\iota', \iota \oplus \iota') = 1, \end{aligned}$$

which shows the claim. The claim implies that there exists a unique factor M containing N such that $[\bar{\kappa}\kappa] = [\mathrm{id}_P] \oplus [\alpha\xi]$ where $\kappa = \iota_{M,P}$. Since $[\xi] \neq [\alpha\xi]$, the inclusion $N \subset M$ is irreducible.

We next claim that $\kappa\alpha\iota$ is irreducible and $[\kappa\alpha\iota] = [\kappa\iota]$. Indeed, we have

$$\dim(\kappa\alpha\iota, \kappa\alpha\iota) = \dim(\bar{\kappa}\kappa\alpha, \alpha\iota\bar{\iota}) = \dim(\alpha \oplus \alpha\xi\alpha, \alpha \oplus \alpha\xi) = 1,$$

$$\dim(\kappa\alpha\iota, \kappa\iota) = \dim(\bar{\kappa}\kappa\alpha, \iota\bar{\iota}) = \dim(\alpha \oplus \alpha\xi\alpha, \mathrm{id}_P \oplus \xi) = 1,$$

which show the claim. The claim shows that there exists a unitary $v \in M$ such that $v\alpha(x)v^* = x$ holds for every $x \in N$. We set $\pi(y) = v\alpha(y)v^*$ for $y \in P$ and set $Q = \pi(P)$. By construction Q is an intermediate subfactor between N and M such that $\iota_{Q,N} = \pi\iota$, $[\overline{\iota_{Q,N}}\iota_{Q,N}] = [\bar{\iota}\iota]$, and $[M : P] = [M : Q]$.

Suppose $P = Q$. Then π would be an automorphism of P, which satisfies $[\kappa\pi] = [\kappa\alpha]$. Frobenius reciprocity implies that $\pi\alpha^{-1}$ is contained in $\bar{\kappa}\kappa = \mathrm{id}_P \oplus \alpha\xi$ and so $[\pi] = [\alpha]$. Thus there exists a unitary u in P such that $v\alpha(y)v^* = u\alpha(y)u^*$ for all $y \in P$ and so $u^*v \in P' \cap M = \mathbb{C}$. This shows that $v \in P$ and $[\alpha\iota] = [\iota]$, which is a contradiction. Therefore $P \neq Q$.

Since $N \subset P$ is 3-supertransitive, it has no intermediate subfactor and $P \cap Q = N$. Since $\dim(\alpha\xi, (\alpha\xi)^2) = 1$, the inclusion $P \subset M$ is 3-supertransitive too and M is generated by P and Q. Therefore $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ is an irreducible quadrilateral of factors. Theorem 3.10,(1) shows that $\mathfrak{Q}$ is noncommuting. Since P and Q are inner conjugate in M, Theorem 3.10,(1) applied to the dual quadrilateral $\hat{\mathfrak{Q}}$ shows that $\mathfrak{Q}$ is noncocommuting. □

The above theorem shows that quadrilaterals in Class I is completely determined by the subfactor $N \subset P$ and α.

Jones and the first-named author [13] showed that there exists a unique quadrilateral of the hyperfinite II_1 factors such that all the elementary subfactors are the the A_7 subfactor. It is easy to show that this is the only quadrilateral in Class I satisfying $[P : N] < 4$. Other than this example, we know two subfactors satisfying the assumption of Theorem 5.1, namely, the $E_7^{(1)}$ subfactor and the Haagerup subfactors.

Let $N \subset P$ be the $E_7^{(1)}$ subfactor. Then the dual principal graph is as follows:



Note that the category of $P - P$ bimodules for the $E_7^{(1)}$ subfactor is isomorphic to the category $\hat{\mathfrak{A}}_4$ of the unitary representations of the alternating group $\mathfrak{A}_4$. It is observed in [24, Corollary 4.2] that there exists an automorphism of the category $\hat{\mathfrak{A}}_4$ that flips the two representations corresponding to ξ and $\alpha\xi$. Thus $\mathrm{id}_P \oplus \alpha\xi$ has a Q-system and gives rise to an irreducible noncommuting quadrilateral in Class I whose elementary subfactors are the $E_7^{(1)}$ subfactor. Note that this quadrilateral satisfies the assumption of Theorem 4.7 and such a quadrilateral is unique.

Since it is easy to see that no other subfactors of index less than or equal to 4 fit into the statement of Theorem 4.3,(1), we get the following:

Theorem 5.2. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *be a quadrilateral of factors in Class I such that* $[M : P] \leq 4$. *Then one of the following two cases occurs:*

1. *The principal graphs of all the elementary subfactors are* A_7.
2. *The principal graphs of all the elementary subfactors are* $E_7^{(1)}$.

In each case, such a quadrilateral of hyperfinite II_1 *factors exists and is unique up to conjugacy.*

Let $N \subset P$ be the Haagerup subfactor [1]. Since the Haagerup subfactor is not self-dual, we specify the dual principal graph (the induction-reduction graph of the $P - P$ and $P - N$ bimodules) as below.



It is known that $\alpha^3 = \mathrm{id}_P$ and $\alpha\xi = \xi\alpha^{-1}$ holds. Since $\mathrm{id}_P \oplus \xi$ has a Q-system, so does $\alpha^{-1}(\mathrm{id}_P \oplus \xi)\alpha = \mathrm{id}_P \oplus \alpha\xi$. Thus there exists an irreducible noncommuting quadrilateral of factors in Class I whose elementary subfactors are the Haagerup subfactor. Although the quadrilateral arising from $N \subset P$ does not satisfy the assumption of Theorem 4.7, we have $[\kappa\alpha] = [\kappa_Q \pi]$ and so $[\bar{\pi}\eta_Q\pi] = [\alpha^{-1}\alpha\xi\alpha] = [\alpha^2\xi]$. Thus the same proof of Theorem 4.8,(2) works and we can show uniqueness of such a quadrilateral up to flip conjugacy.

We don't know if there are infinitely many mutually non-conjugate quadrilaterals of the hyperfinite II_1 factors in Class I.

5.2. **Class II.** The following two are the main theorems of this subsection:

Theorem 5.3. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *be a quadrilateral of factors in Class II such that* $\mathfrak{Q}$ *is (5,3)-supertransitive. Then*

1. *If* $\mathfrak{Q}$ *is (6,3)-supertransitive, the principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* A_4 *and those of* $N \subset P$ *and* $N \subset Q$ *are* D_6.
2. *If* $\mathfrak{Q}$ *is not (6,3)-supertransitive, the principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* $E_8^{(1)}$.

In each case, such a quadrilateral of the hyperfinite II_1 *factors exists and is unique up to conjugacy.*

Theorem 5.4. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *be a quadrilateral of factors in Class II such that* $[M : P] \leq 4$. *Then one of the following holds:*

1. *The principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* A_4 *and those of* $N \subset P$ *and* $N \subset Q$ *are* D_6. *Such a quadrilateral of the hyperfinite* II_1 *factors exists and is unique up to conjugacy.*
2. *The principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* E_6 *and the dual principal graphs of* $N \subset P$ *and* $N \subset Q$ *are as below.*



   Such a quadrilateral of the hyperfinite II_1 *factors exists.*

3. *The principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* $E_8^{(1)}$. *Such a quadrilateral of the hyperfinite* II_1 *factors exists and is unique up to conjugacy.*

In the sequel, the dimension of the space $(\kappa\iota\bar{\iota}\bar{\kappa}, \kappa\iota\bar{\iota}\bar{\kappa})$ often gives important information about a quadrilateral and we give useful formulae about it first.

Lemma 5.5. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *be an irreducible noncommuting quadrilateral of factors such that* $\mathfrak{Q}$ *is cocommuting and (2,3)-supertransitive. Then* $\iota'\bar{\iota}$ *contains* ξ

and η with multiplicity one, the equality $[\iota'\bar{\iota}] = [\eta] \oplus [\eta\xi]$ holds, and

$$\dim(\kappa\iota\bar{\iota}\bar{\kappa}, \kappa\iota\bar{\iota}\bar{\kappa}) = 5 + \dim(\eta\xi, \eta\xi).$$

If moreover $P \subset M$ is 3-supertransitive, then $[\eta\xi] = [\xi] \oplus [\bar{\kappa}'\kappa_Q\pi]$ and

$$\dim(\kappa\iota\bar{\iota}\bar{\kappa}, \kappa\iota\bar{\iota}\bar{\kappa}) = 6 + \dim(\bar{\kappa}'\kappa_Q\pi, \bar{\kappa}'\kappa_Q\pi) = 6 + \dim(\kappa'\bar{\kappa}', \kappa_Q\bar{\kappa}_Q).$$

Proof. By Corollary 4.6,(1) and Frobenius reciprocity, we have

$$\begin{aligned}\dim(\kappa\iota\bar{\iota}\bar{\kappa}, \kappa\iota\bar{\iota}\bar{\kappa}) &= \dim(\bar{\iota}\bar{\kappa}\kappa\iota, \bar{\iota}\bar{\kappa}\kappa\iota) = \dim(\mathrm{id}_N \oplus \hat{\xi} \oplus \bar{\iota}\iota', \mathrm{id}_N \oplus \hat{\xi} \oplus \bar{\iota}\iota') \\ &= 2 + 2\dim(\iota\hat{\xi}, \iota') + \dim(\bar{\iota}\iota', \bar{\iota}\iota') = 4 + \dim(\bar{\iota}\iota', \bar{\iota}\iota').\end{aligned}$$

Since $[\bar{\iota}\iota\bar{\iota}\iota] = 2[\bar{\iota}\iota] \oplus [\bar{\iota}\iota']$, the sector $[\bar{\iota}\iota']$ is self-conjugate, and so is $[\iota'\bar{\iota}]$ for a similar reason. Thus we get

$$\dim(\bar{\iota}\iota', \bar{\iota}\iota') = \dim(\bar{\iota}\iota', \bar{\iota}'\iota) = \dim(\iota'\bar{\iota}, \iota\bar{\iota}') = \dim(\iota'\bar{\iota}, \iota'\bar{\iota}).$$

The statement follows from Corollary 4.6,(1) now. $\square$

In view of Corollary 4.6,(4), a canonical candidate of a quadrilateral in Class II with the smallest index is that with the D_6 subfactor for $N \subset P$.

Proposition 5.6. *Let $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ be a quadrilateral in Class II or Class III such that the principal graph of $P \subset M$ is A_4. Then $\mathfrak{Q}$ is in Class II and the principal graph of $N \subset P$ is D_6. Such a quadrilateral of the hyperfinite II_1 factors exists and is unique up to conjugacy.*

Proof. From Theorem 4.3, we see $[P : N] = [M : P] + 1 = (5 + \sqrt{5})/2 = 4\cos^2(\pi/10)$, which shows that the principal graph of $N \subset P$ is either A_9 or D_6. Thanks to Theorem 2.7, the former never occurs. If $\mathfrak{Q}$ were in Class III, Corollary 4.6,(3) would imply that $\hat{\eta}\theta\hat{\eta}$ does not contain any automorphism and it contains $\theta\hat{\eta}\theta$. However, this contradicts $d(\hat{\eta}\theta\hat{\eta}) - d(\theta\hat{\eta}\theta) = d(\hat{\eta})^2 - d(\hat{\eta}) = 1$ and $\mathfrak{Q}$ is in Class II.

In the case of the hyperfinite II_1 factors, uniqueness up to flip conjugacy follows from Theorem 4.8,(2) and the uniqueness of the D_6 subfactor. Let $[\xi^2] = [\mathrm{id}_P] \oplus [\xi] \oplus [\xi'] \oplus [\xi'']$ be the irreducible decomposition of ξ^2. To prove uniqueness up to conjugacy, it suffices to show that there exists an automorphism α of P such that $\alpha(N) = N$ and $[\alpha\xi'\alpha^{-1}] = [\xi'']$ (see the proof of Theorem 4.8,(2)). It is shown in [34] that there exists an automorphism β of period two on the A_9 subfactor $\tilde{N} \subset \tilde{P}$ such that

$$(N \subset P) \cong (\tilde{N} \rtimes_\beta \mathbb{Z}/2\mathbb{Z} \subset \tilde{P} \rtimes_\beta \mathbb{Z}/2\mathbb{Z}).$$

The dual action $\hat{\beta}$ of β acts on the higher relative commutants of $N \subset P$ non-trivially. It is routine work to show that $\hat{\beta}$ does the right job.

Finally we show existence. Let $\mathcal{N} \subset \mathcal{M}$ be an inclusion of factors whose principal graph is D_6. We use the following parameterization of sectors associated with $\mathcal{N} \subset$

$\mathcal{M}$.


$\mathcal{M}-\mathcal{M}$
$\mathrm{id}_\mathcal{M}$
ρ
ρ_1
ρ_2
$\mathcal{M}-\mathcal{N}$
ν
μ
$\mathcal{N}-\mathcal{N}$
$\mathrm{id}_\mathcal{N}$
σ
σ_1
σ_2

Recall the fusion rule for the D_6 subfactors [17, Proposition 3.10]:

$$[\rho^2]=[\mathrm{id}_\mathcal{M}]\oplus[\rho]\oplus[\rho_1]\oplus[\rho_2],\quad [\rho_1\rho_2]=[\rho_2\rho_1]=[\rho],$$
$$[\rho_1^2]=[\mathrm{id}_\mathcal{M}]\oplus[\rho_1],\quad [\rho_2^2]=[\mathrm{id}_\mathcal{M}]\oplus[\rho_2],$$
$$[\mu]=[\rho_1\nu]=[\rho_2\nu]=[\nu\sigma_1]=[\nu\sigma_2],$$

where $\nu=\iota_{\mathcal{M},\mathcal{N}}$. We can take representatives ρ_1 and ρ_2 satisfying $\mu=\rho_1\nu=\rho_2\nu$. We set $M=\mathcal{M}$, $P=\rho_1(\mathcal{M})$, $Q=\rho_2(\mathcal{M})$, and $N=\mu(\mathcal{N})=\rho_1(\mathcal{N})=\rho_2(\mathcal{N})$. Then $d(\rho_1)=d(\rho_2)=(1+\sqrt{5})/2$, which means $[M:P]=[M:Q]=4\cos^2\pi/5$, and so the principal graphs of $P\subset M$ and $Q\subset M$ are A_4. By construction, the principal graphs of $N\subset P$ and $N\subset Q$ are D_6. Since μ is irreducible, the inclusion $N\subset M$ is irreducible and $\mathfrak{Q}=\begin{matrix}P&\subset&M\\ \cup&&\cup\\ N&\subset&Q\end{matrix}$ is an irreducible quadrilateral. Since $[\rho_1^2]=[\mathrm{id}_\mathcal{M}]\oplus[\rho_1]$ and $[\rho_2^2]=[\mathrm{id}_\mathcal{M}]\oplus[\rho_2]$, we have $[\iota_{M,P}\overline{\iota_{M,P}}]=[\mathrm{id}_M]\oplus[\rho_1]$ and $[\iota_{M,Q}\overline{\iota_{M,Q}}]=[\mathrm{id}_M]\oplus[\rho_2]$, which implies that $\mathfrak{Q}$ is cocommuting. Therefore $[M:P]\neq[P:N]$ shows that the quadrilateral $\mathfrak{Q}$ is not commuting. □

Remark 5.7. The above construction shows that the principal graph and the dual principal graph of $N\subset M$ are the same, which can be easily computed from

$$[\mu\bar{\mu}]=[\rho_1\nu\bar{\nu}\rho_1]=[\rho_1(\mathrm{id}_M\oplus\rho)\rho_1]=[\mathrm{id}_M]\oplus[\rho_1]\oplus[\rho_2]\oplus2[\rho],$$
$$[\rho_1\mu]=[\rho_1^2\nu]=[\nu]\oplus[\rho_1\nu]=[\nu]\oplus[\mu],$$
$$[\rho_2\mu]=[\rho_2^2\nu]=[\nu]\oplus[\rho_2\nu]=[\nu]\oplus[\mu],$$
$$[\rho\mu]=[\rho\rho_1\nu]=[\rho\nu]\oplus[\rho_2\nu]=[\nu]\oplus2[\mu],$$
$$[\nu\bar{\mu}]=[\nu\bar{\nu}\rho_1]=[\rho_1]\oplus[\rho\rho_1]=[\rho_1]\oplus[\rho_2]\oplus[\rho].$$

By symmetry of μ and $\bar{\mu}$, there exist two more intermediate subfactors L and R between N and M such that the principal graphs of $L\subset M$ and $R\subset M$ are D_6 and those of $N\subset L$ and $N\subset R$ are A_4. Since $\iota_{M,L}\overline{\iota_{M,L}}$ and $\iota_{M,R}\overline{\iota_{M,R}}$ are contained in $\mu\bar{\mu}$, we see that they are equivalent to $\mathrm{id}_M\oplus\rho$. There is no intermediate subfactor other than P,Q,L,R. Indeed, if S is an intermediate subfactor, the endomorphism $\iota_{M,S}\overline{\iota_{M,S}}$ is contained in $\mu\bar{\mu}$. Computation of the indices $[M:S]$ and $[S:N]$ shows that all the possibilities of $\iota_{M,S}\overline{\iota_{M,S}}$ are exhausted by P,Q,L,R. Thus Lemma 3.14 applied to the dual quadrilateral implies the claim.

Proposition 5.8. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *be a quadrilateral in Class II or Class III such that the principal graph of* $P \subset M$ *is* A_5*. Then* $\mathfrak{Q}$ *is in Class III and the principal graph of* $N \subset P$ *is* $E_7^{(1)}$*. When* M *is the hyperfinite* II_1 *factor, such a quadrilateral exists and is unique up to conjugacy.*

Proof. Since the principal graph of $P \subset M$ is A_5, there exists an outer automorphism of period two $\alpha \in \mathrm{Aut}(P)$ such that $[\kappa\eta] = [\kappa] \oplus [\kappa\alpha]$. Since $[\xi] = [\bar{\kappa}\kappa_Q\pi]$, we get

$$[\eta\xi] = [\eta\bar{\kappa}\kappa_Q\pi] = [\bar{\kappa}\kappa_Q\pi] \oplus [\alpha\bar{\kappa}\kappa_Q\pi] = [\xi] \oplus [\alpha\xi].$$

Therefore Corollary 4.6,(1) shows that the dual principal graph of $N \subset P$ is



which is $E_7^{(1)}$. Existence of such a quadrilateral follows from Example 2.6 with $G = \mathfrak{S}_{\{1,2,3,4\}}$ and uniqueness follows from Theorem 4.8. Corollary 4.6,(4) shows that $\mathfrak{Q}$ is in Class III. $\square$

Proposition 5.9. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *be a quadrilateral in Class II or Class III such that the principal graph of* $P \subset M$ *is* E_6*. Then* $\mathfrak{Q}$ *is in Class II and the dual principal graph of* $N \subset P$ *is as in Theorem 5.4 and* $[P : N] = 3 + \sqrt{3}$*. There exists a quadrilateral of the hyperfinite* II_1 *factors satisfying the above property, which arises from the GHJ pair* [13] *(see Example 7.10) for* E_6 *with* $*$ *given by the vertex with the smallest entry of the Perron-Frobenius eigenvector.*

Proof. From Theorem 4.3, we see $[P : N] = [M : P] + 1 = 3 + \sqrt{3}$. We recall the fusion rule for the E_6 subfactors [17, p.968]:

$$[\eta^2] = [\mathrm{id}_P] \oplus [\alpha] \oplus 2[\eta], \quad [\alpha\eta] = [\eta\alpha] = [\eta], \quad [\alpha^2] = [\mathrm{id}_P],$$

$$[\kappa\eta] = [\kappa] \oplus [\kappa'] \oplus [\kappa\alpha], \quad [\bar{\kappa}\kappa'] = [\eta],$$

$$[\hat{\eta}^2] = [\mathrm{id}_M] \oplus [\beta] \oplus 2[\hat{\eta}], \quad [\hat{\eta}\kappa] = [\kappa] \oplus [\kappa'] \oplus [\beta\kappa],$$

with $[\beta\kappa] = [\kappa\alpha]$, $d(\kappa) = \sqrt{2+\sqrt{3}}$, $d(\kappa') = \sqrt{2}$, $d(\alpha) = d(\beta) = 1$, $d(\eta) = d(\hat{\eta}) = 1 + \sqrt{3}$, $d(\xi) = 2 + \sqrt{3}$.

From $[\xi] = [\bar{\kappa}\kappa_Q\pi]$, we get $[\eta\xi] = [\eta\bar{\kappa}\kappa_Q\pi] = [\xi] \oplus [\bar{\kappa}'\kappa_Q\pi] \oplus [\alpha\xi]$. Since

$$\dim(\bar{\kappa}'\kappa_Q\pi, \bar{\kappa}'\kappa_Q\pi) = \dim(\kappa'\bar{\kappa}', \kappa_Q\bar{\kappa}_Q) = \dim(\mathrm{id}_M \oplus \beta, \mathrm{id}_M \oplus \hat{\eta}_Q) = 1,$$

the endomorphism $\bar{\kappa}'\kappa_Q\pi$ is irreducible, and Lemma 5.5 implies $\dim(\kappa\iota\bar{\iota}\bar{\kappa}, \kappa\iota\bar{\iota}\bar{\kappa}) = 8$.

Suppose that $\mathfrak{Q}$ is in Class III. Then thanks to Corollary 4.6,(3), we have

$$[\kappa\iota\bar{\iota}\bar{\kappa}] = [\mathrm{id}_M] \oplus [\theta] \oplus [\hat{\eta}] \oplus [\theta\hat{\eta}] \oplus [\hat{\eta}\theta] \oplus [\hat{\eta}\theta\hat{\eta}],$$

and $\hat{\eta}\theta\hat{\eta}$ does not contain any other irreducible components above. Therefore we must have $\dim(\hat{\eta}\theta\hat{\eta}, \hat{\eta}\theta\hat{\eta}) = 3$. However,

$$\begin{aligned}\dim(\hat{\eta}\theta\hat{\eta}, \hat{\eta}\theta\hat{\eta}) &= \dim(\theta\hat{\eta}^2\theta, \hat{\eta}^2) = \dim(\mathrm{id}_M \oplus \theta\beta\theta \oplus 2\theta\hat{\eta}\theta, \mathrm{id}_M \oplus \beta \oplus 2\hat{\eta}) \\ &= 1 + \dim(\theta\beta\theta, \beta),\end{aligned}$$

which is a contradiction. Therefore $\mathfrak{Q}$ is in Class II. By Corollary 4.6,(4), we get $[\bar{\pi}\eta_Q\pi] = [\bar{\kappa}'\kappa_Q\pi]$ and the principal graph of $N \subset P$ is as stated above. □

We exclude A_7, D_6, and E_8 for the principal graph of $P \subset M$.

Lemma 5.10. *There exists no quadrilateral* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *in Class II or Class III such that the principal graph of* $P \subset M$ *is* A_7.

Proof. Suppose there exists such a quadrilateral $\mathfrak{Q}$. Since $P \subset M$ is the A_7 subfactor, there exist outer automorphisms α of P and β of M and such that

$$[\hat{\eta}^2] = [\mathrm{id}_M] \oplus [\hat{\eta}] \oplus [\beta\hat{\eta}],$$

$$[\bar{\kappa}'\kappa] = [\eta] \oplus [\alpha\eta].$$

Lemma 5.5 shows $[\eta\xi] = [\xi] \oplus [\bar{\kappa}'\kappa_Q\pi]$.

Suppose that $\bar{\kappa}'\kappa_Q\pi$ is irreducible. Then Lemma 5.5 shows that $\dim(\kappa\iota\bar{\kappa}\bar{\iota}, \kappa\iota\bar{\kappa}\bar{\iota}) = 7$. Since $\kappa'\bar{\kappa}$ is reducible, Corollary 4.6,(3),(5) shows that the quadrilateral $\mathfrak{Q}$ should be in Class III and $\hat{\eta}\theta\hat{\eta}$ is decomposed into two mutually inequivalent irreducibles, which implies

$$2 = \dim(\hat{\eta}\theta\hat{\eta}, \hat{\eta}\theta\hat{\eta}) = \dim(\theta\hat{\eta}^2\theta, \hat{\eta}^2) = (\mathrm{id}_M \oplus \theta\hat{\eta}\theta \oplus \theta\beta\hat{\eta}\theta, \mathrm{id}_M \oplus \hat{\eta} \oplus \beta\hat{\eta}).$$

Thus either $[\hat{\eta}] = [\theta\beta\hat{\eta}\theta]$, $[\beta\hat{\eta}] = [\theta\hat{\eta}\theta]$, or $[\beta\hat{\eta}] = [\theta\beta\hat{\eta}\theta]$ should hold. However, if $[\hat{\eta}] = [\theta\beta\hat{\eta}\theta]$, we would get $[\hat{\eta}^2] = [\bar{\hat{\eta}}\hat{\eta}] = [\theta\hat{\eta}^2\theta]$, which is a contradiction. The same reasoning shows that the other two do not occur as well and we conclude that $\bar{\kappa}'\kappa_Q\pi$ is reducible.

Suppose that $\bar{\kappa}'\kappa_Q\pi$ is reducible. Since any irreducible component ζ of $\bar{\kappa}'\kappa_Q\pi$ is contained in $\iota'\bar{\iota}$, Frobenius reciprocity shows that $\zeta\iota$ contains ι'. On the other hand,

$$[\bar{\kappa}'\kappa_Q\pi\iota] = [\bar{\kappa}'\kappa_Q\iota_Q] = [\bar{\kappa}'\kappa\iota] = [\eta\iota] \oplus [\alpha\eta\iota] = [\iota'] \oplus [\alpha\iota'].$$

This implies $[\iota'] = [\alpha\iota']$ and that $\bar{\kappa}'\kappa_Q\pi$ is decomposed into two irreducibles, say, ζ_1 and ζ_2 such that $[\zeta_1\iota] = [\zeta_2\iota] = [\iota']$. This shows $[\iota'\bar{\iota}] = [\xi] \oplus [\eta] \oplus [\zeta_1] \oplus [\zeta_2]$ and the set of $P-P$ sectors appearing the dual principal graph of $N \subset P$ is $\{[\mathrm{id}_P], [\xi], [\eta], [\zeta_1], [\zeta_2]\}$. However, since $[\alpha\iota'] = [\iota']$, the automorphism α appears in $\iota'\bar{\iota}'$, which is a contradiction. Therefore the statement is shown. □

Lemma 5.11. *There exists no quadrilateral* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *in Class II or Class III such that the principal graph of* $P \subset M$ *is* D_6.

Proof. Suppose that $\mathfrak{Q}$ is such a quadrilateral. We first recall the fusion rule for the D_6 subfactor:

$$[\kappa'\bar{\kappa}] = [\hat{\eta}] \oplus [\hat{\eta}_1] \oplus [\hat{\eta}_2], \quad [\kappa'\bar{\kappa}'] = [\mathrm{id}_M] \oplus 2[\hat{\eta}] \oplus [\hat{\eta}_1] \oplus [\hat{\eta}_2],$$
$$[\hat{\eta}^2] = [\mathrm{id}_M] \oplus [\hat{\eta}] \oplus [\hat{\eta}_1] \oplus [\hat{\eta}_2],$$
$$[\hat{\eta}_1^2] = [\mathrm{id}_M] \oplus [\hat{\eta}_1], \quad [\hat{\eta}_2]^2 = [\mathrm{id}_M] \oplus [\hat{\eta}_2], \quad [\hat{\eta}_1\hat{\eta}_2] = [\hat{\eta}_2\hat{\eta}_1] = [\hat{\eta}],$$
$$[\hat{\eta}_1\hat{\eta}] = [\hat{\eta}\hat{\eta}_1] = [\hat{\eta}] \oplus [\hat{\eta}_2], \quad [\hat{\eta}_2\hat{\eta}] = [\hat{\eta}\hat{\eta}_2] = [\hat{\eta}] \oplus [\hat{\eta}_1],$$

$d(\hat{\eta}) = (3+\sqrt{5})/2$, $d(\hat{\eta}_1) = d(\hat{\eta}_2) = (1+\sqrt{5})/2$.

Lemma 5.5 shows $[\eta\xi] = [\xi] \oplus [\bar{\kappa}'\kappa_Q\pi]$. We claim that $\bar{\kappa}'\kappa_Q\pi$ is irreducible. Indeed,

$$\dim(\bar{\kappa}'\kappa_Q\pi, \bar{\kappa}'\kappa_Q\pi) = \dim(\kappa'\bar{\kappa}', \kappa_Q\bar{\kappa}_Q) = \dim(\mathrm{id}_M \oplus 2\hat{\eta} \oplus \hat{\eta}_1 \oplus \hat{\eta}_2, \mathrm{id}_M \oplus \hat{\eta}_Q).$$

Since $\mathfrak{Q}$ is cocommuting, we have $[\hat{\eta}] \neq [\hat{\eta}_Q]$ and we get the claim. Lemma 5.5 and the claim implies that $\dim(\kappa\iota\bar{\iota}\bar{\kappa}, \kappa\iota\bar{\iota}\bar{\kappa}) = 7$. Corollary 4.6,(3),(5) shows that $\mathfrak{Q}$ is in Class III and $\dim(\hat{\eta}\theta\hat{\eta}, \hat{\eta}\theta\hat{\eta}) = 2$, and so

$$\begin{aligned} 2 &= \dim(\theta\hat{\eta}^2\theta, \hat{\eta}^2) = \dim(\mathrm{id}_M \oplus \theta\hat{\eta}\theta \oplus \theta\hat{\eta}_1\theta \oplus \theta\hat{\eta}_2\theta, \mathrm{id}_M \oplus \hat{\eta} \oplus \hat{\eta}_1 \oplus \hat{\eta}_2) \\ &= 1 + \dim(\theta\hat{\eta}_1\theta \oplus \theta\hat{\eta}_2\theta, \hat{\eta}_1 \oplus \hat{\eta}_2). \end{aligned}$$

If $[\theta\hat{\eta}_1\theta] = [\hat{\eta}_2]$, we would have $[\theta\hat{\eta}_2\theta] = [\hat{\eta}_1]$ too as the period of θ is two, which is a contradiction. Thus we may assume $[\theta\hat{\eta}_1\theta] = [\hat{\eta}_1]$ and $[\theta\hat{\eta}_2\theta] \neq [\hat{\eta}_1], [\hat{\eta}_2]$.

Since $\hat{\eta}\theta\hat{\eta}$ contains $\theta\hat{\eta}\theta$ with multiplicity one, there exists an irreducible $\zeta \in \mathrm{End}_0(M)$ inequivalent to $\theta\hat{\eta}\theta$ such that $[\hat{\eta}\theta\hat{\eta}] = [\theta\hat{\eta}\theta] \oplus [\zeta]$. We claim that $[\zeta] = [\hat{\eta}\theta\hat{\eta}_2]$ holds. Indeed, we have

$$\dim(\hat{\eta}\theta\hat{\eta}_2, \hat{\eta}\theta\hat{\eta}_2) = \dim(\hat{\eta}^2, \theta\hat{\eta}_2^2\theta) = \dim(\mathrm{id}_M \oplus \hat{\eta} \oplus \hat{\eta}_1 \oplus \hat{\eta}_2, \mathrm{id}_M \oplus \theta\hat{\eta}_2\theta) = 1,$$

which shows that $\hat{\eta}\theta\hat{\eta}_2$ is irreducible. Since $[\theta\hat{\eta}\theta] \neq [\hat{\eta}\theta\hat{\eta}_2]$ and

$$\dim(\hat{\eta}\theta\hat{\eta}, \hat{\eta}\theta\hat{\eta}_2) = \dim(\hat{\eta}^2, \theta\hat{\eta}_2\hat{\eta}\theta) = \dim(\mathrm{id}_M \oplus \hat{\eta} \oplus \hat{\eta}_1 \oplus \hat{\eta}_2, \theta\hat{\eta}_1\theta \oplus \theta\hat{\eta}\theta) = 1,$$

we get the claim. Since

$$[\kappa\iota\bar{\iota}\bar{\kappa}] = [\mathrm{id}_M] \oplus [\theta] \oplus [\hat{\eta}] \oplus [\theta\hat{\eta}] \oplus [\hat{\eta}\theta] \oplus [\theta\hat{\eta}\theta] \oplus [\zeta],$$

and $[\theta\kappa\iota] = [\kappa\iota]$, we get $[\theta\zeta] = [\zeta]$. However,

$$\dim(\hat{\eta}\theta\hat{\eta}_2, \theta\hat{\eta}\theta\hat{\eta}_2) = \dim(\hat{\eta}\theta\hat{\eta}, \theta\hat{\eta}_2^2\theta) = \dim(\theta\hat{\eta}\theta \oplus \zeta, \mathrm{id}_M \oplus \theta\hat{\eta}_2\theta) = 0,$$

which is a contradiction. Thus the statement is proven. □

Lemma 5.12. *There exists no quadrilateral* $\mathfrak{Q} = \begin{matrix} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{matrix}$ *in Class II or Class III such that the principal graph of* $P \subset M$ *is* E_8.

Proof. Suppose that $\mathfrak{Q}$ is such a quadrilateral. Since the E_8 factor is 4-supertransitive, there exists an irreducible $\hat{\eta}' \in \mathrm{End}_0(M)$ such that $[\hat{\eta}^2] = [\mathrm{id}_M] \oplus [\hat{\eta}] \oplus [\hat{\eta}']$. Note that we have $d(\hat{\eta}) \neq d(\hat{\eta}')$ and $\hat{\eta}$ is the only irreducible sector σ contained in $\hat{\eta}'^2$ (also in $\kappa'\bar{\kappa}'$) such that $d(\hat{\eta}) = d(\sigma)$ (see [17, Section 3.3]).

As before, we have $[\xi\eta] = [\xi] \oplus [\bar{\kappa}'\kappa_Q\pi]$ and

$$\dim(\bar{\kappa}'\kappa_Q\pi, \bar{\kappa}'\kappa_Q\pi) = \dim(\kappa'\bar{\kappa}', \kappa_Q\bar{\kappa}_Q) = 1 + \dim(\kappa'\bar{\kappa}', \hat{\eta}_Q) = 1.$$

Thus $\mathfrak{Q}$ is in Class III and

$$2=\dim(\hat{\eta}\theta\hat{\eta},\hat{\eta}\theta\hat{\eta})=\dim(\theta\hat{\eta}^2\theta,\hat{\eta}^2)=\dim(\mathrm{id}_M\oplus\theta\hat{\eta}\theta\oplus\theta\hat{\eta}'\theta,\mathrm{id}_M\oplus\hat{\eta}\oplus\hat{\eta}'),$$

which shows $[\theta\hat{\eta}'\theta]=[\hat{\eta}']$. However, this implies $[\theta\hat{\eta}'^2\theta]=[\hat{\eta}'^2]$, which contradicts $[\theta\hat{\eta}\theta]\neq[\hat{\eta}]$ as $\hat{\eta}'^2$ contains $\hat{\eta}$. □

We are ready to prove Theorem 5.3 now.

Proof of Theorem 5.3. Let $\mathfrak{Q}$ be in Class II and (5,3)-supertransitive such that the principal graph of $P\subset M$ is not A_4. Corollary 4.6,(5) shows $\dim(\kappa\iota\bar{\iota}\bar{\kappa},\kappa\iota\bar{\iota}\bar{\kappa})\geq 8$ and Lemma 5.5 shows $\dim(\bar{\kappa}'\kappa_Q\pi,\bar{\kappa}'\kappa_Q\pi)\geq 2$. Since the principal graph of $P\subset M$ is not A_5, the depth of $P\subset M$ is at least 5. Thus we use the following parameterization of $M-M$, $M-P$, and $P-P$ sectors arising from $P\subset M$.



Corollary 4.6,(5) shows

$$[\kappa\iota\bar{\iota}\bar{\kappa}]=[\mathrm{id}_M]\oplus[\hat{\eta}]\oplus[\hat{\eta}_Q]\oplus[\hat{\eta}']\oplus 2[\kappa_Q\pi\bar{\kappa}].$$

Since $\mathfrak{Q}$ is cocommuting, $[\hat{\eta}]\neq[\hat{\eta}_Q]$. Since the principal graph of $P\subset M$ is neither A_5 nor A_7, we have $d(\hat{\eta}_Q)\neq d(\hat{\eta}')$, $d(\kappa_Q\pi)\neq d(\kappa')$ and $d(\kappa_Q\pi)\neq d(\kappa'')$. Since

$$\dim(\kappa_Q\pi\bar{\kappa},\hat{\eta}')=\dim(\kappa_Q\pi,\hat{\eta}'\kappa)=\dim(\kappa_Q\pi,\kappa'\oplus\kappa'')=0,$$

we get $[\kappa_Q\pi\bar{\kappa}]\neq[\hat{\eta}']$. Thus $\dim(\kappa\iota\bar{\iota}\bar{\kappa},\kappa\iota\bar{\iota}\bar{\kappa})=8$ and $\dim(\bar{\kappa}'\kappa_Q\pi,\bar{\kappa}'\kappa_Q\pi)=2$. Thanks to Corollary 4.6,(4),(5), the endomorphism $\bar{\kappa}'\kappa_Q\pi$ is decomposed into $\bar{\pi}\eta_Q\pi$ and an irreducible endomorphism of P, say ξ'. Since

$$[\bar{\kappa}'\kappa_Q\pi\iota]=[\bar{\kappa}'\kappa\iota]=[\eta\iota]\oplus[\eta'\iota]=[\iota']\oplus[\eta'\iota],$$

and $[\bar{\pi}\eta_Q\pi\iota]=[\iota']$, we get $[\xi'\iota]=[\eta'\iota]$, which contains ι' as ξ' is contained in $\iota'\bar{\iota}$. Thus Frobenius reciprocity implies

$$1=\dim(\iota',\eta'\iota)=\dim(\iota'\bar{\iota},\eta')=\dim(\xi\oplus\eta\oplus\bar{\pi}\eta_Q\pi\oplus\xi',\eta').$$

If $[\xi] = [\eta']$, we would have $[\xi'\iota] = [\iota] \oplus [\iota']$ and by Frobenius reciprocity $[\xi'] = [\xi]$, a contradiction. Thus $[\xi'] = [\eta']$.



We claim $[\eta\bar{\pi}\eta_Q\pi] = [\xi] \oplus [\eta']$. Thanks to Corollary 4.6(5), we know that $\kappa_Q\pi\bar{\kappa}$ is self-conjugate. Thus

$$\begin{aligned}[\xi^2] &= [\bar{\kappa}\kappa_Q\pi\bar{\kappa}\kappa_Q\pi] = [\bar{\kappa}\kappa\bar{\pi}\bar{\kappa}_Q\kappa_Q\pi] \\ &= [(\mathrm{id}_P \oplus \eta)(\mathrm{id}_P \oplus \bar{\pi}\eta_Q\pi)] = [\mathrm{id}_P] \oplus [\eta] \oplus [\bar{\pi}\eta_Q\pi] \oplus [\eta\bar{\pi}\eta_Q\pi],\end{aligned}$$

which shows the claim.

Let $[\eta'\iota] = [\iota'] \oplus \bigoplus_{i=1}^k m_i[\iota_i]$, and $[\iota_i\bar{\iota}] = m_i[\eta'] \oplus \bigoplus_{j=1}^l m_{ij}[\xi_j]$ be the irreducible decompositions. We compute $[\eta\xi]$ in two ways:

$$\begin{aligned}[\eta\xi][\xi] &= [\xi^2] \oplus [\bar{\pi}\eta_Q\pi\xi] \oplus [\eta'\xi] \\ &= [\mathrm{id}_P] \oplus [\xi] \oplus [\eta] \oplus [\bar{\pi}\eta_Q\pi] \oplus [\eta'] \oplus [\xi] \oplus [\eta] \oplus [\eta'] \\ \oplus & \quad [\xi] \oplus [\eta] \oplus [\bar{\pi}\eta_Q\pi] \oplus \sum_{i=1}^k m_i^2[\eta'] \oplus \bigoplus_{j=1}^l \sum_{i=1}^k m_i m_{ij}[\xi_j] \\ &= [\mathrm{id}_P] \oplus 3[\xi] \oplus 3[\eta] \oplus 2[\bar{\pi}\eta_Q\pi] \oplus (2 + \sum_{i=1}^k m_i^2)[\eta'] \oplus \bigoplus_{j=1}^l \sum_{i=1}^k m_i m_{ij}[\xi_j],\end{aligned}$$

$$\begin{aligned}[\eta][\xi^2] &= [\eta] \oplus [\eta\xi] \oplus [\eta^2] \oplus [\eta\bar{\pi}\eta_Q\pi] \oplus [\eta\eta'] \\ &= [\eta] \oplus [\xi] \oplus [\bar{\pi}\eta_Q\pi] \oplus [\eta'] \oplus [\mathrm{id}_P] \oplus [\eta] \oplus [\eta'] \oplus [\xi] \oplus [\eta'] \oplus [\eta\eta'] \\ &= [\mathrm{id}_P] \oplus 2[\xi] \oplus 2[\eta] \oplus [\bar{\pi}\eta_Q\pi] \oplus 3[\eta'] \oplus [\eta\eta'].\end{aligned}$$

This shows

$$[\eta\eta'] = [\xi] \oplus [\eta] \oplus [\bar{\pi}\eta_Q\pi] \oplus (\sum_{i=1}^k m_i^2 - 1)[\eta'] \oplus \bigoplus_{j=1}^l \sum_{i=1}^k m_i m_{ij}[\xi_j].$$

Since $P \subset M$ is 5-supertransitive, the multiplicity of η' in $\eta\eta'$ is 1, which implies $k = 2$ and $m_1 = m_2 = 1$. This shows

$$[\bar{\kappa}\kappa''] = [\eta'] \oplus [\bar{\pi}\eta_Q\pi] \oplus [\xi] \oplus \bigoplus_{j=1}^l (m_{1j} + m_{2j})[\xi_j]$$

and $P \subset M$ cannot be 6-supertransitive. Frobenius reciprocity implies that $\kappa\bar{\pi}\eta_Q\pi$ and $\kappa\xi$ contain κ'' with multiplicity one and

$$[\kappa\xi] = [\kappa\bar{\kappa}\kappa_Q\pi] = [\kappa_Q\pi] \oplus [\hat{\eta}\kappa_Q\pi],$$

On the other hand, the endomorphisms $\kappa\bar{\pi}\eta_Q\pi$ and $\hat{\eta}\kappa_Q\pi$ are irreducible as we have

$$\begin{aligned}\dim(\kappa\bar{\pi}\eta_Q\pi, \kappa\bar{\pi}\eta_Q\pi) &= \dim(\bar{\kappa}\kappa\bar{\pi}\eta_Q\pi, \bar{\pi}\eta_Q\pi) = \dim(\bar{\pi}\eta_Q\pi \oplus \eta\bar{\pi}\eta_Q\pi, \bar{\pi}\eta_Q\pi)\\ &= \dim(\bar{\pi}\eta_Q\pi \oplus \xi \oplus \eta', \bar{\pi}\eta_Q\pi) = 1,\end{aligned}$$

$$\dim(\hat{\eta}\kappa_Q\pi, \hat{\eta}\kappa_Q\pi) = \dim(\hat{\eta}^2, \kappa_Q\bar{\kappa}_Q) = \dim(\mathrm{id}_M \oplus \hat{\eta} \oplus \hat{\eta}', \mathrm{id}_M \oplus \hat{\eta}_Q) = 1,$$

and so $[\kappa''] = [\kappa\bar{\pi}\hat{\eta}_Q\pi] = [\hat{\eta}\kappa_Q\pi]$. By induction, we can get $d(\eta) = d(\kappa)^2 - 1$, $d(\kappa') = d(\kappa)(d(\kappa)^2 - 2)$, $d(\eta') = d(\kappa)^4 - 3d(\kappa^2) + 1$, $d(\kappa'') = d(\kappa)(d(\kappa)^4 - 4d(\kappa)^2 + 3)$. Therefore $d(\kappa'') = d(\kappa\bar{\pi}\hat{\eta}_Q\pi)$ implies $d(\kappa)^4 - 5d(\kappa)^2 + 4 = 0$ and $[M : P] = d(\kappa^2) = 4$. This shows that the principal graph of $P \subset M$ is $E_8^{(1)}$.



The above computation shows that ξ is determined solely by the inclusion $P \subset M$. Since $N \subset P$ is 3-supertransitive, a Q-system for $\mathrm{id}_P \oplus \xi$ is unique up to equivalence and so M is uniquely determined by $P \subset M$ up to inner conjugacy in P. Therefore uniqueness of such a quadrilateral of the hyperfinite II_1 factors up to conjugacy follows from Theorem 3.14.

We show existence now. Let R be a factor and α be an outer action of the alternating group $\mathfrak{A}_5$. We regard $\mathfrak{A}_4$ as a subgroup of $\mathfrak{A}_5$ and fix it. We choose two mutually inequivalent 2-dimensional irreducible projective unitary representations σ_1 and σ_2. Since σ_1 and σ_2 carry a unique non-trivial element in $H^2(\mathfrak{A}_5, \mathbb{T})$, we may assume that they have the same cocycle ω. Since $\mathfrak{A}_4$ has a unique 2-dimensional irreducible projective unitary representation up to equivalence, we may assume that the restriction of σ_1 and that of σ_2 to $\mathfrak{A}_4$ coincide. Let $M_2(\mathbb{C})$ be the 2 by 2 matrix algebra and let $\beta_g = \alpha_g \otimes \mathrm{Ad}\sigma_1(g)$ for $g \in \mathfrak{A}_5$, which is an action of $\mathfrak{A}_5$ on $R \otimes M_2(\mathbb{C})$. We set

$$N = (R \otimes \mathbb{C}) \rtimes_\beta \mathfrak{A}_4 \subset P = (R \otimes \mathbb{C}) \rtimes_\beta \mathfrak{A}_5 \subset M = (R \otimes M_2(\mathbb{C})) \rtimes_\beta \mathfrak{A}_5.$$

Then the principal graph of $P \subset M$ is $E_8^{(1)}$ [11, p.224], the inclusion $N \subset P$ is 3-supertransitive [39], and it is routine work to show that $N \subset M$ is irreducible. Let $u_g = 1 \otimes \sigma_2(g)\sigma_1(g)^*$, which is a β-cocycle, and let $\{\lambda_g\}_{g \in \mathfrak{A}_5}$ be the implementing unitary in $M = (R \otimes M_2(\mathbb{C})) \rtimes_\beta \mathfrak{A}_5$. Since u_g commutes with $R \otimes \mathbb{C}$, we can introduce a homomorphism $\pi : P \to M$ by setting $\pi(x) = x$ for $x \in R \otimes \mathbb{C}$ and $\pi(\lambda_g) = u_g\lambda_g$ for $g \in \mathfrak{A}_5$. We set $Q = \pi(P)$. The restriction of π to N is the identity map because $u_g = 1$ for $g \in \mathfrak{A}_4$. Since σ_1 and σ_2 are not equivalent, there exists $g \in \mathfrak{A}_5 \setminus \mathfrak{A}_4$ such that u_g is not a scalar, and so $P \neq Q$. Since $\mathfrak{A}_4$ is maximal in $\mathfrak{A}_5$ there is no intermediate subfactor between $N \subset P$. Since the principal graph of $P \subset M$ is $E_8^{(1)}$, there is no intermediate subfactor between $P \subset M$. Therefore we conclude that $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ is a quadrilateral, which is noncommuting thanks to Theorem 3.10,(1). Theorem 4.3 shows that $\mathfrak{Q}$ is cocommuting as we have $[M : P] = [P : N]-1$.

To finish the proof, we show that the principal graph of $Q \subset M$ is also $E_8^{(1)}$. Since $(N \subset Q) \cong (R \rtimes_\alpha \mathfrak{A}_4 \subset R \rtimes_\alpha \mathfrak{A}_5)$, the category of $Q-Q$ bimodules arising from $N \subset Q$ is equivalent to the unitary dual of $\mathfrak{A}_5$. Lemma 4.1 shows that this category contains the category of $Q-Q$ bimodules arising from $Q \subset M$. Since $[M : Q] = 4$, we conclude that the principal graph of $Q \subset M$ is $E_8^{(1)}$. □

To prove Theorem 5.4, we have to exclude two more cases.

Proposition 5.13. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *be a quadrilateral in Class II or III such that the principal graph of* $P \subset M$ *is* $E_7^{(1)}$. *Then* $\mathfrak{Q}$ *is in Class III. There exists a factor* R *and an outer action of* $G = \mathfrak{S}_5$ *on* R *with two maximal subgroups* $H \neq K$ *of* G *such that* $M = R^{H\cap K}$, $P = R^H$, $Q = R^K$, *and* $N = R^G$. *Such a quadrilateral of the hyperfinite* II_1 *factors exists and is unique up to conjugacy.*

Proof. We use the following parameterization of $M-M$, $M-P$, and $P-P$ sectors arising from $P \subset M$:



The category of $M-M$ bimodules and $P-P$ bimodules arising from $P \subset M$ are isomorphic to the unitary dual of $\mathfrak{S}_4$, which gives the fusion rule of the above sectors. We have

$$[\eta^2] = [\mathrm{id}_P] \oplus [\eta] \oplus [\eta'] \oplus [\alpha\eta], \quad [\eta'^2] = [\mathrm{id}_P] \oplus [\alpha] \oplus [\eta'],$$

$$[\alpha\eta] = [\eta\alpha], \quad [\alpha\eta'] = [\eta'\alpha] = [\eta'], \quad [\alpha^2] = [\mathrm{id}_P],$$

$$[\hat{\eta}^2] = [\mathrm{id}_M] \oplus [\hat{\eta}] \oplus [\hat{\eta}'] \oplus [\beta\hat{\eta}], \quad [\hat{\eta}'^2] = [\mathrm{id}_M] \oplus [\beta] \oplus [\hat{\eta}'],$$

$$[\beta\hat{\eta}] = [\hat{\eta}\beta], \quad [\beta\hat{\eta}'] = [\hat{\eta}\beta'] = [\hat{\eta}'], \quad [\beta^2] = [\mathrm{id}_M],$$

$d(\eta) = d(\hat{\eta}) = 3$, $d(\eta') = d(\hat{\eta}') = 2$, $d(\alpha) = d(\beta) = 1$.

First we claim that $\bar{\kappa}'\kappa_Q\pi$ is irreducible, which shows that $\mathfrak{Q}$ is in Class III thanks to Corollary 4.6,(4) and Lemma 5.5. Indeed, we have

$$\begin{aligned}\dim(\bar{\kappa}'\kappa_Q\pi,\bar{\kappa}'\kappa_Q\pi) &= \dim(\kappa'\bar{\kappa}',\kappa_Q\bar{\kappa}_Q)\\ &= \dim(\mathrm{id}_M\oplus\beta\oplus 2\hat{\eta}\oplus 2\beta\hat{\eta}\oplus\hat{\eta}',\mathrm{id}_M\oplus\hat{\eta}_Q)\\ &= 1+2\dim(\beta\hat{\eta},\hat{\eta}_Q),\end{aligned}$$

which shows that either $[\beta\hat{\eta}]\neq[\hat{\eta}_Q]$ and $\bar{\kappa}'\kappa_Q\pi$ is irreducible or $[\beta\hat{\eta}]=[\hat{\eta}_Q]$ and $\bar{\kappa}'\kappa_Q\pi$ is a direct sum of three irreducible distinct sectors. Suppose that the latter holds and $[\bar{\kappa}'\kappa_Q\pi]=[\xi_1]\oplus[\xi_2]\oplus[\xi_3]$. Then $\xi_i\iota$ would contain ι' for each $i=1,2,3$. On the other hand, we have

$$[\bar{\kappa}'\kappa_Q\pi\iota]=[\bar{\kappa}'\kappa\iota]=[\eta\iota]\oplus[\eta'\iota]\oplus[\alpha\eta\iota]=[\iota']\oplus[\eta'\iota]\oplus[\alpha\iota'].$$

Since $d(\eta')=2$, $d(\iota)=\sqrt{5}$, and $d(\iota')=3\sqrt{5}$, this is impossible and we get the claim.

Lemma 5.5 and Corollary 4.6,(3),(5) imply

$$\begin{aligned}2 &= \dim(\hat{\eta}\theta\hat{\eta},\hat{\eta}\theta\hat{\eta})=\dim(\hat{\eta}^2,\theta\hat{\eta}^2\theta)\\ &= \dim(\mathrm{id}_M\oplus\hat{\eta}\oplus\hat{\eta}'\oplus\beta\hat{\eta},\mathrm{id}_M\oplus\theta\hat{\eta}\theta\oplus\theta\hat{\eta}'\theta\oplus\theta\beta\hat{\eta}\theta)\\ &= 1+\dim(\hat{\eta},\theta\beta\hat{\eta}\theta)+\dim(\beta\hat{\eta},\theta\hat{\eta}\theta)+\dim(\beta\hat{\eta},\theta\beta\hat{\eta}\theta)+\dim(\hat{\eta}',\theta\hat{\eta}'\theta).\end{aligned}$$

If $[\hat{\eta}]=[\theta\beta\hat{\eta}\theta]$, we would have $[\hat{\eta}^2]=[\overline{\hat{\eta}}\hat{\eta}]=[\theta\hat{\eta}^2\theta]$, which is impossible. For the same reason, we get $[\beta\hat{\eta}]\neq[\theta\hat{\eta}\theta]$, $[\beta\hat{\eta}]\neq[\theta\beta\hat{\eta}\theta]$ and so $[\hat{\eta}']=[\theta\hat{\eta}'\theta]$.

In [21, Corollary 4.8], the second-named author showed that there exists a factor R and an outer action μ of $\mathfrak{S}_4$ on R such that $M=R^{\mathfrak{S}_3}$ and $P=R^{\mathfrak{S}_4}$. Let $\nu=\iota_{R,M}$. Then we have $[\bar{\nu}\nu]=[\mathrm{id}_M]\oplus 2[\sigma]\oplus[\varphi]$ where $d(\sigma)=2$, $d(\varphi)=1$ and

$$[\sigma^2]=[\mathrm{id}_M]\oplus[\varphi]\oplus[\sigma],\quad [\varphi\sigma]=[\sigma\varphi]=[\sigma],\quad [\varphi^2]=[\mathrm{id}_M].$$

We claim that $[\hat{\eta}']=[\sigma]$ and $[\beta]=[\varphi]$ hold. Indeed, since $d(\hat{\eta})=3$ and $\hat{\eta}^2$ does not contain any non-trivial automorphism, $[\mathrm{id}_M],[\varphi],[\sigma],[\hat{\eta}]$, and $[\varphi\hat{\eta}]$ are all distinct sectors. Since $P\subset R$ is of depth 2 and $[R:P]=24$, we get

$$\begin{aligned}24 &= \dim(\nu\kappa\bar{\kappa}\bar{\nu},\nu\kappa\bar{\kappa}\bar{\nu})=\dim(\bar{\nu}\nu\kappa\bar{\kappa},\kappa\bar{\kappa}\bar{\nu}\nu)\\ &= \dim((\mathrm{id}_M\oplus\varphi\oplus 2\sigma)(\mathrm{id}_M\oplus\hat{\eta}),(\mathrm{id}_M\oplus\hat{\eta})(\mathrm{id}_M\oplus\varphi\oplus 2\sigma))\\ &= \dim(\mathrm{id}_M\oplus\varphi\oplus 2\sigma\oplus\hat{\eta}\oplus\varphi\hat{\eta},\mathrm{id}_M\oplus\varphi\oplus 2\sigma\oplus\hat{\eta}\oplus\hat{\eta}\varphi)\\ &+ 2\dim(\mathrm{id}_M\oplus\varphi\oplus 2\sigma\oplus\hat{\eta}\oplus\varphi\hat{\eta},\hat{\eta}\sigma)+2\dim(\sigma\hat{\eta},\mathrm{id}_M\oplus\varphi\oplus 2\sigma\oplus\hat{\eta}\oplus\hat{\eta}\varphi)\\ &+ 4\dim(\sigma\hat{\eta},\hat{\eta}\sigma)\\ &= 7+\dim(\varphi\hat{\eta},\hat{\eta}\varphi)+2\dim(\sigma\oplus\varphi\sigma\oplus 2\sigma^2,\hat{\eta})+2\dim(\hat{\eta}^2,\sigma)+2\dim(\varphi\hat{\eta},\hat{\eta}\sigma)\\ &+ 2\dim(\hat{\eta},\sigma\oplus\sigma\varphi\oplus 2\sigma^2)+2\dim(\sigma,\hat{\eta}^2)+2\dim(\sigma\hat{\eta},\hat{\eta}\varphi)+4\dim(\sigma\hat{\eta},\hat{\eta}\sigma),\end{aligned}$$

which shows $[\varphi\hat{\eta}]=[\hat{\eta}\varphi]$ and

$$\begin{aligned}4 &= 2\dim(\sigma^2,\hat{\eta})+\dim(\hat{\eta}^2,\sigma)+\dim(\hat{\eta}^2\varphi,\sigma)+\dim(\sigma\hat{\eta},\hat{\eta}\sigma)\\ &= 2\dim(\hat{\eta}^2,\sigma)+\dim(\sigma\hat{\eta},\hat{\eta}\sigma)=2\dim(\mathrm{id}_M\oplus\hat{\eta}\oplus\hat{\eta}'\oplus\beta\hat{\eta},\sigma)+\dim(\sigma\hat{\eta},\hat{\eta}\sigma)\\ &= 2\dim(\hat{\eta}',\sigma)+\dim(\sigma\hat{\eta},\hat{\eta}\sigma).\end{aligned}$$

If $[\hat{\eta}']\neq[\sigma]$, we would have,

$$\dim(\sigma\hat{\eta},\sigma\hat{\eta})=\dim(\sigma^2,\hat{\eta}^2)=\dim(\mathrm{id}\oplus\varphi\oplus\sigma,\mathrm{id}\oplus\hat{\eta}\oplus\hat{\eta}'\oplus\beta\hat{\eta})=1,$$

which means that $\sigma\hat{\eta}$ is irreducible. A similar argument shows that $\hat{\eta}\sigma$ is irreducible too and $\dim(\sigma\hat{\eta},\hat{\eta}\sigma)$ is at most 1, which is a contradiction. Thus we get $[\sigma]=[\hat{\eta}']$ and in consequence, the equality $[\varphi]=[\beta]$ holds.

Since $\mathfrak{Q}$ is in Class III, the principal graph of $Q\subset M$ is $E_7^{(1)}$ too and we can apply the same argument to $Q\subset M$. Then there exists a factor $\tilde{R}$ and an outer action of $\mathfrak{S}_4$ on $\tilde{R}$ such that $M=\tilde{R}^{\mathfrak{S}_3}$ and $Q=\tilde{R}^{\mathfrak{S}_4}$. Note that we have $[\theta\kappa]=[\kappa_Q\pi]$, which implies $[\hat{\eta}_Q]=[\theta\hat{\eta}\theta]$ and

$$[\hat{\eta}_Q^2]=[\mathrm{id}_M]\oplus[\theta\hat{\eta}\theta]\oplus[\theta\hat{\eta}'\theta]\oplus[\theta\beta\hat{\eta}\theta]=[\mathrm{id}_M]\oplus[\hat{\eta}_Q]\oplus[\hat{\eta}']\oplus[\theta\beta\hat{\eta}\theta].$$

Thus the above argument shows $[\overline{\iota_{\tilde{R},M}}\iota_{\tilde{R},M}]=[\overline{\iota_{R,M}}\iota_{R,M}]$. The second-named author and Kosaki [26, Theorem 3.3] showed that the cohomology $H^2(M\subset R)$ is identified with $H^2(\hat{\mathfrak{S}}_3,\mathbb{T})$ introduced by Wassermann [55]. Since the order of $\mathfrak{S}_3$ is square free, there is only one ergodic action of $\mathfrak{S}_3$ on a 6-dimensional C^*-algebra and the latter is trivial. Therefor we may assume $R=\tilde{R}$. We denote by μ^Q the action of $\mathfrak{S}_4$ on R whose fixed point algebra is Q. We set $H=\mu_{\mathfrak{S}_4}$ and $K=\mu^Q_{\mathfrak{S}_4}$.

We claim that $N\subset R$ is irreducible. Indeed, we known that $\hat{\eta}\theta\hat{\eta}$ is a direct sum of $\theta\hat{\eta}\theta$ and an irreducible, say ζ, with $d(\zeta)=6$. Thus we have

$$\begin{aligned}\dim(\nu\kappa\iota,\nu\kappa\iota) &= \dim(\bar{\nu}\nu,\kappa\iota\bar{\iota}\bar{\kappa})\\ &= \dim(\mathrm{id}_M\oplus\beta\oplus 2\hat{\eta}',\mathrm{id}_M\oplus\theta\oplus\hat{\eta}\oplus\theta\hat{\eta}\oplus\hat{\eta}\theta\oplus\theta\hat{\eta}\theta\oplus\zeta)\\ &= 1+\dim(\beta,\theta).\end{aligned}$$

Since $[\beta\hat{\eta}]=[\hat{\eta}\beta]$, if $[\beta]=[\theta]$, the endomorphism $\hat{\eta}\theta\hat{\eta}$ would contain an automorphism, which contradicts Corollary 4.6,(3). Therefore the claim is shown.

Let G be the group generated by H and K in $\mathrm{Aut}(R)$. Then the fixed point algebra for R^G is $P\cap Q=N$ and the action of G on R is outer for $N\subset R$ is irreducible. Since $N\subset P$ is 3-supertransitive, the action of G on G/H is 3-transitive. Therefore from $[G:H]=[P:N]=[M:P]+1=5$, we conclude that G is isomorphic to $\mathfrak{S}_5$ because any 3-transitive transforation group of the five point set is either $\mathfrak{A}_5$ or $\mathfrak{S}_5$. Uniqueness in the case of the hyperfinite II_1 factors follows from Theorem 4.8. ☐

Proposition 5.14. *Let* $\mathfrak{Q}=\begin{array}{ccc}P&\subset&M\\ \cup&&\cup\\ N&\subset&Q\end{array}$ *be a quadrilateral in Class II or III such that the principal graph of* $P\subset M$ *is* $E_6^{(1)}$. *Then* $\mathfrak{Q}$ *is in Class III. There exists a factor* R *and an outer action of* $G=\mathfrak{A}_5$ *on* R *with two maximal subgroups* $H\neq K$ *of* G *such that* $M=R^{H\cap K}$, $P=R^H$, $Q=R^K$, *and* $N=R^G$. *Such a quadrilateral of the hyperfinite* II_1 *factors exists and is unique up to conjugacy.*

Proof. We employ the same strategy as in the proof of the previous proposition using $\mathfrak{A}_4$ instead of $\mathfrak{S}_4$. Since the principal graph of $P\subset M$ is $E_6^{(1)}$, there exists outer automorphisms of period 3 $\alpha\in\mathrm{Aut}(P)$ and $\beta\in\mathrm{Aut}(M)$ such that

$$[\eta^2]=[\mathrm{id}_P]\oplus[\alpha]\oplus[\alpha^2]\oplus 2[\eta],\quad [\alpha\eta]=[\eta\alpha]=[\eta],$$

$$[\hat{\eta}^2]=[\mathrm{id}_M]\oplus[\beta]\oplus[\beta^2]\oplus 2[\hat{\eta}],\quad [\beta\hat{\eta}]=[\hat{\eta}\beta]=[\hat{\eta}].$$



Since $[M:Q]=4$, Theorem 5.3 and Proposition 5.13 show that the principal graph of $Q\subset M$ is also $E_6^{(1)}$. Therefore there exists outer automorphism $\beta_Q\in\mathrm{Aut}(M)$ of period 3 such that

$$[\hat\eta_Q^2]=[\mathrm{id}_M]\oplus[\beta_Q]\oplus[\beta_Q^2]\oplus 2[\hat\eta_Q],\quad [\beta_Q\hat\eta_Q]=[\hat\eta_Q\beta_Q]=[\hat\eta_Q].$$

Since $[\kappa\eta]=[\kappa]\oplus[\kappa\alpha]\oplus[\kappa\alpha^2]$, we have $[\eta\xi]=[\eta\bar\kappa\kappa_Q\pi]=[\xi]\oplus[\alpha\xi]\oplus[\alpha^2\xi]$, and Lemma 5.5 shows $\dim(\kappa\iota\bar\iota\bar\kappa,\kappa\iota\bar\iota\bar\kappa)=8$. If $\mathfrak{Q}$ is in Class III, Corollary 4.6,(3) implies

$$\begin{aligned}3 &= \dim(\hat\eta\theta\hat\eta,\hat\eta\theta\hat\eta)=\dim(\hat\eta^2,\theta\hat\eta^2\theta)\\ &= \dim(\mathrm{id}_M\oplus\beta\oplus\beta^2\oplus 2\hat\eta,\mathrm{id}_M\oplus\theta\beta\theta\oplus\theta\beta^2\theta\oplus 2\theta\hat\eta\theta),\end{aligned}$$

which shows either $[\beta]=[\beta_Q]$ or $[\beta]=[\beta_Q^2]$. Since the second cohomology of the cyclic group $\{\mathrm{id}_M,\beta,\beta^2\}\cong\mathfrak{A}_3$ is trivial, a similar proof, using either [16] or [21], as in the case of $E_7^{(1)}$ works .

Suppose now that $\mathfrak{Q}$ is in Class II. (Note that since the cohomology of the $E_6^{(1)}$ subfactor is not trivial, we cannot apply Corollary 4.6,(4) as before.) Corollary 4.6,(4) shows that $\bar\pi\eta_Q\pi$ is contained in ξ^2, which is

$$[\xi^2]=[\mathrm{id}_P]\oplus[\xi]\oplus[\alpha\xi]\oplus[\alpha^2\xi]\oplus[\eta],$$

and so $[\eta]=[\bar\pi\eta_Q\pi]$. The proof of Theorem 4.3 shows

$$[\kappa\iota\bar\iota\bar\kappa]=[\mathrm{id}_M]\oplus[\hat\eta]\oplus[\kappa_Q\pi\bar\kappa]\oplus[\hat\eta\kappa_Q\pi\bar\kappa].$$

Since $\dim(\kappa_Q\pi\bar\kappa,\kappa_Q\pi\bar\kappa)=\dim(\bar\pi\bar\kappa_Q\kappa_Q\pi,\bar\kappa\kappa)=2$, the endomorphism $\kappa\bar\pi\bar\kappa_Q$ is decomposed two irreducible endomorphism, say ρ and σ, neither of which is an automorphism as we assumed that $\mathfrak{Q}$ is in Class II. By Frobenius reciprocity, the morphisms $\rho\kappa_Q\pi$ and $\sigma\kappa_Q\pi$ contain κ with multiplicity one. Since

$$[\kappa\bar\pi\bar\kappa_Q\kappa_Q\pi]=[\kappa(\mathrm{id}\oplus\eta)]=2[\kappa]\oplus[\beta\kappa]\oplus[\beta^2\kappa],$$

we may assume $[\rho\kappa_Q\pi]=[\kappa]\oplus[\beta\kappa]$ and $[\sigma\kappa_Q\pi]=[\kappa]\oplus[\beta^2\kappa]$. By Frobenius reciprocity again, the morphism $\beta\kappa\bar\pi\bar\kappa_Q$ contains ρ and the morphism $\beta^2\kappa\bar\pi\bar\kappa_Q$ contains σ, and so $\kappa\bar\pi\bar\kappa_Q$ contains $\beta^2\rho$ and $\beta\sigma$. If $[\beta\sigma]=[\sigma]$, we also have $[\beta^2\rho]=[\rho]$ and so $[\beta\kappa\bar\pi\bar\kappa_Q]=[\kappa\bar\pi\bar\kappa_Q]$. However, this would imply

$$\begin{aligned}2 &= \dim(\beta\kappa\bar\pi\bar\kappa_Q,\kappa\bar\pi\bar\kappa_Q)=\dim(\kappa\alpha\bar\pi\bar\kappa_Q,\kappa\bar\pi\bar\kappa_Q)=\dim(\bar\kappa\kappa\alpha,\bar\pi\bar\kappa_Q\kappa_Q\pi)\\ &= \dim(\alpha\oplus\eta,\mathrm{id}_P\oplus\eta),\end{aligned}$$

which is a contradiction. Thus we get $[\rho]=[\beta\sigma]$ and $d(\sigma)=2$ and

$$\begin{aligned}[][\kappa\iota\bar{\iota}\bar{\kappa}] &= [\mathrm{id}_M]\oplus[\hat{\eta}]\oplus[\sigma]\oplus[\beta\sigma]\oplus[\hat{\eta}\sigma]\oplus[\hat{\eta}\beta\sigma] \\ &= [\mathrm{id}_M]\oplus[\hat{\eta}]\oplus[\sigma]\oplus[\beta\sigma]\oplus 2[\hat{\eta}\sigma].\end{aligned}$$

Since $d(\sigma)=2$, the endomorphism $\hat{\eta}^2$ does not contain σ and by Frobenius reciprocity $\hat{\eta}\sigma$ does not contain $\hat{\eta}$. Therefore the multiplicity of $\hat{\eta}$ in $\kappa\iota\bar{\iota}\kappa$ is one, and so the multiplicity of $\hat{\eta}_Q$ is also one by symmetry. However, since $d(\hat{\eta}_Q)=3$ and $[\hat{\eta}]\neq[\hat{\eta}_Q]$, this is impossible. Therefore $\mathfrak{Q}$ is in Class III. ☐

Now the proof of Theorem 5.4 follows from the above propositions, lemmas, and Theorem 5.3.

We don't know if there are infinitely many mutually non-conjugate quadrilaterals of the hyperfinite II_1 factors in Class II.

5.3. **Class III.**

Theorem 5.15. *Let* $\mathfrak{Q}=\begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q\end{array}$ *be a quadrilateral of factors in Class III such that* $\mathfrak{Q}$ *is (5,3)-supertransitive. Then*

1. *If* $\mathfrak{Q}$ *is (6,3)-supertransitive, the principal graphs of* $P\subset M$ *and* $Q\subset M$ *are* A_5 *and those of* $N\subset P$ *and* $N\subset Q$ *are* $E_7^{(1)}$*.*
2. *If* $\mathfrak{Q}$ *is not (6,3)-supertransitive, the dual principal graph of* $P\subset M$ *contains one of the principal graphs of the Asaeda-Haagerup subfactor* [1], *and in consequence* $[M:P]\geq(5+\sqrt{17})/2$*.*

Proof. Assume that the principal graph of $P\subset M$ is not A_5. Since it is neither A_3 nor A_4, the depth of $P\subset M$ is at least 5. We use the same symbols $\hat{\eta}',\kappa',\kappa''$, and η' for $M-M$, $M-P$, and $P-P$ sectors arising from $P\subset M$ as in the proof of Theorem 5.3. Corollary 4.6,(3) implies

$$2\leq\dim(\hat{\eta}\theta\hat{\eta},\hat{\eta}\theta\hat{\eta})=\dim(\mathrm{id}_M\oplus\hat{\eta}\oplus\hat{\eta}',\mathrm{id}_M\oplus\theta\hat{\eta}\theta\oplus\theta\hat{\eta}'\theta)=1+\dim(\hat{\eta}',\theta\hat{\eta}'\theta),$$

which shows that $[\hat{\eta}']=[\theta\hat{\eta}'\theta]$ and the endomorphism $\hat{\eta}\theta\hat{\eta}$ is a direct sum of $\theta\hat{\eta}\theta$ and an irreducible endomorphism, which we call $\hat{\zeta}$. Therefore we have

$$[\kappa\iota\bar{\iota}\bar{\kappa}]=[\mathrm{id}_M]\oplus[\theta]\oplus[\hat{\eta}]\oplus[\theta\hat{\eta}]\oplus[\hat{\eta}\theta]\oplus[\theta\hat{\eta}\theta]\oplus[\hat{\zeta}].$$

Since $\theta\kappa\iota=\kappa\iota$, we have $[\theta\hat{\zeta}]=[\hat{\zeta}\theta]=[\hat{\zeta}]$, which is also self-conjugate.

We claim $[\theta\hat{\eta}']\neq[\hat{\eta}']$. Suppose, on the contrary, that $[\theta\hat{\eta}']=[\hat{\eta}']$ holds. Then

$$[\kappa']\oplus[\kappa'']=[\hat{\eta}'\kappa]=[\theta\hat{\eta}'\kappa]=[\theta\kappa']\oplus[\theta\kappa''].$$

If $[\theta\kappa']=[\kappa'']$, then the principal graph of $P\subset M$ would be A_9 and θ would commute with $\hat{\eta}$, which is a contradiction. Thus $[\theta\kappa']=[\kappa']$ holds. However, this implies

$$[\hat{\eta}]\oplus[\hat{\eta}']=[\kappa'\bar{\kappa}]=[\theta\kappa'\bar{\kappa}]=[\theta\hat{\eta}]\oplus[\theta\hat{\eta}']=[\theta\hat{\eta}]\oplus[\hat{\eta}'],$$

and $[\theta\hat{\eta}]=[\hat{\eta}]$, which is a contradiction again. Therefore the claim is proven.

Let $[\kappa''\bar{\kappa}] = [\hat{\eta}'] \oplus \bigoplus_{i=1}^{n} m_i[\hat{\eta}_i]$ be the irreducible decomposition. We compute $\hat{\eta}\theta\hat{\eta}^2$ in two ways:

$$[\hat{\eta}\theta\hat{\eta}][\hat{\eta}] = [\theta\hat{\eta}\theta\hat{\eta}] \oplus [\hat{\zeta}\hat{\eta}] = [\hat{\eta}\theta] \oplus [\theta\hat{\zeta}] \oplus [\hat{\zeta}\hat{\eta}] = [\hat{\eta}\theta] \oplus [\hat{\zeta}] \oplus [\hat{\zeta}\hat{\eta}],$$

$$\begin{aligned}[\hat{\eta}\theta][\hat{\eta}^2] &= [\hat{\eta}\theta] \oplus [\hat{\eta}\theta\hat{\eta}] \oplus [\hat{\eta}\theta\hat{\eta}'] = [\hat{\eta}\theta] \oplus [\theta\hat{\eta}\theta] \oplus [\hat{\zeta}] \oplus [\hat{\eta}\hat{\eta}'\theta] \\ &= 2[\hat{\eta}\theta] \oplus [\theta\hat{\eta}\theta] \oplus [\hat{\eta}'\theta] \oplus [\hat{\zeta}] \oplus \bigoplus_{i=1}^{n} m_i[\hat{\eta}_i\theta],\end{aligned}$$

which implies

$$[\hat{\zeta}\hat{\eta}\theta] = [\hat{\eta}] \oplus [\theta\hat{\eta}] \oplus [\hat{\eta}'] \oplus \bigoplus_{i=1}^{n} m_i[\hat{\eta}_i].$$

Since $[\theta][\hat{\zeta}\hat{\eta}\theta] = [\hat{\zeta}\hat{\eta}\theta]$ and $[\theta][\hat{\eta}'] \neq [\hat{\eta}']$, we may assume that $m_1 = 1$, $[\theta\hat{\eta}'] = [\hat{\eta}_1]$ and

$$\bigoplus_{i=2}^{n} m_i[\theta\hat{\eta}_i] = \bigoplus_{i=2}^{n} m_i[\hat{\eta}_i].$$

By Frobenius reciprocity, the endomorphism $\hat{\eta}'\theta\hat{\eta}$ contains $\hat{\zeta}$ with multiplicity one. On the other hand, we have

$$[\hat{\eta}'\theta\hat{\eta}] = [\theta\hat{\eta}'\hat{\eta}] = [\theta\hat{\eta}] \oplus [\theta\hat{\eta}'] \oplus [\hat{\eta}'] \oplus \bigoplus_{i=2}^{n} m_i[\theta\hat{\eta}_i] = [\theta\hat{\eta}] \oplus [\theta\hat{\eta}'] \oplus [\hat{\eta}'] \oplus \bigoplus_{i=2}^{n} m_i[\hat{\eta}_i].$$

Corollary 4.6,(3) shows $[\hat{\zeta}] \neq [\theta\hat{\eta}]$, and $[\theta\hat{\eta}'] \neq [\hat{\eta}']$ implies $[\hat{\zeta}] \neq [\hat{\eta}'], [\theta\hat{\eta}']$ as $[\theta\hat{\zeta}] = [\hat{\zeta}]$ holds. Thus we may assume that $m_2 = 1$ and $[\hat{\zeta}] = [\hat{\eta}_2]$. This in particular shows that $\mathfrak{Q}$ cannot be $(6,3)$-supertransitive.

Now we have

$$[\hat{\zeta}\hat{\eta}] = [\hat{\eta}\theta] \oplus [\theta\hat{\eta}\theta] \oplus [\hat{\eta}'] \oplus [\theta\hat{\eta}'] \oplus [\hat{\zeta}] \oplus \bigoplus_{i=3}^{n} m_i[\hat{\eta}_i\theta],$$

and $[\hat{\zeta}] \neq [\hat{\eta}_i\theta]$ for $i \geq 3$. Since the multiplicity of $\hat{\zeta}$ in $\hat{\zeta}\hat{\eta}$ is one, there are exactly two vertices in the dual principal graph of $P \subset M$ connected to $\hat{\zeta}$, one of which is κ''. Therefore there exists an irreducible morphism κ''' such that $[\hat{\zeta}\kappa] = [\kappa''] \oplus [\kappa''']$ with $[\kappa'''] \neq [\kappa'']$. We claim $[\kappa'''] = [\theta\hat{\eta}\theta\kappa] = [\hat{\eta}\theta\kappa]$. Indeed, computing $\dim(\theta\hat{\eta}\theta\kappa, \theta\hat{\eta}\theta\kappa)$, $\dim(\hat{\eta}\theta\kappa, \hat{\eta}\theta\kappa)$, and $\dim(\hat{\eta}\theta\kappa, \theta\hat{\eta}\theta\kappa)$, we can show that $\theta\hat{\eta}\theta\kappa$ and $\hat{\eta}\theta\kappa$ are irreducible and $[\theta\hat{\eta}\theta\kappa] = [\hat{\eta}\theta\kappa]$. Since $\theta\hat{\eta}\theta\kappa\bar{\kappa}$ contains $\hat{\zeta}$ with multiplicity one, Frobenius reciprocity implies that $\hat{\zeta}\kappa$ contains $\theta\hat{\eta}\theta\kappa$. If $[\theta\hat{\eta}\theta\kappa] = [\kappa'']$ we would have $d(\hat{\eta})d(\kappa) = d(\kappa'')$, which implies $[M : P] = 4$. Since $P \subset M$ is 5-supertransitive, the principal graph of $P \subset M$ should be $E_8^{(1)}$. However, since the $M-M$ sectors arising from $P \subset M$ is noncommutative, we get a contradiction. Therefore the claim is proven.

The above computation shows that the dual principal graph is as follows.



which contains one of the principal graphs of the Asaeda-Haagerup subfactor [1]. □

Remark 5.16. We do not known whether there exists a (5,3)-supertransitive quadrilateral of factors in Class III with $[M : P] = (5 + \sqrt{17})/2$. If such a quadrilateral of hyperfinite II_1 factors exists, it must be unique up to conjugacy. Indeed, the above argument shows that the inclusion $P \subset M$ must be Asaeda-Haagerup subfactor and $\xi = \bar{\kappa}\theta\kappa$ is determined by $P \subset M$. Since $N \subset P$ is 3-supsertransitive, a Q-system for $\mathrm{id}_P \oplus \xi$ is unique up to equivalence and M is uniquely determined by $P \subset M$ up to inner conjugacy in P. Since $Q = \theta(P)$, where the representative θ of the sector $[\theta]$ is chosen so that the restriction $\theta|_N$ is trivial, the quadrilateral is uniquely determined by $P \subset M$. For existence, the same criterion using a Q-system as in the case of the Haagerup subfactor below (Theorem 5.19) should work in principle, though it involves heavy computation in practice.

Theorem 5.15 and the propositions and lemmas in the previous subsection imply the following theorem:

Theorem 5.17. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *be a quadrilateral of factors in Class III such that* $[M : P] \leq 4$. *Then one of the following holds:*

1. *The principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* A_5 *and those of* $N \subset P$ *and* $N \subset Q$ *are* $E_7^{(1)}$. *The quadrilateral is given by the construction in Example 2.6 with* $\mathfrak{S}_{\{1,2,3,4\}}$.
2. *The principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* $E_6^{(1)}$. *The quadrilateral is given by the construction in Example 2.6 with* $\mathfrak{A}_{\{1,2,3,4,5\}}$.
3. *The principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* $E_7^{(1)}$. *The quadrilateral is given by the construction in Example 2.6 with* $\mathfrak{S}_{\{1,2,3,4,5\}}$.

In each case, such a quadrilateral of the hyperfinite II_1 *factors exists and is unique up to conjugacy.*

Corollary 5.18. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *be an irreducible noncommuting (6,3)-supertransitive quadrilateral of factors. Then one of the following holds:*

1. *The principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* A_3 *and those of* $N \subset P$ *and* $N \subset Q$ *are* A_5.

(2) *The principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* A_4 *and those of* $N \subset P$ *and* $N \subset Q$ *are* D_6.

(3) *The principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* A_5 *and those of* $N \subset P$ *and* $N \subset Q$ *are* $E_7^{(1)}$.

(4) *The principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* A_7 *and those of* $N \subset P$ *and* $N \subset Q$ *are* A_7.

In each case, such a quadrilateral of the hyperfinite II_1 *factors exists and is unique up to conjugacy.*

Thanks to Example 2.6, there are countably many mutually non-conjugate (3,3)-supertransitive quadrilaterals of the hyperfinite II_1 factors in Class III.

5.4. **Class IV.**

Theorem 5.19. *There exists a quadrilateral of factors* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *in Class IV with* $[M:P] = (3+\sqrt{13})/2$. *Such a quadrilateral of the hyperfinite* II_1 *factors is unique up to flip conjugacy.*

Proof. Let $P \subset M$ be the Haagerup subfactor whose principal graph and the dual principal graph are as follows.



It is known that θ and $\hat{\eta}$ satisfies the following relation [1]:

$$[\theta^3] = [\mathrm{id}_M], \quad [\theta^{-1}\hat{\eta}] = [\hat{\eta}\theta], \quad [\hat{\eta}^2] = [\mathrm{id}_M] \oplus [\hat{\eta}] \oplus [\theta\hat{\eta}] \oplus [\theta^2\hat{\eta}].$$

In the Appendix, we will show that there exists a unique Q-system for $\mathrm{id}_P \oplus \bar{\kappa}\theta\kappa$ up to equivalence. Therefore there exists a unique subfactor N of P up to inner conjugacy such that $[\iota\bar{\iota}] = [\mathrm{id}_P] \oplus [\bar{\kappa}\theta\kappa]$, where $\iota : N \hookrightarrow P$ is the inclusion map. Since

$$\dim(\kappa\iota, \kappa\iota) = \dim(\bar{\kappa}\kappa, \iota\bar{\iota}) = \dim(\mathrm{id}_P \oplus \eta, \mathrm{id}_P \oplus \bar{\kappa}\theta\kappa) = 1,$$

the inclusion $N \subset M$ is irreducible. Since

$$\begin{aligned} [\kappa\iota\bar{\iota}\bar{\kappa}] &= [\kappa\bar{\kappa}] \oplus [\kappa\bar{\kappa}\theta\kappa\bar{\kappa}] = [\mathrm{id}_M] \oplus [\hat{\eta}] \oplus [(\mathrm{id}_M \oplus \hat{\eta})\theta(\mathrm{id}_M \oplus \hat{\eta})] \\ &= [\mathrm{id}_M] \oplus [\hat{\eta}] \oplus [\theta] \oplus [\theta\hat{\eta}] \oplus [\theta^2\hat{\eta}] \oplus [\theta^2\hat{\eta}^2] \\ &= [\mathrm{id}_M] \oplus [\theta] \oplus [\theta^2] \oplus 2[\hat{\eta}] \oplus 2[\theta\hat{\eta}] \oplus 2[\theta^2\hat{\eta}], \end{aligned}$$

we can take representative θ of the sector $[\theta]$ such that $\theta\kappa\iota = \kappa\iota$. Then θ^3 is an inner automorphism satisfying $\theta^3\kappa\iota = \kappa\iota$ and so $\theta^3 = \mathrm{id}_M$ as $N \subset M$ is irreducible. We set $Q = \theta(P)$. Using a similar argument as in the proof of Theorem 5.1, we can show that $\begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ is the desired quadrilateral. Uniqueness up to flip conjugacy follows from uniqueness of the Q-system for $\mathrm{id} \oplus \bar{\kappa}\theta\kappa$. □

We do not know if there exists a quadrilateral of the hyperfinite II_1 factors in Class IV different from the above example.

6. Classification II

Theorem 6.1. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *be an irreducible noncommuting quadrilateral of factors such that the indices of all the elementary subfactors are less than or equal to 4. Then one of the following occurs:*

1. *The principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* A_3 *and those of* $N \subset P$ *and* $N \subset Q$ *are* A_5*. In this case* N *is the fixed point algebra of an outer action of the symmetric group* $\mathfrak{S}_3$ *on* M*.*
2. *The principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* A_4 *and those of* $N \subset P$ *and* $N \subset Q$ *are* D_6*.*
3. *The principal graphs of all the elementary subfactors are* A_7*.*
4. *The principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* A_3 *and those of* $N \subset P$ *and* $N \subset Q$ *are* $D_6^{(1)}$*. In this case* N *is the fixed point algebra of an outer action of the dihedral group* $\mathfrak{D}_8$ *of order 8 on* M*.*
5. *The principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* D_4 *and those of* $N \subset P$ *and* $N \subset Q$ *are* $E_6^{(1)}$*. In this case* N *is the fixed point algebra of an outer action of the alternating group* $\mathfrak{A}_4$ *on* M*.*
6. *The principal graphs of* $P \subset M$ *and* $Q \subset M$ *are* A_5 *and those of* $N \subset P$ *and* $N \subset Q$ *are* $E_7^{(1)}$*. In this case the quadrilateral* $\mathfrak{Q}$ *is given by the constructions in Example 2.6 with* $\mathfrak{S}_{\{1,2,3,4\}}$*.*
7. *The principal graphs of all the elementary subfactors are* $E_7^{(1)}$*.*

In each case, such a quadrilateral of the hyperfinite II_1 *factors is unique up to conjugacy.*

We use the symbols $\iota = \iota_{P,N}$, $\iota_Q = \iota_{Q,N}$, $\kappa = \iota_{M,P}$, $\kappa_Q = \iota_{M,Q}$. When $N \subset P$ is 2-supertransitive, we also use the symbols ξ and $\hat{\xi}$ as in Section 4.

Lemma 6.2. *Let* $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *be an irreducible noncommuting quadrilateral of factors such that the indices of all the elementary subfactors are less than or equal to 4. Then*

1. *The principal graph of* $N \subset P$ *is neither* A_3*,* D_4 *nor* $D_4^{(1)}$*.*

(2) *If the principal graph of* $N \subset P$ *is* $E_6^{(1)}$, *then that of* $P \subset M$ *is* D_4. *There exists an outer action action of the alternating group* $\mathfrak{A}_4$ *on* M *such that* N *is the fixed point algebra of the action.*

(3) *If the principal graph of* $N \subset P$ *is* $D_n^{(1)}$, *then* $n = 6$ *and the principal graph of* $P \subset M$ *is* A_3. *There exists an outer action action of the dihedral group* $\mathfrak{D}_8$ *of order 8 on* M *such that* N *is the fixed point algebra of the action.*

(4) *If* $N \subset P$ *and* $N \subset Q$ *are 2-supertransitive, then the principal graph of* $P \subset M$ *is not* $D_n^{(1)}$.

Proof. (1) follows from Remark 3.15.

(2) Since the principal graph of $N \subset P$ is $E_6^{(1)}$, there exists an outer automorphism $\alpha \in \mathrm{Aut}(P)$ of order 3 such that

$$[\xi^2] = [\mathrm{id}_P] \oplus [\alpha] \oplus [\alpha^2] \oplus 2[\xi].$$

The proof of Lemma 4.1 shows that $\dim(\bar{\kappa}\kappa, \xi) = 0$ and

$$2 \leq \dim(\bar{\kappa}\kappa, \xi^2) = \dim(\bar{\kappa}\kappa, \mathrm{id}_P \oplus \alpha \oplus \alpha^2).$$

Therefore $\bar{\kappa}\kappa$ contains α and so the principal graph of $P \subset M$ is D_4. The rest of the statement follows from [16], [21].

(3) Thanks to (1), the number n is at least 5 and we have irreducible decomposition

$$[\iota\bar{\iota}] = [\mathrm{id}_P] \oplus [\alpha] \oplus [\xi], \quad [\bar{\iota}\iota] \oplus [\tau] \oplus [\hat{\xi}],$$

where $d(\xi) = d(\hat{\xi}) = 2$ and $\alpha \in \mathrm{Aut}(P)$ and $\tau \in \mathrm{Aut}(N)$ are outer automorphisms of order 2 satisfying $[\alpha\xi] = [\xi]$, $[\tau\hat{\xi}] = [\hat{\xi}]$. Since $[\kappa\iota\tau] = [\kappa\iota]$, the crossed product $N \rtimes_\tau \mathbb{Z}/2\mathbb{Z}$ is identified with the intermediate subfactor of $N \subset P$ generated by N and a unitary $u \in M$ satisfying $uxu^* = \tau(x)$ for all $x \in N$. If $\bar{\iota}_Q\iota_Q$ contained τ as well, the two inclusions $N \subset P$ and $N \subset Q$ would have a common intermediate subfactor $N \rtimes \tau\mathbb{Z}/2\mathbb{Z}$, which is a contradiction as we have $P \cap Q = N$. Since $\mathfrak{Q}$ is noncommuting $\bar{\iota}_Q\iota_Q$ must contain $\hat{\xi}$ and the principal graph of $N \subset Q$ is either A_5 or $D_m^{(1)}$ with $m \geq 5$.

Assume that the principal graph of $N \subset Q$ is $D_m^{(1)}$. Then we have the irreducible decomposition $[\bar{\iota}_Q\iota_Q] = [\mathrm{id}_N] \oplus [\tau_Q] \oplus [\hat{\xi}]$ such that $[\tau_Q] \neq [\tau]$ and τ_Q is contained in $\hat{\xi}^2$. If n were not equal to 6, the endomorphism $\hat{\xi}^2$ would contain exactly two automorphisms id_P and τ, which would imply $[\tau] = [\tau_Q]$, a contradiction. Thus we get $n = m = 6$ and $[\hat{\xi}^2] = [\mathrm{id}_N] \oplus [\tau] \oplus [\tau_Q] \oplus [\tau']$, where τ' is an automorphisms of N. It is known that there are 4 different subfactors with the principal graph $D_6^{(1)}$ [25]. We show that the only one case among them is possible.

Since the dual principal graph of $N \subset P$ is also $D_6^{(1)}$, we have $[\xi^2] = [\mathrm{id}] \oplus [\alpha] \oplus [\alpha_1] \oplus [\alpha_2]$ where α_1 and α_2 are automorphisms of P. The proof of Lemma 4.1 shows

$$2 \leq \dim(\bar{\iota}\bar{\kappa}\kappa\iota, \hat{\xi}) = 1 + \dim(\bar{\kappa}\kappa, \alpha_1 \oplus \alpha_2).$$

If $\bar{\kappa}\kappa$ contained both α_1 and α_2, it would contain α too, which contradicts irreducibility of $N \subset M$. Thus we may assume that $\bar{\kappa}\kappa$ contains only α_1, whose period must be two in consequence. Therefore either $[M : P] = 2$ or the principal graph of $P \subset M$ is

$D_k^{(1)}$. Let $L = P \rtimes_{\alpha_1} \mathbb{Z}/2\mathbb{Z}$, which is regarded as an intermediate subfactor of P and M. Then it is known [26, Chapter 8] that $N \subset L$ is given by the fixed point algebra of an outer action of either $\mathfrak{D}_8$ or the Kac-Paljutkin algebra on L and

$$[\bar{\iota}\bar{\sigma}\sigma\iota] = [\mathrm{id}_N] \oplus [\tau] \oplus [\tau_Q] \oplus [\tau'] \oplus 2[\hat{\xi}],$$

where we set $\sigma = \iota_{L,P}$. We claim that Q is an intermediate subfactor of $N \subset L$. Since $d(\hat{\xi}) = 2$, Theorem 2.9,(1) shows that the multiplicity of $\hat{\xi}$ in $\bar{\iota}\bar{\kappa}\kappa\iota$ is two and $\bar{\iota}\bar{\kappa}\kappa\iota$ contains automorphisms with multiplicity at most one. On the other hand, the endomorphism $\bar{\iota}_Q\iota_Q$ is contained in $\bar{\iota}\bar{\sigma}\sigma\iota$, and so [28, Theorem 3.9] shows the claim. Since $\mathfrak{Q}$ is a quadrilateral we get $M = L$. Since $[M : P] = [M : Q] = 2$, there exist period two automorphisms $\beta, \beta_1 \in \mathrm{Aut}(M)$ such that $P = M^{\beta}$ and $Q = M^{\beta_1}$. If $N \subset M$ were given by the Kac-Paljutkin algebra, we would have

$$[\kappa\iota\bar{\iota}\bar{\kappa}] = [\mathrm{id}_M] \oplus [\beta_1] \oplus [\beta_2] \oplus [\beta_1\beta_2] \oplus 2[\rho],$$

with $d(\rho) = 2$, and the two automorphisms β_1 and β_2 would generate a group G of order 4 in $\mathrm{Aut}(M)$. However, this means $P \cap Q = M^G \neq N$, which is a contradiction. Therefore N is the fixed point algebra of an outer action of $\mathfrak{D}_8$.

Now we assume that the principal graph of $N \subset P$ is A_5 and we show that $\mathfrak{Q}$ would be commuting, which contradicts the assumption. Since $[M : P][P : N] = [M : Q][Q : N]$ we have $[M : P] = 3$ and $[M : Q] = 4$. In this case, we have

$$[\bar{\iota}_Q\iota_Q] = [\mathrm{id}_N] \oplus [\hat{\xi}], \quad [\iota_Q\bar{\iota}_Q] = [\mathrm{id}_Q] \oplus [\xi_Q],$$

$$[\hat{\xi}^2] = [\mathrm{id}_N] \oplus [\tau] \oplus [\hat{\xi}], \quad [\tau^2] = [\mathrm{id}_M], \quad [\tau\hat{\xi}] = [\hat{\xi}\tau] = [\hat{\xi}],$$

$$[\xi_Q^2] = [\mathrm{id}_Q] \oplus [\alpha_Q] \oplus [\xi_Q], \quad [\alpha_Q^2] = [\mathrm{id}_Q], \quad [\alpha_Q\xi_Q] = [\xi_Q\alpha_Q] = [\xi_Q].$$

As in [18, Lemma 3.2], we can choose representatives of $[\alpha_Q]$ and $[\tau]$ such that $\alpha_Q^2 = \mathrm{id}_Q$, $\tau^2 = \mathrm{id}_N$, and $\alpha_Q\iota_Q = \iota_Q\tau$ hold.

We claim $[\kappa_Q\alpha_Q] = [\kappa_Q]$. Indeed, since $\kappa\iota = \kappa_Q\iota_Q$ and $[\iota\tau] = [\iota]$, we have $[\kappa_Q\alpha_Q\iota_Q] = [\kappa_Q\iota_Q\tau] = [\kappa_Q\iota_Q]$ and

$$1 = \dim(\kappa_Q\alpha_Q\iota_Q, \kappa_Q\iota_Q) = \dim(\kappa_Q\alpha_Q, \kappa_Q\iota_Q\bar{\iota}_Q) = \dim(\kappa_Q\alpha_Q, \kappa_Q \oplus \kappa_Q\xi_Q).$$

Since we have

$$\dim(\kappa_Q\alpha_Q, \kappa_Q\xi_Q) = \dim(\kappa_Q, \kappa_Q\xi_Q\alpha_Q) = \dim(\kappa_Q, \kappa_Q\xi_Q) = \dim(\bar{\kappa}_Q\kappa_Q, \xi_Q) = 0,$$

we get the claim.

The claim implies that we can regard $L := Q \rtimes_{\alpha_Q} \mathbb{Z}/2\mathbb{Z}$ as an intermediate subfactor of $Q \subset M$. On the other hand, we can regard $R := N \rtimes_{\tau} \mathbb{Z}/2\mathbb{Z}$ as an intermediate subfactor of $N \subset P$. Since we have $\alpha_Q\iota_Q = \iota_Q\tau$, we have the inclusion $R \subset L$. Moreover [18, Theorem 3.1] shows that N is the fixed point algebra of an outer action of $\mathfrak{S}_3$ on L. In particular $\begin{matrix} R & \subset & L \\ \cup & & \cup \\ N & \subset & Q \end{matrix}$ is a commuting square as we have $[L : R] = 3$ and $[L : Q] = 2$. By Galois correspondence, the subfactor R is the fixed point subalgebra of L under a subgroup of $\mathfrak{S}_3$ isomorphic to $\mathbb{Z}/3\mathbb{Z}$, and by duality, there exists an order three outer automorphism $\beta \in \mathrm{Aut}(R)$ such that $L = R \rtimes_{\beta} \mathbb{Z}/3\mathbb{Z}$.

Since $[P : R] = 2$, there exits an order two automorphism $\varphi \in \mathrm{Aut}(R)$ such that $P = R \rtimes_\varphi \mathbb{Z}/2\mathbb{Z}$. Since $\overline{\iota_{M,R}}\iota_{M,R}$ contains φ and β and $[M : R] = 6$, we have $M = N \rtimes \Gamma$, where Γ is the group generated by $[\varphi]$ and $[\beta]$, whose order must be 6. In particular $\begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ R & \subset & L \end{array}$ is a commuting square. Therefore we have $E_Q E_P = E_Q E_L E_P = E_Q E_R = E_N$, which shows that $\mathfrak{Q}$ is commuting, a contradiction. Thus the statement is proven.

(4) Suppose that $N \subset P$ and $N \subset Q$ are 2-supertransitive and the principal graph of $P \subset M$ is $D_n^{(1)}$. If $\mathfrak{Q}$ were cocommuting, Corollary 3.12 would imply $4 = [M : P] \leq [P : N] - 1$ which is a contradiction. If $\mathfrak{Q}$ were noncocommuting, the statement (3) applied to the dual quadrilateral $\hat{\mathfrak{Q}}$ would imply $M = N \rtimes \mathfrak{D}_8$. However, this shows that $\mathfrak{Q}$ is commuting and we get a contradiction. Therefore the lemma is proven. □

Proof of Theorem 6.1. Assume first that $N \subset P$ and $N \subset Q$ are 2-supertransitive. Then Lemma 6.2,(4) shows that $P \subset M$ is either 2-supertransitive or $M = P \rtimes G$ where G is a finite abelian group of order 2,3, or 4. If the latter occurs, then the proof of Lemma 4.1 implies that ξ^2 contains a non-trivial automorphism, and so the principal graph is either A_5, E_6, or $E_6^{(1)}$. Applying Lemma 6.2,(1) to the dual quadrilateral $\hat{\mathfrak{Q}}$, we see that $\mathfrak{Q}$ is cocommuting. Since the E_6 subfactor has trivial second cohomology, Corollary 3.12 implies that E_6 never occurs. For the same reason, we have $[M : P] = [P : N] - 1 = 2$ for A_5 and we get case (1) from Theorem 2.7. For $E_6^{(1)}$, we get case (5) from Lemma 6.2,(2). If $P \subset M$ and $Q \subset M$ are 2-supertransitive, either Theorem 4.3 or Lemma 6.2,(2) is applied and we get cases (1),(2),(3),(6), and (7) from Theorem 5.2, 5.4, and 5.17.

Assume now that $N \subset P$ is not 2-supertransitive. Then Lemma 6.2,(1) implies that the principal graph of $N \subset P$ is $D_n^{(1)}$ with $n \geq 5$, and so Lemma 6.2,(4) implies the case (4). □

Remark 6.3. Except for the case (4), we have $\Theta(P, Q) = \cos^{-1} 1/([P : Q] - 1)$ thanks to Lemma 2.5 and Corollary 3.11. In the case (4), we have $\Theta(P, Q) = \pi/4$ thanks to [54, Theorem 6.1].

7. α-INDUCTION AND GHJ PAIRS

7.1. **α-induction and angles.** Let $\mathcal{N}$ be a properly infinite factor and $\{\lambda_i\}_{i \in I}$ be a finite system of irreducible endomorphisms in $\mathrm{End}_0(\mathcal{N})$. We assume:
(1) there exists $0 \in I$ such that $\lambda_0 = \mathrm{id}_{\mathcal{N}}$,
(2) for $i \neq j \in I$, we have $[\lambda_i] \neq [\lambda_j]$,
(3) for each $i \in I$, there exists $\bar{i} \in I$ such that $[\overline{\lambda_i}] = [\lambda_{\bar{i}}]$,
(4) there exist non-negative integers N_{ij}^k such that

$$[\lambda_i \lambda_j] = \bigoplus_{k \in I} N_{ij}^k [\lambda_k],$$

(5) the system $\{\lambda_i\}_{i\in I}$ has a braiding $\{\varepsilon(\lambda_i,\lambda_j)\}_{i,j\in I}$ (see [3, Definition 2.2] for the definition),
(6) $\mathcal{N}\subset\mathcal{M}$ is an irreducible inclusion of factors of finite index such that

$$[\overline{\iota_{\mathcal{M},\mathcal{N}}}\iota_{\mathcal{M},\mathcal{N}}]=\bigoplus_{j\in J}n_j[\lambda_j],$$

where J is a subset of I and n_j is a positive integer.

We naturally extend the braiding to the category generated by $\{\lambda_i\}_{i\in I}$ and use the same symbol $\varepsilon(\rho,\sigma)$ for the extension. We set $\varepsilon^+(\rho,\sigma)=\varepsilon(\rho,\sigma)$ and $\varepsilon^-(\rho,\sigma)=\varepsilon(\sigma,\rho)^*$. For simplicity, we use the following notation: $\nu=\iota_{\mathcal{M},\mathcal{N}}$, $\gamma=\nu\bar{\nu}\in\mathrm{End}_0(\mathcal{M})$, $\hat{\gamma}=\bar{\nu}\nu\in\mathrm{End}_0(\mathcal{N})$. Then α-induction $\alpha^{\pm}_{\lambda_i}\in\mathrm{End}_0(\mathcal{M})$ is defined by

$$\alpha^{\pm}_{\lambda_i}=\bar{\nu}^{-1}\cdot\mathrm{Ad}\varepsilon^{\pm}(\lambda_i,\hat{\gamma})\cdot\lambda_i\cdot\bar{\nu}.$$

(See [44], [58].) Note that for $x\in\mathcal{N}$ we have $\bar{\nu}(\nu(x))=\hat{\gamma}(x)$ and $\alpha^{\pm}_{\lambda_i}(x)=\lambda_i(x)$, which means $\alpha^{\pm}_{\lambda_i}\nu=\nu\lambda_i$.

Let $\mathcal{H}_j=(\nu,\nu\lambda_j)$. Then thanks to Theorem 2.9, we have

$$\mathcal{M}=\bigoplus_{j\in J}\mathcal{N}\mathcal{H}_j$$

as a linear space. We choose an orthonormal basis $\{t(j)_k\}_{k=1}^{n_j}$ of $\mathcal{H}_j$.

Lemma 7.1. *Let the notation be as above. Then for $t\in\mathcal{H}_j$ we have*

$$\alpha^{\pm}_{\lambda_i}(t)=\varepsilon^{\pm}(\lambda_i,\lambda_j)^*t.$$

Proof. Since $\bar{\nu}(t)\in(\hat{\gamma},\hat{\gamma}\lambda_j)$, the statement follows from the equation

$$\varepsilon(\lambda_i,\hat{\gamma})\lambda_i(\bar{\nu}(t))\varepsilon(\lambda_i,\hat{\gamma})^*=$$



$$=\hat{\gamma}(\varepsilon(\lambda_i,\lambda_j)^*)\bar{\nu}(t).$$

□

From the above lemma, it is easy to show $(\rho,\sigma)\subset(\alpha^{\pm}_{\rho},\alpha^{\pm}_{\sigma})$.

We fix $i_0\in I$ and set $\lambda=\lambda_{i_0}$, $M=\mathcal{M}$, $P=\alpha^+_{\lambda}(\mathcal{M})$, $Q=\alpha^-_{\lambda}(\mathcal{M})$, $N=P\cap Q$, and $R=\lambda(\mathcal{N})$. Since $\alpha^{\pm}_{\lambda}\nu=\nu\lambda$, we have $R\subset N$. In general these two factors do not coincide and we give a description of N now.

We set

$$J_{i_0}=\{j\in J;\ \varepsilon(\lambda_j,\lambda)\varepsilon(\lambda,\lambda_j)\text{ is a scalar}\},$$

and set $m_j=\varepsilon(\lambda_j,\lambda)\varepsilon(\lambda,\lambda_j)$ for $j\in J_{i_0}$. Recall that $\phi_\rho=r_\rho^*\bar{\rho}(\cdot)r_\rho$ is the standard left inverse of ρ. Since $\varepsilon(\lambda_j,\lambda)\varepsilon(\lambda,\lambda_j)\in(\lambda\lambda_j,\lambda\lambda_j)$, we have $\phi_\lambda(\varepsilon(\lambda_j,\lambda)\varepsilon(\lambda,\lambda_j))\in$

(λ_j, λ_j), which is a scalar. It is easy to show that $j \in J_{i_0}$ if and only if

$$|\phi_\lambda(\varepsilon(\lambda_j, \lambda)\varepsilon(\lambda, \lambda_j))| = 1.$$

Lemma 7.2. *Let the notation be as above. Then*

$$N = \alpha_\lambda^+(\sum_{j \in J_{i_0}} \mathcal{N}\mathcal{H}_j) = \alpha_\lambda^-(\sum_{j \in J_{i_0}} \mathcal{N}\mathcal{H}_j).$$

In particular, the von Neumann algebra N is a factor.

Proof. Let $x, y \in \mathcal{M}$ satisfy $\alpha_\lambda^+(x) = \alpha_\lambda^-(y)$. Then Theorem 2.9 shows that we have expansion of x and y as

$$x = \sum_{j \in J} \sum_{k=1}^{n_j} a(j)_k t(j)_k,$$

$$y = \sum_{j \in J} \sum_{k=1}^{n_j} b(j)_k t(j)_k,$$

where $a(j)_k$ and $b(j)_k$ are elements in $\mathcal{N}$ uniquely determined by x and y respectively. Thus we get

$$\sum_{j \in J} \sum_{k=1}^{n_j} \lambda(a(j)_k)\varepsilon(\lambda, \lambda_j)^* t(j)_k = \alpha_\lambda^+(x) = \alpha_\lambda^-(y) = \sum_{j \in J} \sum_{k=1}^{n_j} \lambda(b(j)_k)\varepsilon(\lambda_j, \lambda) t(j)_k,$$

which implies $\lambda(a(j)_k)\varepsilon(\lambda, \lambda_j)^* = \lambda(b(j)_k)\varepsilon(\lambda_j, \lambda)$ for all $j \in J$ and $1 \leq k \leq n_j$. Applying ϕ_λ to both $\lambda(a(j)_k) = \lambda(b(j)_k)\varepsilon(\lambda_j, \lambda)\varepsilon(\lambda, \lambda_j)$ and $\lambda(a(j)_k)(\varepsilon(\lambda_j, \lambda)\varepsilon(\lambda, \lambda_j))^* = b(j)_k$, we get

$$a(j)_k = b(j)_k \phi_\lambda(\varepsilon(\lambda_j, \lambda)\varepsilon(\lambda, \lambda_j)),$$

$$a(j)_k \overline{\phi_\lambda(\varepsilon(\lambda_j, \lambda)\varepsilon(\lambda, \lambda_j))} = b(j)_k,$$

which shows that $a(k)_j \neq 0$ only if $|\phi_{\lambda_j}(\varepsilon(\lambda, \lambda_j)\varepsilon(\lambda_j, \lambda))| = 1$, which is equivalent to $j \in J_{i_0}$. On the other hand, when $j \in J_{i_0}$ we have $\varepsilon(\lambda_j, \lambda) = m_j \varepsilon(\lambda, \lambda_j)^*$ and

$$\alpha_\lambda^+(\sum_{j \in J_{i_0}} \sum_{k=1}^{n_j} a(j)_k t(j)_k) = \alpha_\lambda^-(\sum_{j \in J_{i_0}} \sum_{k=1}^{n_j} m_j^{-1} a(j)_k t(j)_k), \quad \forall a(j)_k \in \mathcal{N},$$

which shows the first statement. Since $\sum_{j \in J_{i_0}} \mathcal{N}\mathcal{H}_j$ is an intermediate von Neumann algebra between $\mathcal{N}$ and $\mathcal{M}$, it must be a factor, and so is N. □

Assume that $P \subset M$ has no non-trivial intermediate subfactor. Then $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ is a quadrilateral though it is not irreducible in general.

Theorem 7.3. *Let the notation be as above. Then*

$$\mathrm{Ang}(P, Q) = \{\cos^{-1} |\phi_\lambda(\varepsilon(\lambda_j, \lambda)\varepsilon(\lambda, \lambda_j))|;\ j \in J\} \setminus \{0, \frac{\pi}{2}\}.$$

In particular, when the braiding is non-degenerate, we have

$$\mathrm{Ang}(P,Q)=\{\cos^{-1}\frac{|S_{00}S_{i_0j}|}{|S_{0i_0}S_{0j}|};\ j\in J\}\setminus\{0,\frac{\pi}{2}\},$$

where S_{ij} *is as in* [3, Proposition 2.4].

Proof. Since $\mathcal{M}=\sum_{j\in J}\mathcal{N}\mathcal{H}_j$, we have

$$P=\alpha^+_\lambda(\mathcal{M})=\sum_{j\in J}\lambda(\mathcal{N})\alpha^+_\lambda(\mathcal{H}_j),$$

$$Q=\alpha^-_\lambda(\mathcal{M})=\sum_{j\in J}\lambda(\mathcal{N})\alpha^-_\lambda(\mathcal{H}_j).$$

Thanks to Remark 2.12, to compute the eigenvalues of $E_PE_QE_P$ it suffices to compute E_PE_Q on $\alpha^+_\lambda(\mathcal{H}_j)$. We claim that for every $t\in\mathcal{H}_j$ the equalities

$$E_Q(\alpha^+_\lambda(t))=\phi_\lambda(\varepsilon(\lambda,\lambda_j)^*\varepsilon(\lambda_j,\lambda)^*)\alpha^-_\lambda(t),$$

$$E_P(\alpha^-_\lambda(t))=\phi_\lambda(\varepsilon(\lambda_j,\lambda)\varepsilon(\lambda,\lambda_j))\alpha^+_\lambda(t),$$

hold. Note that we have $E_Q=\alpha^-_\lambda\phi_{\alpha^-_\lambda}$ and $\phi_{\alpha^-_\lambda}=r^*_\lambda\alpha^-_{\bar\lambda}(\cdot)r_\lambda$. Thus

$$\begin{aligned}\phi_{\alpha^-_\lambda}(\alpha^+_\lambda(t)) &= r^*_\lambda\alpha^-_{\bar\lambda}(\alpha^+_\lambda(t))r_\lambda=r^*_\lambda\alpha^-_{\bar\lambda}(\varepsilon(\lambda,\lambda_j)^*t)r_\lambda=r^*_\lambda\bar\lambda(\varepsilon(\lambda,\lambda_j)^*)\varepsilon(\lambda_j,\lambda)tr_\lambda\\ &= r^*_\lambda\bar\lambda(\varepsilon(\lambda,\lambda_j)^*)\varepsilon(\lambda_j,\lambda)\lambda_j(r_\lambda)t.\end{aligned}$$

Since

$$r^*_\lambda\bar\lambda(\varepsilon(\lambda,\lambda_j)^*)\varepsilon(\lambda_j,\lambda)\lambda_j(r_\lambda)=\frac{1}{d(\lambda_j)}$$



$$=\frac{1}{d(\lambda_j)}$$



$$=\phi_\lambda(\varepsilon(\lambda,\lambda_j)^*\varepsilon(\lambda_j,\lambda)^*),$$

we get the first equation. The second one follows from a similar computation. Using the claim, we get

$$E_PE_Q(\alpha^+_\lambda(t))=|\phi_\lambda(\varepsilon(\lambda_j,\lambda)\varepsilon(\lambda,\lambda_j))|^2\alpha^+_\lambda(t),$$

which proves the statement. □

7.2. **GHJ pairs.** We first recall the construction of the GHJ subfactors, which first appeared in the book of Goodman, de la Harpe and Jones [11]. Our presentation is based on [9].

Let G be one of the Dynkin diagrams of type A, D, or E, with a distinguished vertex $*$ and let $\tilde{A}_n$ be the string algebra $\mathrm{String}^{(n)}G$ (see [9, p.554] for the definition). For $i\geq 1$, we denote by e_i the Jones projection in $\tilde{A}_{i+1}\cap\tilde{A}'_{i-1}$ defined by [9, Definition 11.5]. Then the Jones projections $\{e_k\}_{k=1}^\infty$ satisfy the Temperley-Lieb relations:
(1) $e_ie_{i\pm1}e_i=\tau e_i$
(2) $e_ie_j=e_je_i$ if $|i-j|\geq 2$
(3) $tr(e_iw)=\tau tr(w)$ if w is a word on $e_1,\ldots,e_{i-1}$.

Here $\tau = 1/(4\cos^2\pi/m)$, where m is an integer associated to G called *the Coxeter number*. For each n, the Coxeter number of A_n is $n+1$, the Coxeter number of D_n is $2n-2$, and the Coxeter numbers of E_6, E_7, and E_8 are 12, 18, and 30 respectively.

Let $\tilde{B}_i$ be the subalgebra of $\tilde{A}_i$ generated by 1 and $e_1, ..., e_{i-1}$, for each $i = 2, 3, ...$. Then the towers $\tilde{B}_i \subset \tilde{A}_i$ form commuting squares, and the von Neumann algebra $\tilde{B}$ generated by $\cup_{i=1}^{\infty}\tilde{B}_i$ is a subfactor of the factor $\tilde{A}$ generated by $\cup_{i=1}^{\infty}\tilde{A}_i$ in the GNS representation of the unique trace on $\cup_{i=1}^{\infty}\tilde{A}_i$. This is an irreducible subfactor with finite index, called the *GHJ subfactor* for $(G, *)$. The principal graphs of the GHJ subfactors were computed by Okamoto in [50].

Because of the Temperley-Lieb relations, there is a unitary representation of the braid group inside the algebras $\tilde{B}_i$, sending the usual braid group generators σ_i to $g_i = (t+1)e_i - 1$, where $t = e^{2\pi i/m}$, m again being the Coxeter number of G. Let $v_i = g_1 g_2 \cdots g_{i-1}$ and $w_i = g_1^{-1} g_2^{-1} \cdots g_{i-1}^{-1}$. We set $\tilde{C}_i = v_i \tilde{A}_{i-1} v_i^*$ and $\tilde{D}_i = w_i \tilde{A}_{i-1} w_i^*$. Then the towers $\tilde{C}_i \subset \tilde{A}_i$ and $\tilde{D}_i \subset \tilde{A}_i$ form commuting squares, and the resulting subfactor $\tilde{P} \subset \tilde{M}$ and $\tilde{Q} \subset \tilde{M}$ are irreducible and $[\tilde{M} : \tilde{P}] = [\tilde{M} : \tilde{Q}] = 4\cos^2\pi/k$. For $G = A_n$ the principal graphs of $\tilde{P} \subset \tilde{M}$ and $\tilde{Q} \subset \tilde{M}$ are A_n, for $G = D_{2n+1}$ they are A_{4n-1}, for $G = D_{2n}$ they are D_{2n}, for $G = E_7$ they are A_{17} , and for $G = E_n$ with $n = 6, 8$ they are E_n. The pair $\tilde{P}$ and $\tilde{Q}$ is called the *GHJ pair* [13]. (Though $*$ is assumed to be an endpoint in [13, Definition 6.2.5], we don't pose this condition here.) We set $\tilde{N} = \tilde{P} \cap \tilde{Q}$ and set $\tilde{R}$ to be the subfactor generated by $\{e_i\}_{i=2}^{\infty}$. By construction, we have $\tilde{R} \subset \tilde{N}$ though these two factors do not necessarily coincide in general. For a special case, the angles between $\tilde{P}$ and $\tilde{Q}$ are computed in [13, Theorem 6.3.2].

Now we show that $\begin{array}{ccc} \tilde{P} & \subset & \tilde{M} \\ \cup & & \cup \\ \tilde{N} & \subset & \tilde{Q} \end{array}$ is essentially the same object as $\begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ discussed in the previous subsection for an appropriate $\{\lambda_i\}_{i\in I}$ and $\mathcal{N} \subset \mathcal{M}$. Consequently, the angles between $\tilde{P}$ and $\tilde{Q}$ can be easily computed by Theorem 7.3. Our argument below is inspired by [2, Appendix].

Let $\mathcal{N}$ be the AFD type III$_1$ factor and let $\{\lambda_i\}_{i=0}^{k}$ be a system of irreducible endomorphisms isomorphic to the irreducible sectors for the $SU(2)_k$ WZW model (see [2] for details). Such a system can be obtained either by the loop group construction [55] or by combination of [57] and [14] using the quantum $SU(2)$ at a root of unity. Then the system $\{\lambda_i\}_{i=0}^{k}$ satisfies the assumptions (1)-(5) of the previous subsection. The principal graph of $\lambda_1(\mathcal{N}) \subset \mathcal{N}$ is A_{k+1} and

$$[\lambda_i\lambda_j] = \bigoplus_{0\le 2l\le \min\{|i+j|,k\}-|i-j|} [\lambda_{|i-j|+2l}],$$

$$d(\lambda_i) = \frac{\sin\frac{(i+1)\pi}{k+2}}{\sin\frac{\pi}{k+2}},$$

$$S_{ij} = \sqrt{\frac{2}{k+2}}\sin\Big(\frac{(i+1)(j+1)\pi}{k+2}\Big).$$

Assume that an inclusion of factors $\mathcal{N} \subset \mathcal{M}$ satisfies the assumption (6). We set $i_0 = 1$ and we use the same notation as in the previous subsection. It is known that such an inclusion is completely classified by a pair $(G, *)$ (up to graph automorphism) where G is one of the Coxeter graphs of type A, D, or E [36, Theorem 2.1], so that we can identify $\tilde{A}_n$ with $(\nu\lambda^n, \nu\lambda^n)$ and $\tilde{B}_n$ with $\nu((\lambda^n, \lambda^n))$. Since $\nu\lambda = \alpha_\lambda^\pm \nu$, we have the inclusion relation $\alpha_\lambda^\pm(\tilde{A}_n) \subset (\alpha_\lambda^\pm \nu\lambda^n, \alpha_\lambda^\pm \nu\lambda^n) = (\nu\lambda^{n+1}, \nu\lambda^{n+1}) = \tilde{A}_{n+1}$. Recall that the trace tr on $\tilde{A}_n$ is given by $\phi_{\nu\lambda^n}$.

Lemma 7.4. *Let the notation be as above. Then*

1. *For $x \in \tilde{A}_n$, we have $\alpha_\lambda^\pm(x) = \varepsilon^\pm(\lambda, \lambda^n)^* x \varepsilon^\pm(\lambda, \lambda^n)$. In consequence, the restriction of $\alpha_\lambda^\pm$ to $\tilde{A}_n$, as a map from $\tilde{A}_n$ to $\tilde{A}_{n+1}$, is trace preserving.*
2. *For $x \in \tilde{A}_n$ we have $E_P(x) \subset \alpha_\lambda^+(\tilde{A}_n)$ and $E_Q(x) \subset \alpha_\lambda^-(\tilde{A}_n)$. The restrictions of E_P and E_Q to $\tilde{A}_n$ is trace preserving.*

Proof. (1) Let $x \in \tilde{A}_n$. Then $\bar{\nu}(x) \in (\hat{\gamma}\lambda^n, \hat{\gamma}\lambda^n)$ and

$$\varepsilon(\lambda, \hat{\gamma})\lambda(\bar{\nu}(x))\varepsilon(\lambda, \hat{\gamma})^* =$$



$$= \bar{\nu}(\varepsilon(\lambda, \lambda^n)^* x \varepsilon(\lambda, \lambda^n)),$$

which shows $\alpha_\lambda^+(x) = \varepsilon(\lambda, \lambda^n)^* x \varepsilon(\lambda, \lambda^n)$. The second equality can be shown in the same way.

(2) Since $E_P(x) = \alpha_\lambda^+ \phi_{\alpha_\lambda^+}(x) = \alpha_\lambda^+(r_\lambda^* \alpha_\lambda^+(x) r_\lambda)$, it suffices to show $r_\lambda^* \alpha_\lambda^+(x) r_\lambda \in \tilde{A}_n$ for $x \in \tilde{A}_n$, which can be easily shown by using (1). This also shows

$$\begin{aligned} \mathrm{tr}(E_P(x)) &= \mathrm{tr}(\alpha_\lambda^+ \phi_{\alpha_\lambda^+}(x)) = \mathrm{tr}(\phi_{\alpha_\lambda^+}(x)) = \phi_{\nu\lambda^n}\phi_{\alpha_\lambda^+}(x) = \phi_{\alpha_\lambda^+ \nu\lambda^n}(x) \\ &= \phi_{\nu\lambda^{n+1}}(x) = \mathrm{tr}(x). \end{aligned}$$

□

Since $\varepsilon(\lambda, \lambda^n)^* = \varepsilon(\lambda, \lambda)^* \lambda(\varepsilon(\lambda, \lambda)^*) \cdots \lambda^{n-1}(\varepsilon(\lambda, \lambda)^*)$, we can identify $\alpha_\lambda^+(\tilde{A}_{n-1})$ with $\tilde{C}_n$ and $\alpha_\lambda^-(\tilde{A}_{n-1})$ with $\tilde{D}_n$ (for an appropriate choice of the braiding).

We now show that $\begin{matrix} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{matrix}$ and $\begin{matrix} \tilde{P} & \subset & \tilde{M} \\ \cup & & \cup \\ \tilde{N} & \subset & \tilde{Q} \end{matrix}$ are conjugate after tensor product with the AFD type III_1 factor.

Lemma 7.5. *Let $\mathcal{L}$ be a factor and $\sigma \in \mathrm{Mor}_0(\mathcal{N}, \mathcal{L})$. We set $\mathcal{A}_n = (\sigma\lambda^n, \sigma\lambda^n)$ and $\mathcal{B}_n = \sigma((\lambda^n, \lambda^n))$. Let $\mathcal{A}$ and $\mathcal{B}$ be the II_1 factors generated by $\cup_n \mathcal{A}_n$ and $\cup_n \mathcal{B}_n$ respectively on the GNS Hilbert space of the unique trace on $\cup_n \mathcal{A}_n$. Then $\mathcal{A} \cap \mathcal{B}' = (\sigma, \sigma)$.*

Proof. Since (λ^n, λ^n) is generated by Jones projections, the statement follows from the flatness of the Jones projections [9, Chapter 12]. □

Theorem 7.6. *Let the notation be as above. Then* $\begin{array}{ccc} \tilde{P}\otimes R_\infty & \subset & \tilde{M}\otimes R_\infty \\ \cup & & \cup \\ \tilde{N}\otimes R_\infty & \subset & \tilde{Q}\otimes R_\infty \end{array}$ *and* $\begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$ *are conjugate where* R_∞ *is the AFD type* III_1 *factor.*

Proof. First we show that $R \subset M$ and $\tilde{R} \subset \tilde{M}$ have the same standard invariant. We fix a positive integer n larger than half of the depth of $R \subset M$, which is finite now. Let $\rho = \nu\lambda$. Then $R = \rho(\mathcal{N})$ and

$$\begin{array}{ccccc} (\bar{\rho}\rho)^{n-1}\bar{\rho}((\rho,\rho)) & \subset & (\bar{\rho}\rho)^{n-1}((\bar{\rho}\rho,\bar{\rho}\rho)) & \subset \cdots \subset & ((\bar{\rho}\rho)^n,(\bar{\rho}\rho)^n) \\ \cup & & \cup & & \cup \\ \mathbb{C} & \subset & (\bar{\rho}\rho)^{n-1}((\bar{\rho},\bar{\rho})) & \subset \cdots \subset & ((\bar{\rho}\rho)^{n-1}\bar{\rho},(\bar{\rho}\rho)^{n-1}\bar{\rho}) \end{array}$$

is the standard invariant of $R \subset M$ (or rather $(\bar{\rho}\rho)^n(\mathcal{N}) \subset (\bar{\rho}\rho)^{n-1}\bar{\rho}(\mathcal{M})$). We set

$$\mathcal{A}_{2j,m} = (\bar{\rho}\rho)^{n-j}(((\bar{\rho}\rho)^j\lambda^m,(\bar{\rho}\rho)^j\lambda^m)),$$

$$\mathcal{A}_{2j+1,m} = (\bar{\rho}\rho)^{n-j-1}\bar{\rho}((\rho(\bar{\rho}\rho)^j\lambda^m,\rho(\bar{\rho}\rho)^j\lambda^m)).$$

Note that $\bar{\rho}\rho$ is a direct sum of irreducibles in $\{\lambda_i\}_{i=0}^k$. When m is larger than the depth of $\lambda(\mathcal{N}) \subset \mathcal{N}$, the inclusion graph for $\mathcal{A}_{j,m} \subset \mathcal{A}_{j+1,m}$ is the reflection of that for $\mathcal{A}_{j+1,m} \subset \mathcal{A}_{j+2,m}$ thanks to Frobenius reciprocity. Therefore

$$\mathcal{A}_{j,m} \subset \mathcal{A}_{j+1,m} \subset \mathcal{A}_{j+2,m}$$

is the basic construction.

Let $\mathcal{A}_{j,\infty}$ be the factor generated by $\cup_m\mathcal{A}_{j,m}$ in the GNS representation of the unique trace of $\cup_m\mathcal{A}_{2n,m}$. Then we may identify $\tilde{R} \subset \tilde{M}$ with $\mathcal{A}_{0,\infty} \subset \mathcal{A}_{1,\infty}$ and

$$\mathcal{A}_{0,\infty} \subset \mathcal{A}_{1,\infty} \subset \mathcal{A}_{2,\infty} \subset \cdots \subset \mathcal{A}_{2n,\infty}$$

is the Jones tower. Lemma 7.5 implies

$$\mathcal{A}_{2j,\infty} \cap \mathcal{A}'_{0,\infty} = (\bar{\rho}\rho)^{n-j}(((\bar{\rho}\rho)^j,(\bar{\rho}\rho)^j)),$$

$$\mathcal{A}_{2j+1,\infty} \cap \mathcal{A}'_{0,\infty} = (\bar{\rho}\rho)^{n-j-1}\bar{\rho}((\rho(\bar{\rho}\rho)^j,\rho(\bar{\rho}\rho)^j)),$$

which shows that $R \subset M$ and $\tilde{R} \subset \tilde{M}$ have the same principal graph. On the other hand, we have

$$\mathcal{A}_{2j,\infty} \cap \mathcal{A}'_{1,\infty} \supset (\bar{\rho}\rho)^{n-j}(((\bar{\rho}\rho)^{j-1}\bar{\rho},(\bar{\rho}\rho)^{j-1}\bar{\rho})),$$

$$\mathcal{A}_{2j+1,\infty} \cap \mathcal{A}'_{1,\infty} \supset (\bar{\rho}\rho)^{n-j-1}\bar{\rho}((\rho\bar{\rho})^j,(\rho\bar{\rho})^j)).$$

Since $[M : R] = [\tilde{M} : \tilde{R}]$, the equality holds in the above inclusions and the two subfactors $R \subset M$ and $\tilde{R} \subset \tilde{M}$ have the same standard invariant. Therefore we conclude that the two inclusions $R \subset M$ and $\tilde{R} \otimes R_\infty \subset \tilde{M} \otimes R_\infty$ are conjugate thanks to Popa's result [53] with [19] (see [47] too).

To finish the proof, it suffices to show that $e_P = e_{\tilde{P}}$ and $e_Q = e_{\tilde{Q}}$ hold in the above identification of the standard invariants, which can be done by using Lemma 7.4,(2). □

Remark 7.7. Since

$$|\phi_\lambda(\varepsilon(\lambda_j,\lambda)\varepsilon(\lambda,\lambda_j))| = \frac{|S_{1j}|}{d(\lambda_1)d(\lambda_j)|S_{00}|} = \frac{|\sin\frac{\pi}{k+2}\sin\frac{2(j+1)\pi}{k+2}|}{|\sin\frac{2\pi}{k+2}\sin\frac{(j+1)\pi}{k+2}|} = \frac{|\cos\frac{(j+1)\pi}{k+2}|}{|\cos\frac{\pi}{k+2}|},$$

the operator $\varepsilon(\lambda_j,\lambda)\varepsilon(\lambda,\lambda_j)$ is a scalar if and only if $j=0,k$. Thus we have $J_{i_0} = J\cap\{0,k\}$. It is known that $k\in J$ occurs only if k is even. In this case, we may and do assume $\lambda_k^2 = \mathrm{id}_{\mathcal{N}}$ by choosing a representative λ_k satisfying $\lambda_k\lambda_{k/2} = \lambda_{k/2}$, which implies that there exists a unitary $u_k \in \mathcal{H}_k$ satisfying $u_k^2 = 1$. Therefore we may regard $\mathcal{N} + \mathcal{N}\mathcal{H}_k$ as the crossed product $\mathcal{N} \rtimes_{\lambda_k} \mathbb{Z}/2\mathbb{Z}$. We make this assumption and identification in what follows.

The following corollary answers the question raised in [13, comment after Proposition 6.2.4] in the negative. (The subfactor $\tilde{R}$ is denoted by $TL2$ in [13].)

Corollary 7.8. *Let the notation be as above. Then* $\tilde{R} = \tilde{P}\cap\tilde{Q}$ *if and only if* $k\notin J$. *If* $k \in J$, *the index of* $\tilde{R}\subset\tilde{P}\cap\tilde{Q}$ *is 2.*

Proof. Thanks to Theorem 7.6, we may work on R and N instead of $\tilde{R}$ and $\tilde{N}$, which together with Lemma 7.2 show the first statement. When $k \in J$, Lemma 7.2 shows $N = \alpha_\lambda^{\pm}(\mathcal{N} + \mathcal{N}\mathcal{H}_k)$ and the second statement follows from Remark 7.7. □

The following corollary is a generalization of [13, Theorem 6.3.2].

Corollary 7.9. *Let the notation be as above. Then*

$$\mathrm{Ang}(\tilde{P},\tilde{Q}) = \{\cos^{-1}\frac{|\cos\frac{(j+1)\pi}{k+2}|}{|\cos\frac{\pi}{k+2}|};\ j\in J\}\setminus\{0,\frac{\pi}{2}\}.$$

Example 7.10. Let $\mathcal{N}\subset\mathcal{M}$ be the inclusion of the AFD type III_1 factors associated with the conformal embedding $SU(2)_{10}\subset SO(5)_1$ (see [58, p.381]). Then the corresponding GHJ subfactor is given by E_6 with $*$ equal to the vertex of the minimum value of the Perron-Frobenius eigenvector, and so its principal graph is as in Theorem 5.4,(2) (see [9, p.726],[50]). Since $[\hat{\gamma}] = [\lambda_0]\oplus[\lambda_6]$, we have $\tilde{R} = \tilde{N}$, and so $\tilde{N}\subset\tilde{P}$ is conjugate to the GHJ subfactor, while $\tilde{P}\subset\tilde{M}$ is the E_6 subfactor. Since

$$\dim(\nu\lambda,\nu\lambda) = \dim(\hat{\gamma},\lambda^2) = \dim(\lambda_0\oplus\lambda_6,\lambda_0\oplus\lambda_2) = 1,$$

the inclusion $\tilde{N}\subset\tilde{M}$ is irreducible. The angle between $\tilde{P}$ and $\tilde{Q}$ is computed as $\Theta(\tilde{P},\tilde{Q}) = \cos^{-1}(2-\sqrt{3})$ by using either of [13, Theorem 6.3.2], Theorem 4.3, or Corollary 7.9.

It is observed in [13, Proposition 6.2.6] that $\tilde{R}\subset\tilde{M}$ is irreducible if and only if $*$ is an endpoint. However, even when $*$ is not an endpoint, the inclusion $\tilde{N}\subset\tilde{M}$ may be irreducible (though it is very subtle in general to decide whether $\tilde{N}\subset\tilde{M}$ is irreducible or not as we will see below). We use this observation to compute the opposite angles of quadrilaterals constructed in [12]. The following theorem is shown in [12].

Theorem 7.11. *Let* $\begin{array}{ccc} B & \subset & A \\ \cup & & \cup \\ D & \subset & C \end{array}$ *be an irreducible noncommuting quadrilateral of the hyperfinite* II_1 *factors such that* $D \subset B$ *and* $D \subset C$ *are supertransitive and* $[B : D]$ *and* $[C : D]$ *are less than 4. Then the principal graphs of* $D \subset B$ *and* $D \subset C$ *are* A_{2n+1} *with an integer* $n \geq 2$*. For each integer* $n \geq 2$*, such a quadrilateral exists and is unique up to conjugacy.*

We now compute the "opposite angles" of the above quadrilateral.

Theorem 7.12. *Let* $\begin{array}{ccc} \hat{C} & \subset & \hat{D} \\ \cup & & \cup \\ A & \subset & \hat{B} \end{array}$ *be the dual quadrilateral of the quadrilateral in the previous theorem. Then*

$$\mathrm{Ang}(\hat{B}, \hat{C}) = \{\cos^{-1} \frac{\cos \frac{(2j+1)\pi}{2n+2}}{\cos \frac{\pi}{2n+2}};\ j = 1, 2, \cdots, [\frac{n-1}{2}]\},$$

where $[(n-1)/2]$ *is the integer part of* $(n-1)/2$*.*

Proof. We show that the above quadrilateral is given by the GHJ pair of A_{2n+1} with $*$ equal to the midpoint. Thanks to Theorem 7.6, we can use α-induction instead of the GHJ construction for this purpose. Let $\mathcal{N}$ and $\{\lambda_i\}_{i=0}^k$ be as before with $k = 2n$. Let $(\mathcal{M}, \{\tilde{\lambda}_i\}_{i=0}^k)$ be a copy of $(\mathcal{N}, \{\lambda_i\}_{i=0}^k)$. We embed $\mathcal{N}$ into $\mathcal{M}$ by

$$\nu : \mathcal{N} \ni x \mapsto \lambda_n(x) \in \mathcal{M}.$$

Then we have $\bar{\nu} = \tilde{\lambda}_n$ regarded as a map from $\mathcal{M}$ into $\mathcal{N}$, and so $\hat{\gamma} = \lambda_n^2$ and $\gamma = \tilde{\lambda}_n^2$. Since

$$[\lambda_n^2] = \bigoplus_{i=0}^{n} [\lambda_{2i}],$$

we have $J = \{0, 2, 4, \cdots, k\}$ and $J_{i_0} = \{0, k\}$. Therefore thanks to Corollary 7.8,

$$N = \alpha_\lambda^\pm(\mathcal{N} \rtimes_{\lambda_k} \mathbb{Z}/2\mathbb{Z}) = \alpha_\lambda^\pm(\nu(\mathcal{N}) + \nu(\mathcal{N})u_k),$$

where $u_k \in (\nu, \nu\lambda_k) = (\tilde{\lambda}_n, \tilde{\lambda}_n\tilde{\lambda}_k)$. As before, we set $M = \mathcal{M}$, $P = \alpha_\lambda^+(\mathcal{M})$, $Q = \alpha_\lambda^-(\mathcal{M})$, $N = P \cap Q$, and $\mathfrak{Q} = \begin{array}{ccc} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{array}$. Then the principal graphs of $P \subset M$ and $Q \subset M$ are A_{2n+1}. To prove the theorem, it suffices to show that $N \subset M$ is irreducible and $\mathfrak{Q}$ is not cocommuting thanks to Corollary 7.9 and Theorem 7.11.

We claim $\alpha_\lambda^\pm = \mathrm{Ad}\varepsilon^\pm(\lambda, \lambda_n)\tilde{\lambda}$. Indeed, since $\varepsilon^\pm(\lambda, \hat{\gamma}) = \lambda_n(\varepsilon^\pm(\lambda, \lambda_n))\varepsilon^\pm(\lambda, \lambda_n)$ and $\varepsilon^\pm(\lambda, \lambda_n) \in (\lambda\lambda_n, \lambda_n\lambda) = (\lambda\bar{\nu}, \bar{\nu}\tilde{\lambda})$, we have

$$\mathrm{Ad}(\varepsilon^\pm(\lambda, \hat{\gamma}))\lambda\bar{\nu}(x) = \mathrm{Ad}\lambda_n(\varepsilon^\pm(\lambda, \lambda_n))\mathrm{Ad}\varepsilon^\pm(\lambda, \lambda_n)\lambda\bar{\nu}(x) = \mathrm{Ad}\bar{\nu}(\varepsilon^\pm(\lambda, \lambda_n))\bar{\nu}(\tilde{\lambda}(x))$$

for all $x \in \mathcal{M}$, which shows the claim. In particular, the two subfactors P and Q are inner conjugate in M. Therefore if $N \subset M$ is irreducible, Theorem 3.10 applied to the dual quadrilateral $\hat{\mathfrak{Q}}$ shows that $\mathfrak{Q}$ is not cocommuting.

Now our only task is to show $M \cap N' = \mathbb{C}$. Note that N is generated by $\nu(\lambda(\mathcal{N}))$ and $\nu(\varepsilon(\lambda,\lambda_k)^*)u_k$ with $u_k \in (\nu, \nu\lambda_k)$, and so

$$M \cap N' = (\nu\lambda, \nu\lambda) \cap \{\nu(\varepsilon(\lambda,\lambda_k)^*)u_k\}'.$$

Therefore it suffices to show

$$(\lambda_n\lambda, \lambda_n\lambda) \cap \{\lambda_n(\varepsilon(\lambda,\lambda_k)^*)u\}' = \mathbb{C}.$$

where u is a unitary in $(\lambda_n, \lambda_n\lambda_k)$. By Frobenius reciprocity, we have

$$\dim(\lambda_n\lambda, \lambda_n\lambda) = \dim(\lambda_n^2, \lambda^2) = \dim(\lambda_n^2, \lambda_0 \oplus \lambda_2) = 2,$$

and we can choose a basis $\{1, s\}$ of $(\lambda_n\lambda, \lambda_n\lambda)$ with

$s =$ 

,

where $v \in (\lambda_n, \lambda_n\lambda_2)$ and $w \in (\lambda_2\lambda, \lambda)$. On one hand, we have

$s\lambda_n(\varepsilon(\lambda,\lambda_k)^*)u =$ 

$=$ 

,

where $\tilde{v} = \varepsilon(\lambda_2, \lambda_n)^* v$. On the other hand,

$\lambda_n(\varepsilon(\lambda,\lambda_k)^*)us =$ 

$=$ 

.

Since $\dim(\lambda_n, \lambda_2\lambda_n\lambda_k) = 1$, there exists a scalar c such that



$= c$ 

,

which implies $s\lambda_n(\varepsilon(\lambda,\lambda_k)^*)u = c\lambda_n(\varepsilon(\lambda,\lambda_k)^*)us$. The number c is given by Kirillov-Reshetikhin's quantum $6j$-symbol with $q = e^{\pi i/(n+1)}$ (see [37, (5.4)]). Using the notation in [37] (irreducible representations are parameterized by half integers there), we get

$$c = \left\{ \begin{array}{ccc} n & \frac{n}{2} & \frac{n}{2} \\ 1 & \frac{n}{2} & \frac{n}{2} \end{array} \right\}_q .$$

Note that although an explicit formula of the $6j$-symbols depends on the choice of intertwiners in general, the above one does not because the same intertwiners u and $\tilde{v}$ appear in the both sides. This can be computed by [37, Theorem 5.1] and it turns out that we have $c = -1$. Therefore $N \subset M$ is irreducible. □

7.3. **Asymptotic inclusions and angles.** Another rich source of noncommuting quadrilaterals is asymptotic inclusions. We show that a technique similar to the one we used in the previous subsection works in this case too.

Let $\mathcal{N} \subset \mathcal{M}$ be an irreducible inclusion of the hyperfinite II_1 factors with finite index and of finite depth and let

$$\mathcal{N} \subset \mathcal{M} \subset \mathcal{M}_1 \subset \cdots \subset \mathcal{M}_n \subset \cdots$$

be the Jones tower. The algebra $\cup_n \mathcal{M}_n$ has a unique trace and we denote by $\mathcal{M}_\infty$ the factor generated by $\cup_n \mathcal{M}_n$ in the GNS representation of the unique trace of $\cup_n \mathcal{M}_n$. Then the inclusion $\mathcal{M} \vee (\mathcal{M}_\infty \cap \mathcal{M}') \subset \mathcal{M}_\infty$ is called the asymptotic inclusion of $\mathcal{N} \subset \mathcal{M}$ (see [9, Chapter 12.6] for details). It is easy to show that

$$\begin{array}{ccc} \mathcal{N} \vee (\mathcal{M}_\infty \cap \mathcal{N}') & \subset & \mathcal{M}_\infty \\ \cup & & \cup \\ \mathcal{N} \vee (\mathcal{M}_\infty \cap \mathcal{M}') & \subset & \mathcal{M} \vee (\mathcal{M}_\infty \cap \mathcal{M}') \end{array}$$

is an irreducible commuting quadrilateral of factors. We show how to compute the “opposite angles" of this quadrilateral.

Example 7.13. When the principal graph of $\mathcal{N} \subset \mathcal{M}$ is A_n, there exists a sequence of Jones projection $\{e_i\}_{i=1}^\infty$ such that $\mathcal{N} = \{e_i\}_{i=2}^{\infty}{}''$ $\subset \mathcal{M} = \{e_i\}_{i=1}^{\infty}{}''$. We set $e_0 = e_N$, $e_{-1} = e_{\mathcal{M}}$ and $e_{-(i+1)} = e_{\mathcal{M}_i}$ for $i \geq 1$. Then $\mathcal{M}_\infty = \{e_i\}_{i=-\infty}^{\infty}{}''$, $\mathcal{M} \vee (\mathcal{M}_\infty \cap \mathcal{M}') = \{e_i\}''_{i\neq 0}$, $\mathcal{N} \vee (\mathcal{M}_\infty \cap \mathcal{N}') = \{e_i\}''_{i\neq 1}$, and $\mathcal{N} \vee (\mathcal{M}_\infty \cap \mathcal{M}') = \{e_i\}''_{i\neq 0,1}$. The index of the asymptotic inclusion in this case is computed in [8] and the principal graphs are obtained in [10].

It is well-known that $\mathcal{M}_\infty \cap \mathcal{M}'$ is naturally identified with the opposite algebra $\mathcal{M}^{\mathrm{opp}}$ of $\mathcal{M}$ and $\mathcal{M} \vee (\mathcal{M}_\infty \cap \mathcal{M}')$ is naturally identified with $\mathcal{M} \otimes \mathcal{M}^{\mathrm{opp}}$, and so we identify $\mathcal{N} \vee (\mathcal{M}_\infty \cap \mathcal{M}')$ with $\mathcal{N} \otimes \mathcal{M}^{\mathrm{opp}}$ too. However, we never identify $\mathcal{N} \vee (\mathcal{M}_\infty \cap \mathcal{N}')$ with $\mathcal{N} \otimes \mathcal{N}^{\mathrm{opp}}$ as we cannot take common factorizations $\mathcal{M} \otimes \mathcal{M}^{\mathrm{opp}}$ and $\mathcal{N} \otimes \mathcal{N}^{\mathrm{opp}}$ for $\mathcal{M} \vee (\mathcal{M}_\infty \cap \mathcal{M}')$ and $\mathcal{N} \vee (\mathcal{M}_\infty \cap \mathcal{N}')$. By cutting the Hilbert space $L^2(\mathcal{M}_\infty)$ by an appropriate projection in $\mathcal{M}'_\infty$, we may assume that $\mathcal{M}_\infty$ acts on $L^2(\mathcal{M} \otimes \mathcal{M}^{\mathrm{opp}}) = L^2(\mathcal{M}) \otimes L^2(\mathcal{M}^{\mathrm{opp}})$. Since e_N may be confused with $e_N \otimes 1$ in this representation, we use the symbol e for e_N in $\mathcal{M}_\infty$. Let $J = J_{\mathcal{M}} \otimes J_{\mathcal{M}^{\mathrm{opp}}}$ be the canonical conjugation of $\mathcal{M} \otimes \mathcal{M}^{\mathrm{opp}}$. We set

$$M := J\big(\mathcal{N} \vee (\mathcal{M}_\infty \cap \mathcal{M}')\big)'J = J(\mathcal{N} \otimes \mathcal{M}^{\mathrm{opp}})'J = \mathcal{M}_1 \otimes \mathcal{M}^{\mathrm{opp}},$$

$$P := J\big(\mathcal{M} \vee (\mathcal{M}_\infty \cap \mathcal{M}')\big)'J = J(\mathcal{M} \otimes \mathcal{M}^{\mathrm{opp}})'J = \mathcal{M} \otimes \mathcal{M}^{\mathrm{opp}},$$

$$Q := J\big(\mathcal{N} \vee (\mathcal{M}_\infty \cap \mathcal{N}')\big)' J, \quad N := J\mathcal{M}'_\infty J.$$

Note that Q is not equal to $\mathcal{M}_1 \otimes \mathcal{N}^{\mathrm{opp}}$ as we formally (just formally) broke the symmetry between $\mathcal{M} \vee (\mathcal{M}_\infty \cap \mathcal{M}')$ and $\mathcal{N} \vee (\mathcal{M}_\infty \cap \mathcal{N}')$ by representing $\mathcal{M}_\infty$ on $L^2(\mathcal{M} \otimes \mathcal{M}^{\mathrm{opp}})$. However, we still have the following:

Lemma 7.14. *Let the notation be as above. Then Q and $\mathcal{M}_1 \otimes \mathcal{N}^{\mathrm{opp}}$ are inner conjugate in M. In particular, we have $M \otimes_Q M \cong \mathcal{M}_1 \otimes \mathcal{M}_1^{\mathrm{opp}}$ as $M-M$ bimodules.*

Proof. Note that

$$\mathcal{N} \otimes \mathcal{N}^{\mathrm{opp}} \subset \mathcal{N} \otimes \mathcal{M}^{\mathrm{opp}} \subset \mathcal{N} \vee (\mathcal{M}_\infty \cap \mathcal{N}') = (\mathcal{N} \otimes \mathcal{M}^{\mathrm{opp}}) \vee \{e\}$$

is the basic construction with the Jones projection e. Thus

$$Q \subset \mathcal{M}_1 \otimes \mathcal{M}^{\mathrm{opp}} \subset J(\mathcal{N} \otimes \mathcal{N}^{\mathrm{opp}})' J = \mathcal{M}_1 \otimes \mathcal{M}_1^{\mathrm{opp}}$$

is the basic construction too, which shows that Q and $\mathcal{M}_1 \otimes \mathcal{N}^{\mathrm{opp}}$ are inner conjugate in M. □

Our purpose of this subsection is to show the following theorem:

Theorem 7.15. *Let the notation be as above. If the principal graph of $\mathcal{N} \subset \mathcal{M}$ is A_n, then*

$$\mathrm{Ang}(P, Q) = \{\cos^{-1} \frac{\cos \frac{(j+1)\pi}{n+1}}{\cos \frac{\pi}{n+1}};\ j = 1, 2, \cdots [\frac{n-2}{2}]\}.$$

We go back to the general (not necessarily A_n) case. Let $\mathfrak{Q} = \begin{matrix} P & \subset & M \\ \cup & & \cup \\ N & \subset & Q \end{matrix}$ be as above. To compute $\mathrm{Ang}(P, Q)$ we may replace M with

$$M \otimes \mathbb{B}(\ell^2) \otimes \mathbb{B}(\ell^2) \cong \mathcal{M} \otimes \mathbb{B}(\ell^2) \otimes (\mathcal{M} \otimes \mathbb{B}(\ell^2))^{\mathrm{opp}},$$

and so we may and do assume that $\mathcal{N} \subset \mathcal{M}$ is an inclusion of properly infinite factors from now on. In this case, the inclusion $N \subset P$ is described as the Longo-Rehren inclusion [46] whose structure is well-studied [22]. Therefore the structure of $N \subset P \subset M$ is completely understood. It is a little tricky to decide the position of Q, but we can handle it by using Lemma 7.14. We use the symbols ι, ι_Q, κ, and κ_Q as in Section 4.

We recall the notation in [22] to describe the Longo-Rehren inclusion $N \subset P$. Let $\nu : \mathcal{M} \hookrightarrow \mathcal{M}_1$ be the inclusion map and let $\Delta = \{\rho_\xi\}_{\xi \in \Delta_0}$ be the set of irreducibles contained in $\cup_n (\bar{\nu}\nu)^n$. We may assume that $\rho_e = \mathrm{id}_{\mathcal{M}}$ with $e \in \Delta_0$ and $[\overline{\rho_\xi}] = [\rho_{\bar{\xi}}]$. We identify $\mathcal{M}^{\mathrm{opp}}$ with $\mathcal{M}' = J_{\mathcal{M}} \mathcal{M} J_{\mathcal{M}}$ and define $j : \mathcal{M} \to \mathcal{M}^{\mathrm{opp}}$ by $j(x) = J_{\mathcal{M}} x J_{\mathcal{M}}$ for $x \in \mathcal{M}$. For $x \in \mathcal{M}$, the corresponding element $x^{\mathrm{opp}} \in \mathcal{M}^{\mathrm{opp}}$ is identified with $j(x^*)$. We denote by ρ_ξ^{opp} the endomorphism of $\mathcal{M}^{\mathrm{opp}}$ corresponding to ρ_ξ. In our situation, we have $\rho_\xi^{\mathrm{opp}} = j \rho_\xi j^{-1}$. It is known that we have $[\iota\bar{\iota}] = \bigoplus_{\xi \in \Delta_0} [\hat{\rho}_\xi]$, where $\hat{\rho}_\xi = \rho_\xi \otimes \rho_\xi^{\mathrm{opp}}$, and so $d(\iota)^2 = \sum_{\xi \in \Delta_0} d(\xi)^2$. We choose an isometry $V_\xi \in (\rho_\xi \otimes \rho_\xi^{\mathrm{opp}}, \iota\bar{\iota})$ such that $r_{\bar{\iota}} = V_e$, which will be denoted by V for simplicity. We denote $r_\iota \in N$ by W, which is described as follows. Let $\{T(\substack{\zeta\\ \xi,\eta})_i\}_{i=1}^{N_{\xi,\eta}^{\zeta}}$ be an orthonormal basis of $(\rho_\zeta, \rho_\xi\rho_\eta)$, and set

$$T_{\xi,\eta}^{\zeta} = \sum_{i=1}^{N_{\xi,\eta}^{\zeta}} T(\substack{\zeta\\ \xi,\eta})_i \otimes j(T(\substack{\zeta\\ \xi,\eta})_i),$$

which does not depend on the choice of the basis. Then we have

$$W = \sum_{\xi,\eta,\zeta} \frac{\sqrt{d(\xi)d(\eta)}}{d(\iota)\sqrt{d(\zeta)}} V_\xi \hat{\rho}_\xi(V_\eta) T_{\xi,\eta}^{\zeta} V_\zeta^*.$$

To describe the conditional expectation E_Q, we use the following lemma:

Lemma 7.16. *Let $\mathcal{B} \subset \mathcal{A}$ be an irreducible inclusion of property infinite factors with finite index and let $\mu : \mathcal{B} \hookrightarrow \mathcal{A}$ be the inclusion map. Let $[\mu\bar{\mu}] = \bigoplus_{i=0}^{m} n_i[\sigma_i]$ be the irreducible decomposition and let $\{s(i)_l\}_{l=1}^{n_i}$ be an orthonormal basis of $(\mu, \sigma_i\mu)$. Then*

$$E_{\mathcal{B}}(x) = \frac{1}{[\mathcal{A} : \mathcal{B}]} \sum_{i=0}^{m} \sum_{l=1}^{n_i} d(\sigma_i) s(i)_l^* \sigma_i(x) s(i)_l, \quad \forall x \in \mathcal{A}.$$

Proof. Let $\tilde{s}(i)_l = \sqrt{d(\sigma_i)} s(i)_l^* \sigma_i(r_{\bar{\mu}})$. Then $\{\tilde{s}(i)_l\}_{l=1}^{n_i}$ is an orthonormal basis of $(\sigma_i, \mu\bar{\mu})$ thanks to Frobenius reciprocity. Thus

$$E_{\mathcal{B}}(x) = \mu(r_\mu^* \bar{\mu}(x) r_\mu) = \mu(r_\mu^*)\mu\bar{\mu}(x)\mu(r_\mu) = \sum_{i=0}^{m}\sum_{l=1}^{n_i} \mu(r_\mu^*)\tilde{s}(i)_l \sigma_i(x) \tilde{s}(i)_l^* \mu(r_\mu).$$

On the other hand,

$$\mu(r_\mu^*)\tilde{s}(i)_l = \sqrt{d(\sigma_i)}\mu(r_\mu^*) s(i)_l^* \sigma_i(r_{\bar{\mu}}) = \sqrt{d(\sigma_i)} s(i)_l^* \sigma_i(\mu(r_\mu^*) r_{\bar{\mu}}) = \frac{\sqrt{d(\sigma_i)}}{d(\mu)} s(i)_l^*,$$

which shows the statement. □

Recall that to compute $\mathrm{Ang}(P, Q)$, it suffices to determine the eigenvalues of $E_P E_Q$ restricted to P.

Lemma 7.17. *Let $[\bar{\nu}\nu] = \bigoplus_{\xi\in\Delta_1} n_\xi[\rho_\xi]$ be the irreducible decomposition. Then for $x \in P$, we have*

$$E_P E_Q(x) = \frac{d(\iota)^2}{[\mathcal{M} : \mathcal{N}]} \sum_{\xi\in\Delta_1} \frac{n_\xi}{d(\xi)} W^* V_\xi \hat{\rho}_\xi(x) V_\xi^* W.$$

Proof. Thanks to Lemma 7.14, we have $[\kappa_Q\bar{\kappa}_Q] = \bigoplus_{\xi\in\Delta_1} n_\xi[\mathrm{id}_{\mathcal{M}_1} \otimes \rho_\xi^{\mathrm{opp}}]$. Thus to compute E_Q using Lemma 7.16, it suffices to obtain $(\kappa_Q, (\mathrm{id}_{\mathcal{M}_1\otimes\rho_\xi^{\mathrm{opp}}})\kappa_Q)$ for $\xi \in \Delta_1$. Although it is not so easy to capture this space directly, we have

$$\begin{aligned}
(\kappa_Q, (\mathrm{id}_{\mathcal{M}_1} \otimes \rho_\xi^{\mathrm{opp}})\kappa_Q) &\subset (\kappa_Q\iota_Q, (\mathrm{id}_{\mathcal{M}_1} \otimes \rho_\xi^{\mathrm{opp}})\kappa_Q\iota_Q) \\
&= ((\nu \otimes \mathrm{id}_{\mathcal{M}^{\mathrm{opp}}})\iota, (\nu \otimes \rho_\xi^{\mathrm{opp}})\iota) \\
&\cong (\iota\bar{\iota}, \bar{\nu}\nu \otimes \rho_\xi^{\mathrm{opp}}).
\end{aligned}$$

Comparing the dimensions of the both sides above, we get equality for the above inclusion.

Let $\{t(\xi)_i\}_{i=1}^{n_\xi}$ be an orthonormal basis of $(\nu,\nu\rho_\xi)$. We claim that

$$\{\frac{d(\iota)}{\sqrt{d(\xi)}}(t(\xi)_i^*\otimes 1)V_\xi^*W\}_{i=1}^{n_\xi}$$

is an orthonormal basis of $((\nu\otimes\mathrm{id}_{\mathcal{M}^{\mathrm{opp}}})\iota,(\nu\otimes\rho_\xi^{\mathrm{opp}})\iota)$. Indeed, it is easy to show that $(t(\xi)_i^*\otimes 1)V_\xi^*W$ belongs to the intertwiner space $((\nu\otimes\mathrm{id}_{\mathcal{M}^{\mathrm{opp}}})\iota,(\nu\otimes\rho_\xi^{\mathrm{opp}})\iota)$, and so the operator $W^*V_\xi(t(\xi)_l t(\xi)_i^*\otimes 1)V_\xi^*W$ is already a scalar, which is equal to

$$\begin{aligned}&E_NE_P(W^*V_\xi(t(\xi)_l t(\xi)_i^*\otimes 1)V_\xi^*W)=E_N(W^*V_\xi E_P(t(\xi)_l t(\xi)_i^*\otimes 1)V_\xi^*W)\\ &= \frac{\langle t(\xi)_l,t(\xi)_i\rangle}{d(\xi)}E_N(W^*V_\xi V_\xi^*W)=\frac{\delta_{i,l}}{d(\xi)}W^*E_N(V_\xi V_\xi^*)W=\frac{d(\xi)\delta_{i,l}}{d(\iota)^2}W^*W\\ &= \frac{d(\xi)\delta_{i,l}}{d(\iota)^2}.\end{aligned}$$

This shows the claim. Thus Lemma 7.16 shows that for $x\in P$ we have

$$\begin{aligned}E_Q(x) &= \frac{1}{[\mathcal{M}:\mathcal{N}]}\sum_{\xi\in\Delta_1}\sum_{i=1}^{n_\xi}d(\iota)^2W^*V_\xi(t(\xi)_i\otimes 1)(\mathrm{id}_{\mathcal{M}}\otimes\rho_\xi^{\mathrm{opp}})(x)(t(\xi)_i^*\otimes 1)V_\xi^*W\\ &= \frac{d(\iota)^2}{[\mathcal{M}:\mathcal{N}]}\sum_{\xi\in\Delta_1}\sum_{i=1}^{n_\xi}W^*V_\xi\hat{\rho}_\xi(x)(t(\xi)_i t(\xi)_i^*\otimes 1)V_\xi^*W.\end{aligned}$$

Therefore we get

$$\begin{aligned}E_PE_Q(x) &= \frac{d(\iota)^2}{[\mathcal{M}:\mathcal{N}]}\sum_{\xi\in\Delta_1}\sum_{i=1}^{n_\xi}W^*V_\xi\hat{\rho}_\xi(x)E_P(t(\xi)_i t(\xi)_i^*\otimes 1)V_\xi^*W\\ &= \frac{d(\iota)^2}{[\mathcal{M}:\mathcal{N}]}\sum_{\xi\in\Delta_1}\frac{n_\xi}{d(\xi)}W^*V_\xi\hat{\rho}_\xi(x)V_\xi^*W.\end{aligned}$$

□

To compute the eigenvalues of $E_PE_Q|_P$, we need the crossed product-like decomposition as in Theorem 2.9 based on the irreducible decomposition of $\bar{\iota}\iota$. It is known that every sector contained in a power of $\bar{\iota}\iota$ is of the form $\widetilde{\sigma}^\alpha$, where σ is a finite direct sum of endomorphisms in Δ and α is a parameter of the corresponding half-braiding $\varepsilon_\sigma^\alpha(\xi)\in(\sigma\rho_\xi,\rho_\xi\sigma)$ (see [22, Section 4] for details). Let

$$U^\alpha(\sigma)=\sum_{\xi\in\Delta_0}V_\xi(\varepsilon_\sigma^\alpha(\xi)\otimes 1)(\sigma\otimes\mathrm{id}_{\mathcal{M}^{\mathrm{opp}}})(V_\xi^*).$$

Then $\widetilde{\sigma}^\alpha$ is the restriction of $\mathrm{Ad}U^\alpha(\sigma)(\sigma\otimes\mathrm{id}_{\mathcal{M}^{\mathrm{opp}}})$ to N. Irreducible $\widetilde{\sigma}^\alpha$ is contained in $\bar{\iota}\iota$ if and only if σ contains $\mathrm{id}_{\mathcal{M}}$. In this case, we have

$$(\iota,\iota\widetilde{\sigma}^\alpha)=U^\alpha(\sigma)\big((\mathrm{id}_{\mathcal{M}},\sigma)\otimes\mathbb{C}\big).$$

Therefore to compute $\mathrm{Ang}(P,Q)$, it suffices to compute E_PE_Q on this space thanks to Remark 2.11.

Lemma 7.18. *Let $\widetilde{\sigma}^\alpha$ be an irreducible endomorphism of N contained in $\bar{\iota}\iota$. Then for $v \in (\mathrm{id}_{\mathcal{M}}, \sigma)$,*

$$E_P E_Q(U^\alpha(\sigma)(v \otimes 1)) = U^\alpha(\sigma)(\phi_{\bar{\nu}\nu}(\varepsilon_\sigma^\alpha(\bar{\nu}\nu)v) \otimes 1).$$

The map $(\mathrm{id}_{\mathcal{M}}, \sigma) \ni v \mapsto \phi_{\bar{\nu}\nu}(\varepsilon_\sigma^\alpha(\bar{\nu}\nu)v)$ depends only on the sector of $\bar{\nu}\nu$ and

$$\phi_{\bar{\nu}\nu}(\varepsilon_\sigma^\alpha(\bar{\nu}\nu)v) = \sum_{\xi \in \Delta_1} \frac{n_\xi d(\xi)}{[\mathcal{M} : \mathcal{N}]} \phi_{\rho_\xi}(\varepsilon_\sigma^\alpha(\xi)v).$$

Proof. It is routine work to show the second statement. Since we have

$$U^\alpha(\sigma)(v \otimes 1) = \sum_{\eta \in \Delta_0} V_\eta(\varepsilon_\sigma^\alpha(\eta)v \otimes 1)V_\eta^*,$$

Lemma 7.17 implies

$$\begin{aligned}
&E_P E_Q(U^\alpha(\sigma)(v \otimes 1)) \\
&= \frac{d(\iota)^2}{[\mathcal{M} : \mathcal{N}]} \sum_{\xi \in \Delta_1} \sum_{\eta \in \Delta_0} \frac{n_\xi}{d(\xi)} W^* V_\xi \hat{\rho}_\xi(V_\eta(\varepsilon_\sigma^\alpha(\eta)v \otimes 1)V_\eta^*)V_\xi^* W \\
&= \frac{1}{[\mathcal{M} : \mathcal{N}]} \sum_{\eta, \zeta \in \Delta_0} \sum_{\xi \in \Delta_1} \sum_{i=1}^{N_{\xi,\eta}^\zeta} \frac{n_\xi d(\eta)}{d(\zeta)} V_\zeta \big(T\tbinom{\zeta}{\xi,\eta}_i^* \rho_\xi(\varepsilon_\sigma^\alpha(\eta)v)T\tbinom{\zeta}{\xi,\eta}_i \otimes 1\big)V_\zeta^*.
\end{aligned}$$

Thanks to the half-braiding relation, we have

$$T\tbinom{\zeta}{\xi,\eta}_i^* \rho_\xi(\varepsilon_\sigma^\alpha(\eta)v) = \varepsilon_\sigma^\alpha(\zeta)\sigma(T\tbinom{\zeta}{\xi,\eta})_i^*)\varepsilon_\sigma^\alpha(\xi)^* \rho_\xi(v).$$

Thus Frobenius reciprocity implies

$$\begin{aligned}
&\sum_{\eta \in \Delta_0} \sum_{i=1}^{N_{\xi,\eta}^\zeta} d(\eta) T\tbinom{\zeta}{\xi,\eta}_i^* \rho_\xi(\varepsilon_\sigma^\alpha(\eta)v) T\tbinom{\zeta}{\xi,\eta}_i \\
&= \varepsilon_\sigma^\alpha(\zeta) \sum_{\eta \in \Delta_0} \sum_{i=1}^{N_{\xi,\eta}^\zeta} d(\eta)\sigma(T\tbinom{\zeta}{\xi,\eta}))_i^*)\varepsilon_\sigma^\alpha(\xi)^* \rho_\xi(v) T\tbinom{\zeta}{\xi,\eta}_i \\
&= d(\xi)d(\zeta)\varepsilon_\sigma^\alpha(\zeta) \sum_{\eta \in \Delta_0} \sum_{i=1}^{N_{\xi,\eta}^\zeta} \sigma(r_{\rho_{\bar{\xi}}}^* \rho_\xi(T\tbinom{\eta}{\bar{\xi},\zeta}))_i))\varepsilon_\sigma^\alpha(\xi)^* \rho_\xi(vT\tbinom{\eta}{\bar{\xi},\zeta}_i^*)r_{\rho_{\bar{\xi}}} \\
&= d(\xi)d(\zeta)\varepsilon_\sigma^\alpha(\zeta) \sum_{\eta \in \Delta_0} \sum_{i=1}^{N_{\xi,\eta}^\zeta} \sigma(r_{\rho_{\bar{\xi}}}^* \rho_\xi(T\tbinom{\eta}{\bar{\xi},\zeta}))_i))\varepsilon_\sigma^\alpha(\xi)^* \rho_\xi(\sigma(T\tbinom{\eta}{\bar{\xi},\zeta}_i^*)v)r_{\rho_{\bar{\xi}}} \\
&= d(\xi)d(\zeta)\varepsilon_\sigma^\alpha(\zeta) \sum_{\eta \in \Delta_0} \sum_{i=1}^{N_{\xi,\eta}^\zeta} \sigma(r_{\rho_{\bar{\xi}}}^* \rho_\xi(T\tbinom{\eta}{\bar{\xi},\zeta}_i T\tbinom{\eta}{\bar{\xi},\zeta}_i^*))\varepsilon_\sigma^\alpha(\xi)^* \rho_\xi(v)r_{\rho_{\bar{\xi}}} \\
&= d(\xi)d(\zeta)\varepsilon_\sigma^\alpha(\zeta)\sigma(r_{\rho_{\bar{\xi}}}^*)\varepsilon_\sigma^\alpha(\xi)^* \rho_\xi(v)r_{\rho_{\bar{\xi}}}.
\end{aligned}$$

Using the half-braiding relation again, we see that this is equal to

$$d(\xi)d(\zeta)\varepsilon^{\alpha}_{\sigma}(\zeta)r^*_{\rho_{\bar{\xi}}}\rho_{\xi}(\varepsilon^{\alpha}_{\sigma}(\bar{\xi})v)r_{\rho_{\bar{\xi}}}=d(\xi)d(\zeta)\varepsilon^{\alpha}_{\sigma}(\zeta)\phi_{\rho_{\bar{\xi}}}(\varepsilon^{\alpha}_{\sigma}(\bar{\xi})v),$$

which finishes the proof. □

Proof of Theorem 7.15. Assume that the principal graph of $\mathcal{N}\subset\mathcal{M}$ is A_n. The structure of $N-N$ sectors associated with the inclusion $N\subset P$ is described in [22, Section 7]. We may assume that there exists a system of endomorphism $\{\lambda_i\}_{i=0}^{n-1}$ of $\mathcal{M}$ isomorphic to the irreducible sectors of the $SU(2)_{n-1}$ WZW model such that $\mathcal{N}=\lambda_1(\mathcal{M})$. Then we have $[\bar{\nu}\nu]=[\lambda_1^2]$ and $\Delta=\{\lambda_{2j}\}_{j=0}^{[(n-1)/2]}$. When $i-j$ is even, we set $\sigma_{i,j}=\lambda_i\lambda_j$ and $\varepsilon_{\sigma_{i,j}}(\lambda_l)=\varepsilon^+(\lambda_i,\lambda_l)\lambda_i(\varepsilon^-(\lambda_j,\lambda_l))$. Then $\{\varepsilon_{\sigma_{i,j}}(\lambda_l)\}_l$ is a half-braiding for $\sigma_{i,j}$ and we denote by $\widetilde{\sigma_{i,j}}$ the corresponding endomorphism of N.

Assume first that n is even. In this case, the endomorphism $\widetilde{\sigma_{2i,2i}}$ is irreducible and

$$[\bar{\iota}\iota]=\bigoplus_{i=0}^{n/2-1}[\widetilde{\sigma_{2i,2i}}].$$

Since $(\mathrm{id}_{\mathcal{M}},\sigma_{2i,2i})=\mathbb{C}r_{\lambda_{2i}}$, we get

$$\mathrm{Ang}(P,Q)=\{\cos^{-1}\sqrt{r^*_{\lambda_{2i}}\phi_{\lambda_1^2}(\varepsilon_{\sigma_{2i,2i}}(\lambda_1^2)r_{\lambda_{2i}})};\ i=0,1,\cdots,\frac{n-2}{2}\}\setminus\{0,\frac{\pi}{2}\}.$$

It is easy to show $r^*_{\lambda_{2i}}\phi_{\lambda_1^2}(\varepsilon_{\sigma_{2i,2i}}(\lambda_1^2)r_{\lambda_{2i}})=\phi_{\lambda_1}(\varepsilon(\lambda_{2i},\lambda_1)\varepsilon(\lambda_1,\lambda_{2i}))^2$ and we get the statement.

Assume now that $n=2s+1$ is odd. Then $\widetilde{\sigma_{i,i}}$ is irreducible for $i=0,1,2,\cdots s-1$ while $\widetilde{\sigma_{s,s}}$ is decomposed into two sectors, say μ_0 and μ_1 such that

$$[\bar{\iota}\iota]=\bigoplus_{i=0}^{s-1}[\widetilde{\sigma_{i,i}}]\oplus[\mu_0].$$

Then a similar computation as above shows

$$\mathrm{Ang}(P,Q)=\{\cos^{-1}|\phi_{\lambda_1}(\varepsilon(\lambda_i,\lambda_1)\varepsilon(\lambda_1,\lambda_i))|;\ i=1,2,\cdots,s\},$$

(since $E_PE_Q((\iota,\iota\widetilde{\sigma_{s,s}}))=0$ we have $E_PE_Q((\iota,\iota\mu_0))=0$ too) which shows the statement. □

8. Appendix

In the proof of Theorem 5.19 we use the fact that there exists a unique Q-system for $\mathrm{id}_P\oplus\bar{\kappa}\theta\kappa$ up to equivalence. We give a proof of this statement here.

The next lemma follows from Frobenius reciprocity [20, Section 2.3], which says that the right normalization of the norm of an element $v\in(\tau,\rho\sigma)$ is $\left(\frac{d(\rho)d(\sigma)}{d(\tau)}\right)^{1/4}$.

Lemma 8.1. *Let $\mathcal{L}$, $\mathcal{M}$, and $\mathcal{N}$ be properly infinite factors and $\rho\in\mathrm{Mor}_0(\mathcal{N},\mathcal{M})$, $\sigma\in\mathrm{Mor}_0(\mathcal{L},\mathcal{N})$, and $\tau\in(\mathcal{L},\mathcal{M})$ be irreducible morphisms. If $v\in(\tau,\rho\sigma)$ satisfies*

$||v|| = \big(\frac{d(\rho)d(\sigma)}{d(\tau)}\big)^{1/4}$, *then*



$$= \Big(\frac{d(\tau)d(\sigma)}{d(\rho)}\Big)^{1/4},$$



$$= \Big(\frac{d(\tau)d(\rho)}{d(\sigma)}\Big)^{1/4}.$$

We recall the construction of the Haagerup subfactor in [23, Section 7]. Let $\mathcal{O}_4$ be the Cuntz algebra generated by isometries $\{S_0, T_0, T_1, T_2\}$ and let $d = (3+\sqrt{13})/2$ (which will be $d(\rho)$). We introduce an endomorphism ρ and a period 3 automorphism α of $\mathcal{O}_4$ by setting

$$\alpha(S_0) = S_0, \quad \alpha(T_i) = T_{i+2}, \tag{8.1}$$

$$\rho(S_0) = \frac{1}{d}S_0 + \frac{1}{\sqrt{d}}\sum_{i\in\mathbb{Z}/3\mathbb{Z}} T_iT_i, \tag{8.2}$$

$$\rho(T_i) = \frac{1}{\sqrt{d}}S_0T_{-i}^* + T_{-i}S_0S_0^* + \sum_{j,k\in\mathbb{Z}/3\mathbb{Z}} A(i+j,i+k)T_jT_{i+j+k}T_k^*, \tag{8.3}$$

where i is understood as an element of $\mathbb{Z}/3\mathbb{Z}$ and

$$A(0,0) = 1 - \frac{1}{d-1},$$

$$A(0,1) = A(0,2) = A(1,0) = A(2,0) = A(1,1) = A(2,2) = \frac{-1}{d-1},$$

$$A(1,2) = \overline{A(2,1)} = \frac{1+\sqrt{4d-1}\sqrt{-1}}{2(d-1)}.$$

Then ρ and α extend to an irreducible endomorphism and an outer automorphism respectively of the weak closure M of $\mathcal{O}_4$ in the GNS representation of some KMS state, which will be denoted by the same symbols. Moreover we have $S_0 \in (\mathrm{id}_M, \rho^2)$, $T_0 \in (\rho, \rho^2)$, $T_1 \in (\alpha\rho, \rho^2)$, and $T_2 \in (\alpha^2\rho, \rho^2)$ and $\alpha\rho = \rho\alpha^2$. It is easy to show that ρ satisfies the condition of Lemma 3.5, namely

$$\rho(T_0)S_0 = T_0S_0, \tag{8.4}$$

$$\sqrt{d}S_0 + (d-1)T_0^2 = \sqrt{d}\rho(S_0) + (d-1)\rho(T_0)T_0. \tag{8.5}$$

Therefore there exist a Q-system for $\mathrm{id}_M \oplus \rho$ and a subfactor $P \subset M$ such that $[\kappa\bar{\kappa}] = [\mathrm{id}_M] \oplus [\rho]$, where $\kappa : P \hookrightarrow M$ is the inclusion map. Note that $\rho = \hat{\eta}$

and $\alpha = \theta$ in the notation of Theorem 5.19. Lemma 3.5 shows that we can choose isometries $v = r_{\bar{\kappa}}$, $w = \kappa(r_\kappa)$, and $v_1 \in (\rho, \kappa\bar{\kappa})$ such that (3.3) holds:

$$w = \frac{v}{\sqrt{d+1}} + \frac{v_1\rho(v)v_1^*}{\sqrt{d+1}} + \sqrt{\frac{d}{d+1}}v_1\rho(v_1)S_0v^* + \sqrt{\frac{d-1}{d+1}}v_1\rho(v_1)T_0v_1^*. \tag{8.6}$$

To finish the proof of Theorem 5.19, it suffices to show that there exists a unique Q-system for $\mathrm{id}_P \oplus \bar{\kappa}\alpha\kappa$ up to equivalence. For this purpose we solve the equations (3.1) and (3.2) with $\sigma := \bar{\kappa}\alpha\kappa$. For this, we use the following notation:



$$:= \left(\frac{d(\kappa)^2}{d}\right)^{1/4} v_1^* = \left(\frac{d+1}{d}\right)^{1/4} v_1^*,$$



$$:= \left(\frac{d+1}{d}\right)^{1/4} v_1,$$



$:=$



,



$:=$



,



$:=$



,



$:=$



.

Every vertex expressing an element in $(\alpha\rho, \rho\alpha^2)$, $(\rho, \alpha\rho\alpha)$ etc. such as



,



will always mean 1. We choose S_0 as $r_\rho = \bar{r_\rho}$ and set



$$:= d^{1/4}T_0,$$



$$= d^{1/4}T_0^*.$$

Then



$$= d^{1/2}S_0^*\rho(d^{1/4}T_0) = d^{1/4}T_0^* =$$



,



$$= \rho(d^{1/2}S_0^*)d^{1/4}T_0 = d^{1/4}T_0^* =$$



.

In the same way, we have



Equation (8.4) means



We use a similar expression for $d^{1/4}T_1 \in (\alpha\rho, \rho^2) = (\rho\alpha^2, \rho^2)$ and $d^{1/4}T_2 \in (\alpha^2\rho, \rho^2) = (\rho\alpha, \rho^2)$.

Since we have



we simply express this intertwiner by  . In a similar way, we have



and we simply express it by  . The diagram  is also interpreted in the same way.

We set

$R =$  $\in (\mathrm{id}_P, \sigma^2).$

Then thanks to Lemma 8.1, we have

$$||R|| = \big(d(\kappa)^2\frac{d(\kappa)d(\rho)}{d(\kappa)}\frac{d(\kappa)^2}{d(\rho)}\big)^{1/4} = d(\kappa) = \sqrt{d+1}.$$

All we have to show is the following:

Theorem 8.2. *Let the notation be as above. Then there exist exactly two elements* $S \in (\sigma, \sigma^2)$ *satisfying* $||S|| = d(\sigma)^{1/4} = (d+1)^{1/4}$ *and*

$$\sigma(S)R = SR, \tag{8.7}$$

$$\frac{\sqrt{d+1}}{d}(R-\sigma(R)) = \sigma(S)S - S^2. \tag{8.8}$$

(Note that if S satisfies the above condition, so does $-S$.)

To prove the theorem, we choose a basis of (σ,σ^2). Note that we have

$$\begin{aligned}\dim(\sigma,\sigma^2) &= \dim(\bar{\kappa}\alpha\kappa, \bar{\kappa}\alpha\kappa\bar{\kappa}\alpha\bar{\kappa}) = \dim(\kappa\bar{\kappa}\alpha\kappa\bar{\kappa}, \alpha\kappa\bar{\kappa}\alpha) \\ &= \dim(\alpha\oplus\alpha\rho\oplus\alpha^2\rho\oplus\alpha^2\rho^2, \alpha^2\oplus\rho) = 2.\end{aligned}$$

We set

$$S_1 =$$



$$\in (\sigma,\sigma^2),$$

$$S_2 =$$



$$\in (\sigma,\sigma^2).$$

Then since $r_{\bar{\kappa}}$ and v_1 are orthogonal, so are S_1 and S_2.

Lemma 8.3. *Let the notation be as above. Then $||S_1|| = ||S_2|| = d(\sigma)^{1/4} = (d+1)^{1/4}$.*

Proof. For S_1, we have

$$||S_1||^2 = d(\kappa)$$



$$.$$

Since this is already a scalar, it is equal to



$$=$$



$$= \frac{d(\kappa)}{\sqrt{d}}$$



$$= d(\kappa).$$

For S_2,

$$||S_2||^2 = \frac{d(\kappa)}{\sqrt{d}}$$



$$= \frac{1}{\sqrt{d}}$$



$$= \frac{d(\kappa)}{d}$$



$$= d(\kappa).$$

□

The above lemma shows that every element $S \in (\sigma, \sigma^2)$ satisfying $||S|| = d(\sigma)^{1/4}$ is uniquely expressed as $S = aS_1 + bS_2$, where a and b are complex numbers satisfying $|a|^2 + |b|^2 = 1$. Therefore (8.7) and (8.8) are equivalent to the following equations respectively:

$$aS_1R + bS_2R = a\sigma(S_1)R + b\sigma(S_2)R,$$

$$\begin{aligned}\frac{\sqrt{d+1}}{d}(R - \sigma(R)) &= a^2(\sigma(S_1)S_1 - S_1^2) + ab(\sigma(S_1)S_2 - S_1S_2 + \sigma(S_2)S_1 - S_2S_1) \\ &+ b^2(\sigma(S_2)S_2 - S_2^2).\end{aligned}$$

The following lemma will be frequently used in what follows.

Lemma 8.4. *Let the notation be as above. Then*



Proof. The first equality is easy. For the second, we have



Thanks to Equation (8.6), we have



$$= \left(\frac{d(\kappa)^2}{d}\right)^{3/4} d(\kappa)^{1/2} v_1^*\rho(v_1^*)wv_1 = \frac{\sqrt{d^2-1}}{d} d^{1/4}T_0,$$

which shows the statement. □

Lemma 8.5. *Let the notation be as above. Then we have*

$$S_1R = \sigma(S_1)R, \quad S_2R = \sigma(S_2)R.$$

In particular, (8.7) always holds.

Proof. We introduce a linear isomorphism $F : (\mathrm{id}_P, \sigma^3) \to (\kappa\bar{\kappa}, \alpha\kappa\bar{\kappa}\alpha\kappa\bar{\kappa}\alpha)$ by

$$F(x) =$$



.

Since every intertwiner above belongs to the space $(\mathrm{id}_P, \sigma^3)$, it suffices to show the equalities after applying F to the intertwiners.

For $F(S_1R)$, we have

$$F(S_1R) =$$



$$=$$



.

Using Lemma 8.4 with consideration of $[\alpha^2(\mathrm{id}_M \oplus \rho)\alpha] = [\mathrm{id}_M] \oplus [\alpha\rho]$, we get

$$F(S_1R) = \frac{1}{d}$$



.

In a similar way, we have

$$F(\sigma(S_1)R) = \frac{1}{d}$$



.

Since we have



$$= d\alpha^2(S_0^*)S_0 = d,$$



$$= d\alpha(S_0^*)S_0 = d,$$

we get $F(S_1R) = F(\sigma(S_1)R)$.

For $F(S_2R)$, we have

$F(S_2R) =$ 

$=$ 

.

Using Lemma 8.4 with consideration of $[\rho\ \mathrm{id}_M\alpha] = [\alpha\rho]$ and $[\rho\rho\alpha] = [\alpha] \oplus [\rho] \oplus [\alpha\rho] \oplus [\alpha^2\rho]$, we get

$F(S_2R) = \dfrac{d-1}{d}$ 

.

In a similar way,

$F(\sigma(S_2)R) = \dfrac{d-1}{d}$ 

.

Since we have



$$= d^{3/4}\rho(T_0^*)T_2T_0 = d^{3/4}\overline{A(2,1)}T_1 = d^{3/4}A(1,2)T_1,$$



$$= d^{3/4}\alpha(T_0^*)\alpha\rho(T_1)T_0 = d^{3/4}T_2^*\rho(T_2)T_0 = d^{3/4}A(1,2)T_1,$$

we get $F(S_2R) = F(\sigma(S_2)R)$. ☐

Lemma 8.6. *Let $S = aS_1 + bS_2$ with $a, b \in \mathbb{C}$. Then Equation (8.8) is equivalent to*

$$\frac{1}{\sqrt{d}} = a^2 + \sqrt{d-1}ab, \tag{8.9}$$

$$\sqrt{\frac{d-1}{d}} = ab - \frac{b^2}{\sqrt{d-1}} \tag{8.10}$$

$$0 = a^2 - \frac{ab}{\sqrt{d-1}} - A(1,2)b^2, \tag{8.11}$$

$$0 = ab + \frac{1+(d-1)A(1,2)}{(d-1)^{3/2}}b^2. \tag{8.12}$$

Proof. Let $G : (\sigma, \sigma^3) \to (\kappa\bar{\kappa}\alpha\kappa\bar{\kappa}, \alpha\kappa\bar{\kappa}\alpha\kappa\bar{\kappa}\alpha)$ be the linear isomorphism defined in a similar way as in the proof of the previous lemma. We first compute $G(R)$, $G(\sigma(R))$, $G(S_1^2)$ etc.

For $G(R)$, we have

$$G(R) =$$



$$= \frac{1}{d(\kappa)}$$



$$+ \frac{\sqrt{d}}{d(\kappa)}$$



.

Applying Lemma 8.4 to the second term, we get

$$G(R) = \frac{1}{\sqrt{d+1}}$$



$$+ \frac{1}{\sqrt{d+1}}$$



$$+ \sqrt{\frac{d-1}{d+1}}$$



.

The intertwiner $G(\sigma(R))$ is the mirror image of $G(R)$. Since



$$= \sqrt{d}S_0 =$$



,

we get

$$\frac{\sqrt{d+1}}{d}G(R-\sigma(R))$$

$$= \frac{1}{d}$$



$$-\frac{1}{d}$$



$$+ \frac{\sqrt{d-1}}{d}$$



$$-\frac{\sqrt{d-1}}{d}$$



.

For $G(S_1^2)$, Lemma 8.4 implies

$$G(S_1^2) =$$



$$=$$



$$= \frac{1}{\sqrt{d}}$$



$$+\sqrt{\frac{d-1}{d}}$$



$$= \frac{1}{\sqrt{d}}$$



$$+\sqrt{\frac{d-1}{d}}$$



,

where we use $\alpha(S_0) = S_0$. The intertwiner $G(\sigma(S_1)S_1)$ is the mirror image of $G(S_1^2)$. A similar argument shows

$$G(S_2S_1) = \frac{1}{\sqrt{d}}$$


κ $\bar\kappa$ α κ $\bar\kappa$
ρ
ρ
α^2
ρ
α
ρ
ρ
α κ $\bar\kappa$ α κ $\bar\kappa$ α

$$+\sqrt{\frac{d-1}{d}}$$


κ $\bar\kappa$ α κ $\bar\kappa$
ρ
ρ
ρ
ρ
α^2
ρ
α
ρ
ρ
α κ $\bar\kappa$ α κ $\bar\kappa$ α

,

and $G(\sigma(S_2)S_1)$ is its mirror image.

For $G(S_1S_2)$, we have

$$G(S_1S_2) =$$


κ $\bar\kappa$ α κ $\bar\kappa$
ρ
ρ
ρ
ρ
$\bar\kappa$
α
κ
ρ
ρ
α^2
α κ $\bar\kappa$ α κ $\bar\kappa$ α

$$=$$


κ $\bar\kappa$ α κ $\bar\kappa$
$\bar\kappa$
ρ
ρ
ρ
ρ
α
α^2
ρ
ρ
κ
α κ $\bar\kappa$ α κ $\bar\kappa$ α

$$= \frac{1}{d}$$


κ $\bar\kappa$ α κ $\bar\kappa$
ρ
ρ
ρ
α
α^2
ρ
α κ $\bar\kappa$ α κ $\bar\kappa$ α

$$+\frac{\sqrt{d-1}}{d}$$


κ $\bar\kappa$ α κ $\bar\kappa$
ρ
ρ
ρ
α
α^2
ρ
ρ
ρ
α κ $\bar\kappa$ α κ $\bar\kappa$ α

$$+\frac{\sqrt{d-1}}{d}$$


κ $\bar\kappa$ α κ $\bar\kappa$
ρ
ρ
ρ
ρ
ρ
α
α^2
ρ
α κ $\bar\kappa$ α κ $\bar\kappa$ α

$$+\frac{d-1}{d}$$


κ $\bar\kappa$ α κ $\bar\kappa$
ρ
ρ
ρ
ρ
ρ
α
α^2
ρ
ρ
ρ
α κ $\bar\kappa$ α κ $\bar\kappa$ α

.

Since $(\alpha^2\alpha, \alpha\rho) = (\alpha^2\alpha, \rho\alpha\rho) = 0$, we get

$$G(S_1S_2) = \frac{\sqrt{d-1}}{d}$$



$$+ \frac{d-1}{d}$$



.

The intertwiner $G(\sigma(S_1)S_2)$ is the mirror image of $G(S_1S_2)$. A similar argument shows

$$G(S_2^2) = \frac{\sqrt{d-1}}{d}$$



$$+ \frac{\sqrt{d-1}}{d}$$



$$+ \frac{d-1}{d}$$



and $G(\sigma(S_2)S_2)$ is the mirror image of $G(S_2^2)$.

Now (8.8) is equivalent to

$$\frac{1}{d}$$



$$= \frac{a^2}{\sqrt{d}}$$



$$+ \frac{\sqrt{d-1}}{d}ab$$



,

$$\frac{1}{d}$$



$$= \frac{a^2}{\sqrt{d}}$$



$$+ \frac{\sqrt{d-1}}{d}ab$$



,

$$\frac{\sqrt{d-1}}{d}$$



$$= \frac{ab}{\sqrt{d}}$$



$$+ \frac{\sqrt{d-1}}{d}b^2$$



,

$$\frac{\sqrt{d-1}}{d}$$



$$= \frac{ab}{\sqrt{d}}$$



$$+ \frac{\sqrt{d-1}}{d} b^2$$



,

$$0 = \sqrt{\frac{d-1}{d}} a^2$$



$$+ \frac{d-1}{d} ab$$



$$- \frac{\sqrt{d-1}}{d} b^2$$



,

$$0 = \sqrt{\frac{d-1}{d}} a^2$$



$$+ \frac{d-1}{d} ab$$



$$- \frac{\sqrt{d-1}}{d} b^2$$



,

$$0 = \sqrt{\frac{d-1}{d}} ab$$



$$- \sqrt{\frac{d-1}{d}} ab$$



$$+ \frac{d-1}{d} b^2$$



$$- \frac{d-1}{d} b^2$$



.

It is easy to show that the first and the second equations are equivalent to (8.9), the third and fourth are equivalent to (8.10), and the fifth and sixth are equivalent to (8.11). For the last equation, we have



$-$



$$= dT_2\rho\alpha^2(S_0^*)T_0 - d\alpha(T_1)S_0^*\rho\alpha(T_0)$$

$$= dT_2\rho(S_0^*)T_0 - dT_0S_0\rho(T_2) = \sqrt{d}(T_2T_0^* - T_0T_1^*),$$



$-$



$$\begin{aligned}
&= d\rho(T_0^*)T_2\rho(T_2^*)T_0 - d\alpha(T_0^*)\alpha\rho(T_1)T_2^*\rho\alpha(T_0)\\
&= d\rho(T_0^*)T_2\rho(T_2^*)T_0 - dT_2^*\rho(T_2)T_2^*\rho(T_2)\\
&= d\Big(\sum_{l\in\mathbb{Z}/3\mathbb{Z}} A(2,l)T_{2+l}T_l^*\Big)^*\Big(\sum_{k\in\mathbb{Z}/3\mathbb{Z}} A(2,2+k)T_{2+k}T_k^*\Big)^*\\
&- d\Big(\sum_{l\in\mathbb{Z}/3\mathbb{Z}} A(1,l+2)T_{1+l}T_l^*\Big)\Big(\sum_{k\in\mathbb{Z}/3\mathbb{Z}} A(1,k+2)T_{1+k}T_k^*\Big)\\
&= d\sum_{l\in\mathbb{Z}/3\mathbb{Z}} \overline{A(2,l)A(2,l+1)}T_lT_{l+1}^* - d\sum_{k\in\mathbb{Z}/3\mathbb{Z}} A(1,k)A(1,k+2)T_{k-1}T_k^*\\
&= d\Big(\frac{1}{(d-1)^2} + \frac{A(1,2)}{d-1}\Big)(T_2T_0^* - T_0T_1^*).
\end{aligned}$$

Therefore the last equation is equivalent to (8.12). □

Proof of Theorem 8.2. Equation (8.11) shows $b \neq 0$ and so (8.12) implies

$$a = -\frac{B+1}{(d-1)^{3/2}}b. \tag{8.13}$$

where $B=(d-1)A(1,2)$. Using the fact that B satisfies $B^2-B+d=0$, we can see that this is compatible with (8.11). Iterating (8.13) into (8.9), we get

$$\begin{aligned}\frac{1}{\sqrt{d}} &= \frac{(B+1)^2}{(d-1)^3}b^2-\frac{B+1}{d-1}b^2=\frac{1}{(d-1)^3}\big(B^2+2B+1-(d-1)^2(B+1)\big)b^2\\ &= \frac{1}{(d-1)^3}\big(3B-d+1-(d+2)(B+1)\big)b^2\\ &= \frac{1}{(d-1)^3}\big((1-d)B+d-d^2\big)b^2\\ &= -\frac{1}{(d-1)^2}\big(B+d)b^2,\end{aligned}$$

which shows

$$b^2=-\frac{(d-1)^2}{(B+d)\sqrt{d}}. \tag{8.14}$$

We get the same equation from (8.10).

Now all we have to show is that (8.13) and (8.14) are compatible with $|a|^2+|b|^2=1$. The equation (8.13) and $|a|^2+|b|^2=1$ imply

$$\frac{1}{|b|^2}=1+\frac{1}{(d-1)^3}|B+1|^2=1+\frac{d+2}{(d-1)^3}=1+\frac{1}{d-1}=\frac{d}{d-1}.$$

Since $|B+d|^2=d^2+2d=d(d-1)^2$, this is compatible with (8.14). □

REFERENCES

1. Asaeda, M. and Haagerup, U., Exotic subfactors of finite depth with Jones indices $(5+\sqrt{13})/2$ and $(5+\sqrt{17})/2$. *Comm. Math. Phys.* **202** (1999), 1–63.
2. Böckenhauer, J., Evans, D. E., and Kawahigashi, Y. Chiral structure of modular invariants for subfactors. *Comm. Math. Phys.* **210** (2000), 733–784.
3. Böckenhauer, J., Evans, D. E., and Kawahigashi, Y. On α-induction, chiral generators and modular invariants for subfactors. *Comm. Math. Phys.* **208** (1999), no. 429–487.
4. Bisch, D. A note on intermediate subfactors. *Pacific Journal of Mathematics*, **163** (1994), 201–216.
5. Bisch, D., Bimodules, higher relative commutants and the fusion algebra associated to a subfactor. In *Operator algebras and their applications.* Fields Institute Communications, Vol. 13, American Math. Soc. (1997), 13–63.
6. Bisch, D. and Haagerup, U., Composition of subfactors: new examples of infinite depth subfactors. *Ann. Sci. École Norm. Sup.* (4) **29** (1996), 329–383.
7. Bisch, D. and Jones, V. F. R., Algebras associated to intermediate subfactors. *Inventiones Mathematicae*, **128** (1997), 89–157.
8. Choda, M. Index for factors generated by Jones' two sided sequence of projections. *Pacific J. Math.* **139** (1989), 1–16.
9. Evans, D. E. and Kawahigashi, Y., Quantum symmetries on operator algebras. *Oxford University Press* (1998).
10. Evans, D. E. and Kawahigashi, Y., Orbifold subfactors from Hecke algebras. II. Quantum doubles and braiding. *Comm. Math. Phys.* **196** (1998), 331–361.
11. Goodman, F., de la Harpe, P. and Jones, V. F. R., Coxeter graphs and towers of algebras. *MSRI Publications (Springer)* (1989), **14**.

12. Grossman, P. (in press), Forked Temperley-Lieb algebras and intermediate subfactors. *J. Funct. Anal.* (2007), doi:10.1016/j.jfa.2007.03.014
13. Grossman, P. and Jones, V. F. R., Intermediate subfactors with no extra structure. *J. Amer. Math. Soc.* **20** (2007), 219–265.
14. Hayashi, T. and Yamagami, S., Amenable tensor categories and their realizations as AFD bimodules. *J. Funct. Anal.* **172** (2000), 19–75.
15. Hiai, F., Minimizing indices of conditional expectations onto a subfactor. *Publ. Res. Inst. Math. Sci.* **24** (1988), 673–678.
16. Hong, J. H., Subfactors with principal graph $E_6^{(1)}$. *Acta Appl. Math.* **40** (1995), 255–264.
17. Izumi, M., Application of fusion rules to classification of subfactors. *Publ. Res. Inst. Math. Sci.*, **27** (1991), 953–994.
18. Izumi, M., Goldman's type theorem for index 3. *Publ. Res. Inst. Math. Sci.* **28** (1992), 833–843.
19. Izumi, M., On type II and type III principal graphs of subfactors. *Math. Scand.* **73** (1993), 307–319.
20. Izumi, M., Subalgebras of infinite C*-algebras with finite Watatani indices II. Cuntz-Krieger algebras. *Duke J. Math.* **91** (1998), 409-461.
21. Izumi, M., Goldman's type theorems in index theory. *Operator algebras and quantum field theory (Rome, 1996)*, 249–269, Int. Press, Cambridge, MA, 1997.
22. Izumi, M., The structure of sectors associated with Longo-Rehren inclusions. I. General theory. *Comm. Math. Phys.* **213** (2000), 127–179.
23. Izumi, M., The structure of sectors associated with Longo-Rehren inclusions. II. Examples. *Rev. Math. Phys.* **13** (2001), 603–674.
24. Izumi, M., Characterization of isomorphic group-subgroup subfactors. *Int. Math. Res. Not.* 2002, no. 34, 1791–1803.
25. Izumi, M. and Kawahigashi, Y., Classification of subfactors with the principal graph $D_n^{(1)}$. *J. Funct. Anal.* **112** (1993), 257–286.
26. Izumi, M. and Kosaki, H., Kac algebras arising from composition of subfactors: general theory and classification. *Mem. Amer. Math. Soc.* **158** (2002), no. 750.
27. Izumi, M. and Kosaki, H., On a subfactor analogue of the second cohomology. *Reviews in Mathematical Physics*, **14** (2002), 733–757.
28. Izumi, M., Longo, R., and Popa, S. A Galois correspondence for compact groups of automorphisms of von Neumann algebras with a generalization to Kac algebras. *J. Funct. Anal.* **155** (1998), 25–63.
29. Jones, V. F. R. (1980). Actions of finite groups on the hyperfinite type II_1 factor. *Memoirs of the American Mathematical Society*, **237**.
30. Jones, V. F. R., Index for subfactors. *Inventiones Mathematicae*, **72** (1983), 1–25.
31. Jones, V. F. R. (in press). Planar algebras I. *New Zealand Journal of Mathematics.* QA/9909027
32. Jones, V. F. R., Quadratic tangles in planar algebras. In preparation, (2003), http://math.berkeley.edu∼vfr
33. Jones, V. F. R. and Xu, F., Intersections of finite families of finite index subfactors. *International Journal of Mathematics*, **15** (2004), 717–733.
34. Kawahigashi, Y., On flatness of Ocneanu's connections on the Dynkin diagrams and classification of subfactors. *J. Funct. Anal.* **127** (1995), 63–107.
35. Kawahigashi, Y. and Longo, R., Classification of two-dimensional local conformal nets with $c < 1$ and 2-cohomology vanishing for tensor categories. *Comm. Math. Phys.* **244** (2004), 63–97.
36. Kawahigashi, Y. and Longo, R., Pennig, U., and Rehren, K.-H., The classification of non-local chiral CFT with $c < 1$. math.OA/0505130, to appear in Comm. Math. Phys.
37. Kirillov, A. N. and Reshetikhin, N. Yu., Representations of the algebra $U_q(\mathrm{sl}(2))$, q-orthogonal polynomials and invariants of links. Infinite-dimensional Lie algebras and groups (Luminy-Marseille, 1988), 285–339, Adv. Ser. Math. Phys., 7, World Sci. Publ., Teaneck, NJ, 1989.

38. Kosaki, H. Extension of Jones' theory on index to arbitrary factors. *J. Funct. Anal.* **66** (1986), 123–140.
39. Kosaki, H. and Yamagami, S., Irreducible bimodules associated with crossed product algebras. *Internat. J. Math.* **3** (1992), 661–676.
40. Landau, Z., Fuss-Catalan algebras and chains of intermediate subfactors. *Pacific J. Math.*, **197** (2001), 325–367.
41. Landau, Z., Exchange relation planar algebras. *J. of Funct. Anal.*, **195** (2002), 71–88.
42. Longo, R., Index of subfactors and statistics of quantum fields. II. Correspondences, braid group statistics and Jones polynomial. *Comm. Math. Phys.* **130** (1990), 285–309.
43. Longo, R., A duality for Hopf algebras and for subfactors. I. *Comm. Math. Phys.* **159** (1994), 133–150.
44. Longo, R. and Rehren, K.-H. (1995). Nets of subfactors. *Reviews in Mathematical Physics*, **7**, 567–597.
45. Longo, R. and Roberts, J. E. (1997). A theory of dimension. *K-theory*, **11**, 103–159.
46. Masuda, T., An analogue of Longo's canonical endomorphism for bimodule theory and its application to asymptotic inclusions. *Internat. J. Math.* **8** (1997), 249–265.
47. Masuda, T., An analogue of Connes-Haagerup approach for classification of subfactors of type III_1. *J. Math. Soc. Japan*, **57** (2005), 959–1001.
48. Nakamura, M. and Takeda, Z., A Galois theory for finite factors. *Proc. Japan Acad.* **36** (1960), 258–260.
49. Ocneanu, A. *Quantum symmetry, differential geometry of finite graphs and classification of subfactors*, University of Tokyo Seminary Notes 45, (1991), (Notes recorded by Kawahigashi, Y.).
50. Okamoto, S., Invariants for subfactors arising from Coxeter graphs. *Current Topics in Operator Algebras*, World Scientific Publishing, (1991), 84–103.
51. Pimsner, M. and Popa, S., Entropy and index for subfactors. *Annales Scientifiques de l'École Normale Superieur*, **19** (1986), 57–106.
52. Popa, S., Classification of amenable subfactors of type II. *Acta Math.* **172** (1994), 163–255.
53. Popa, S., Classification of subfactors and their endomorphisms. CBMS Regional Conference Series in Mathematics, 86. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1995.
54. Sano, T. and Watatani, Y., Angles between two subfactors. *Journal of Operator Theory*, **32** (1994), 209–241.
55. Wassermann, A., Operator algebras and conformal field theory. III. Fusion of positive energy representations of $\mathrm{LSU}(N)$ using bounded operators. *Invent. Math.* **133** (1998), 467–538.
56. Watatani, Y., Lattices of intermediate subfactors. *J. Funct. Anal.*, **140** (1996), 312–334.
57. Wenzl, H., C^* tensor categories from quantum groups. *J. Amer. Math. Soc.* **11** (1998), 261–282.
58. Xu, Feng, New braided endomorphisms from conformal inclusions. *Comm. Math. Phys.* **192** (1998), 349–403.

Pinhas Grossman, Department of Mathematics, 1326 Stevenson Center, Vanderbilt
University, Nashville, TN, 37203
e-mail: pinhas.grossman@vanderbilt.edu
Masaki Izumi, Department of Mathematics, Graduate School of Science, Kyoto University,
Sakyo-ku, Kyoto 606-8502, Japan
e-mail: izumi@math.kyoto-u.ac.jp